John Kalliongis, Ryo Ohashi

Finite actions on the Klein four-orbifold and prism manifolds

Comment.Math.Univ.Carolin. 58,1 (2017) 49-68.

Abstract: We describe the finite group actions, up to equivalence, which can act on the orbifold $\Sigma(2,2,2)$, and their quotient types. This is then used to consider actions on prism manifolds M(b,d) which preserve a longitudinal fibering, but do not leave any Heegaard Klein bottle invariant. If $\varphi: G \to \text{Homeo}(M(b,d))$ is such an action, we show that M(b,d) = M(b,2) and $M(b,2)/\varphi$ fibers over a certain collection of 2-orbifolds with positive Euler characteristic which are covered by $\Sigma(2,2,2)$. For the standard actions, we compute the fundamental group of $M(b,2)/\varphi$ and indicate when it is a Seifert fibered manifold.

Keywords: finite group action; prism 3-manifold; equivalence of actions; orbifold; Klein four-group

AMS Subject Classification: Primary 57M99; Secondary 57S99

References

- [1] Dummit D., Foote R., Abstract Algebra, Wiley, Hoboken, NJ, 2004.
- [2] Kalliongis J., Ohashi R., Finite group actions on prism manifolds which preserve a Heegaard Klein bottle, Kobe J. Math. 28 (2011), no. 1, 69–89.
- [3] Kalliongis J., Ohashi R., Classifying non-splitting fiber preserving actions on prism manifolds, Topology Appl. 178 (2014), 200–218.
- [4] Kalliongis J., Ohashi R., Finite actions on the 2-sphere and the projective plane, preprint.
- [5] McCullough D., Isometries of elliptic 3-manifolds, J. London Math. Soc. 65 (2002), no. 1, 167–182.
- [6] Orlik P., Seifert Manifolds, Lecture Notes in Mathematics, 291, Springer, Berlin-New York, 1972.
- [7] Scott P., The geometries of 3-manifolds, Bull. London Math. Soc. 5 (1983), 401-48.