

# Groups, transversals, and loops

TUVAL FOGUEL

*Abstract.* A family of loops is studied, which arises with its binary operation in a natural way from some transversals possessing a “normality condition”.

*Keywords:* loops, groups, transversals

*Classification:* Primary 20N05; Secondary 20D99

## §1. Introduction

The study of loops leads in a natural way to the study of transversals of subgroups, for example the works of: Karzel, Kepka, Kiechle, Kinyon, Niemenmaa, Phillips, Sabinin, Ungar, and et al. [6], [10], [11], [13], [15], [16]. Loops have also played an important role in the study of groups such as in Conway’s celebrated construction of the Fischer-Griess monster group using Parker’s Moufang loop of order  $2^{13}$ . This led to Griess’s construction of binary code loops by a double cover elementary abelian 2-group ([3], [8]). From the above one can see that there is a strong connection between the study of group and loops.

Sabinin has shown that every left loop arises with its binary operation in a natural way from some special transversal of a subgroup in certain groups ([16]). In this paper I will look at a family of loops which arises with its binary operation in a natural way from some transversals which possesses a “normality condition”. Subgroups and subsets of groups with normality conditions such as subnormality, seminormality, and etc. have interested me for a long time ([4], [5]).

## §2. AP-loop as transversals of groups

**Definition 2.1.** A *groupoid* ([1]) is a nonempty set with a binary operation. An automorphism of the groupoid  $(S, \odot)$  is a bijection of  $S$  that respects the binary operation  $\odot$  in  $S$ . The set of all automorphisms of  $(S, \odot)$  forms a group denoted by  $Aut(S, \odot)$ .

**Definition 2.2.** A *left loop* is a groupoid  $(S, \odot)$  with an identity element in which the equation  $a \odot x = b$  possesses a unique solution for the unknown  $x$ .  $(S, \odot)$  is a *loop* if  $y \odot a = b$  also possesses a unique solution.

**Definition 2.3.** Given a left loop  $P$  and an ordered pair  $(a, b) \in P \times P$  we get a bijection  $\delta_{a,b} : P \rightarrow P$  defined by  $a \odot (b \odot x) = (a \odot b) \odot \delta_{a,b}x$  for any  $x \in P$ . Here  $\delta_{a,b}$  is a correction to associativity called a *left inner mapping* ([1]). Let  $AS_\delta(P)$  be the group generated by all the bijections  $\delta_{a,b}$  (Note that  $(a \odot b) \odot x = a \odot (b \odot \delta_{a,b}^{-1}x)$ ).

*Note.* It is shown in [16] that if  $L_a$  for  $a \in P$  is the left translation by  $a$ , then  $\delta_{a,b} = L_{(a \odot b)}^{-1} L_a L_b$  where the product is in the permutation group  $L_P$  generated by all left translations.

**Definition 2.4.** We will call a left loop (or a loop) a *left  $A_l$ -loop* (or  *$A_l$ -loop* respectively) if for all  $(a, b) \in P \times P$ ,  $\delta_{a,b} \in \text{Aut}(P, \odot)$ .

**Definition 2.5.** A groupoid  $(P, \odot)$  has the *left inverse property* if for each  $a \in P$ , there is a unique  $a^{-1} \in P$  such that  $a^{-1} \odot (a \odot b) = b$  for all  $b$  in  $P$ .

**Definition 2.6.** A left  $A_l$ -loop  $P$  (or an  $A_l$ -loop) is a *left AP-loop* (or *AP-loop* respectively) if it possesses the left inverse property.

Definitions of two well known AP-loops are presented below:

**Definition 2.7** ( *$A_l$ -Bol-loop = Gyrogroup* [15], [17]). A groupoid  $(G, \odot)$  is an  *$A_l$ -Bol-loop* if its binary operation satisfies the following axioms. In  $G$  there is at least one element, 1, called a left identity, satisfying

(G1)  $1 \odot a = a$  Left Identity  
 for all  $a \in G$ . There is an element  $1 \in G$  satisfying axiom (G1) such that for each  $a$  in  $G$  there is an  $x$  in  $G$ , called a left inverse of  $a$ , satisfying

(G2)  $x \odot a = 1$ . Left Inverse

Moreover, for any  $a, b, z \in G$  there exists a unique element  $\delta_{a,b}z \in G$  such that

(G3)  $a \odot (b \odot z) = (a \odot b) \odot \delta_{a,b}z$ .

If  $\delta_{a,b}$  denotes the map  $\delta_{a,b}: G \rightarrow G$  given by  $z \mapsto \delta_{a,b}z$  then

(G4)  $\delta_{a,b} \in \text{Aut}(G, \odot)$ ,

(G5)  $\delta_{a,b} = \delta_{a \odot b, b}$ . Left Loop Property

**Definition 2.8** (*Gyrocommutative Gyrogroup = K-loop = Bruck-loop* [9], [15], [17]). The  $A_l$ -Bol-loop  $(G, \oplus)$  is a *Bruck loop* if for all  $a, b \in G$ ,

(G6)  $a \odot b = \delta_{a,b}(b \odot a)$ .

In case  $P$  is a left  $A_l$ -loop, since  $\delta_{a,b} \in \text{Aut}(P, \odot)$ , Sabinin’s [16] “semidirect product” becomes:

**Definition 2.9.** Let  $P = (P, \odot)$  be a left  $A_l$ -loop, and let  $AS_\delta(P) \leq H \leq \text{Aut}(P, \odot)$ . The *semidirect product group*

$$P \rtimes_1 H$$

is the set of ordered pairs  $(x, X)$ , where  $x \in P$  and  $X \in H$ , with the binary operation given by

$$(x, X)(y, Y) = (x \odot Xy, \delta_{x, Xy}XY).$$

Below is a corollary to Sabinin’s Theorem 2 [16] about “semidirect products”:

**Corollary 2.10.** *Let  $(P, \odot)$  be a left  $A_l$ -loop, and let  $AS_\delta(P) \leq H \leq \text{Aut}(P, \odot)$ . Then  $P \rtimes_1 H$  is a group.*

**Definition 2.11.** A set  $B$  is a transversal in a group  $G$  (all transversals in this article are left transversals) of a subgroup  $H$  of  $G$  if every  $g \in G$  can be written uniquely as  $g = bh$  where  $b \in B$  and  $h \in H$ . Let  $b_1, b_2 \in B$  be any two elements of  $B$ , and let

$$b_1b_2 = (b_1 \odot b_2)h(b_1, b_2)$$

be the unique decomposition of the element  $b_1b_2 \in G$ , where  $b_1 \odot b_2 \in B$  and  $h(b_1, b_2) \in H$ , determining (i) a binary operation,  $\odot$ , in  $B$ , called the *loop or transversal operation* of  $B$  induced by  $G$ , and (ii) a map  $h: B \times B \rightarrow H$ , called the *transversal map*. The element  $h(b_1, b_2) \in H$  is called the element of  $H$  determined by the two elements  $b_1$  and  $b_2$  of its transversal  $B$  in  $G$ . A *transversal groupoid*  $(B, \odot)$  of  $H$  in  $G$  is a groupoid formed by a transversal  $B$  of  $H$  in  $G$  with its transversal operation  $\odot$ .

**Definition 2.12.** A transversal groupoid  $(B, \odot)$  of a subgroup  $H$  in a group  $G$  is an  *$A_l$ -transversal* of  $H$  in  $G$  if

- (i)  $1_G \in B$ ,  $1_G$  being the identity element of  $G$ ;
- (ii)  $B$  is normalized by  $H$ ,  $H \subseteq N_G(B)$ , that is,  $hBh^{-1} \subseteq B$  for all  $h \in H$ .

*Note.* If an  $A_l$ -transversal is also a subgroup, then it is a normal subgroup. So we see that an  $A_l$ -transversal possesses a “normality condition”.

**Theorem 2.13.** *Let  $(B, \odot)$  be an  $A_l$ -transversal groupoid of a subgroup  $H$  in a group  $G$ . Then, for any  $a, b, x \in B$ ,  $(a \odot b) \odot \delta_{a,b}x = a \odot (b \odot x)$  and  $\delta_{a,b} \in \text{Aut}(B, \odot)$ .*

PROOF: Let  $a, b \in B$  be any two elements of  $B$ , and let  $ab = (a \odot b)h(a, b)$  be the unique decomposition of the element  $ab \in G$ , where  $a \odot b \in B$  and  $h(a, b) \in H$ . Let  $\delta_{a,b}x = x^{h(a,b)} = h(a, b)x(h(a, b))^{-1}$  for all  $x \in B$ .

For all  $a, b, c \in B$  we have in  $G$ ,

$$(2.1) \quad (ab)c = a(bc).$$

Employing the uniqueness of the decomposition for both sides of (2.1) we have

$$(2.2) \quad \begin{aligned} (ab)c &= (a \odot b)h(a, b)c \\ &= (a \odot b)\delta_{a,b}ch(a, b) \\ &= ((a \odot b) \odot \delta_{a,b}c)h(a \odot b, \delta_{a,b}c)h(a, b) \end{aligned}$$

on one hand, and

$$(2.3) \quad \begin{aligned} a(bc) &= a(b \odot c)h(b, c) \\ &= (a \odot (b \odot c))h(a, b \odot c)h(b, c) \end{aligned}$$

on the other hand. It follows from (2.1)–(2.3) and from the uniqueness of the decomposition that

$$(a \odot b) \odot \delta_{a,b}c = a \odot (b \odot c).$$

We now have to show that

$$(x \odot y)^{h(a,b)} = x^{h(a,b)} \odot y^{h(a,b)}$$

for all  $a, b, x, y \in B$ .

More generally, however, we will verify the desired identity for any  $k \in H$  regardless of whether or not  $k$  possesses the form  $k = h(a, b)$ . We will thus show that

$$(x \odot y)^k = x^k \odot y^k$$

for any  $k \in H$ . Clearly, we have in  $G$

$$(xy)^k = x^k y^k.$$

Employing the unique decomposition  $G = BH$ , we have

$$(xy)^k = ((x \odot y)h(x, y))^k = (x \odot y)^k h(x, y)^k$$

on one hand, and

$$x^k y^k = (x^k \odot y^k)h(x^k, y^k)$$

on the other hand. It follows from the above and from the uniqueness of the decomposition  $G = BH$  that

$$(x \odot y)^k = x^k \odot y^k,$$

which completes the proof. □

**Theorem 2.14.** *An  $A_l$ -transversal  $B$  of a subgroup  $H$  in  $G$  that possesses the left inverse property is a left AP-loop under the loop operation.*

PROOF: We have to show that  $(B, \odot)$  satisfies axioms (G1)–(G4) of Definition 2.7 and is a left loop. Axioms (G3) and (G4) are verified in Theorem 2.13, and we get (G2) from the left inverse property.

Given  $b \in B$  we get

$$b = 1b = (1 \odot b)h(1, b)$$

Hence (G1) is verified. Given  $a \odot x = b$  we have a unique solution  $a^{-1} \odot b$ . □

*Note.* If  $B$  is an  $A_l$ -transversal with  $B = B^{-1}$ , then  $B$  is an left AP-loop.

**Definition 2.15.** An  $A_l$ -transversal  $B$  with  $B = B^{-1}$  is called an *AP-transversal* = *gyrotransversal* ([6]).

**Example 2.16.** Let  $P \subset S_n$  where  $P$  consists of the identity permutation and all 2-cycles of the form  $(1, i)$  for  $i = 2, \dots, n$ . Let  $H$  be the stabilizer of 1 in  $S_n$ . Then  $P$  is a transversal of  $H$ , and is, in fact, an *AP-transversal*, but  $P$  is not a loop.

*Note.*  $P = P^{-1}$  and if  $a \in P$  and  $h \in H$ , then  $hah^{-1} \in P$ .

### §3. A family of AP-loops

In the literature there are many examples of  $A_l$ -Bol-loops and  $K$ -loops which are all necessarily *AP-loops*. In this section we will look at a family of *AP-loops* that are not  $A_l$ -Bol-loops (and thus not  $K$ -loops).

**Definition 3.1** (Diagonal transversals). Let  $K$  be a group and let  $G = K \rtimes_l Inn(K)$  be the semidirect product group of  $K$  and  $Inn(K)$ , where  $Inn(K)$  is the inner automorphism group of  $K$  whose generic element  $\alpha_k$  denotes conjugation by  $k \in K$  (i.e.  $\alpha_k x = kxk^{-1}$ ). Then, the *diagonal transversal*  $D$  generated by  $K$  (in  $G$ ) is the subset of  $G$  given by

$$D = \{(k, \alpha_k) | k \in K\} \subset G$$

which is a transversal of  $Inn(K)$  in  $G$ . Any element  $(k, \alpha_k) \in D$  is determined by a corresponding element  $k \in K$ . We therefore use the notation

$$D(k) = (k, \alpha_k)$$

to denote the elements of  $D$ .

**Theorem 3.2.** *A diagonal transversal with its transversal operation is an AP-transversal.*

PROOF: ([6, Theorem 3.2]). □

**Definition 3.3** (Associated Left Gyrogroups) [6]. The associated left gyrogroup of a group  $(K, \cdot)$  from Theorem 3.2 is the left  $A_l$ -loop  $(K, \odot)$ . The operation  $\odot$  is given in terms of the group operation  $\cdot$  by  $a \odot b = ab^a = a^2ba^{-1}$  for all  $a, b \in K$ . This corresponds to the transversal operation of the diagonal transversal.

**Theorem 3.4.** *If  $K$  is a nilpotent group, then  $(K, \odot)$  is an AP-loop.*

PROOF: It will suffice to show that the equation  $x \odot a = b$  has a unique solution for  $x$ . It is true when  $(K, \cdot)$  is abelian, and an induction on the class (divide out the center) does the rest. □

**Theorem 3.5.** *The associated left gyrogroup  $(K, \odot)$  of a group  $(K, \cdot)$  is a group if and only if  $(K, \cdot)$  is nilpotent of class 2.*

PROOF: ([6, Theorem 3.6]). □

**Definition 3.6.** Given  $a, b \in E$  let  $[a, {}_1 b] = [a, b]$  and inductively  $[a, {}_{n+1} b] = [[a, {}_n b], b]$ . A group  $E$  is  $n$ -Engel group ([14])  $[a, {}_n b] = 1$  for all  $a, b \in E$ .

**Theorem 3.7.** Let  $(K, \odot)$  be the associated left gyrogroup of a group  $(K, \cdot)$ . Then  $(K, \odot)$  is a  $A_l$ -Bol-loop = gyrogroup if and only if  $(K, \cdot)$  is central by a 2-Engel group.

PROOF: ([6, Theorem 3.7]). □

Hence, for any nilpotent group  $(K, \cdot)$  of class  $\geq 5$ , the associated groupoid  $(K, \odot)$  is an AP-loop but not  $A_l$ -Bol-loop, since it is nilpotent but not central by a 2-Engel group.

**§4. A multiplication table and a look at normal subgroups of loops**

**Example 4.1** (A  $A_l$ -Bol-loop multiplication table). The lowest order of a nilpotent group of class 3 which is not of class 2 is 16. Using the software package MAGMA ([2]) we found three non-isomorphic nilpotent groups of order 16 which are of class 3 but are not of class 2. Their associated left gyrogroup generate three non- $K$ -loops (i.e., non-Bruck-loops)  $A_l$ -Bol-loops of order 16, denoted by  $K_{16}$ ,  $L_{16}$ , and  $M_{16}$ . The multiplication table of  $K_{16}$ , is presented in Table I, where the elements  $k_i \in K_{16}$ ,  $i = 1, 2, \dots, 16$ , are denoted by their subscripts.

**Table I** (*The  $A_l$ -Bol-loop  $K_{16}$* )

$\circ$		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
—		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
1		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
2		2	1	4	3	6	5	8	7	10	9	12	11	14	13	16	15
3		3	4	2	1	7	8	6	5	12	11	9	10	16	15	13	14
4		4	3	1	2	8	7	5	6	11	12	10	9	15	16	14	13
5		5	6	7	8	4	3	1	2	16	15	13	14	10	9	12	11
6		6	5	8	7	3	4	2	1	15	16	14	13	9	10	11	12
7		7	8	6	5	1	2	3	4	14	13	16	15	11	12	10	9
8		8	7	5	6	2	1	4	3	13	14	15	16	12	11	9	10
9		9	10	11	12	13	14	15	16	1	2	3	4	5	6	7	8
10		10	9	12	11	14	13	16	15	2	1	4	3	6	5	8	7
11		11	12	10	9	15	16	14	13	4	3	1	2	8	7	5	6
12		12	11	9	10	16	15	13	14	3	4	2	1	7	8	6	5
13		13	14	15	16	12	11	9	10	7	8	6	5	1	2	3	4
14		14	13	16	15	11	12	10	9	8	7	5	6	2	1	4	3
15		15	16	14	13	9	10	11	12	5	6	7	8	4	3	1	2
16		16	15	13	14	10	9	12	11	6	5	8	7	3	4	2	1

$K_{16}$  has only one non-identity left inner mapping,  $A$ , whose transformation table is given in Table II.

**Table II** (*The automorphism  $A$  of  $K_{16}$* )

$1 \rightarrow 1$	$5 \rightarrow 5$	$9 \rightarrow 10$	$13 \rightarrow 14$
$2 \rightarrow 2$	$6 \rightarrow 6$	$10 \rightarrow 9$	$14 \rightarrow 13$
$3 \rightarrow 3$	$7 \rightarrow 7$	$11 \rightarrow 12$	$15 \rightarrow 16$
$4 \rightarrow 4$	$8 \rightarrow 8$	$12 \rightarrow 11$	$16 \rightarrow 15$

The left inner mapping  $\delta_{a,b}$  generated by any  $a, b \in K_{16}$  is either  $A$  or the identity automorphism denoted by 1. The left inner mapping table for  $\delta_{a,b}$  is presented in Table III.

**Table III** (*The left inner mapping  $\delta_{a,b}$  of  $K_{16}$* )

$\delta$		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
—		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
1		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
2		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
3		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
4		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
5		1	1	1	1	1	1	1	1	$A$	$A$	$A$	$A$	$A$	$A$	$A$	$A$
6		1	1	1	1	1	1	1	1	$A$	$A$	$A$	$A$	$A$	$A$	$A$	$A$
7		1	1	1	1	1	1	1	1	$A$	$A$	$A$	$A$	$A$	$A$	$A$	$A$
8		1	1	1	1	1	1	1	1	$A$	$A$	$A$	$A$	$A$	$A$	$A$	$A$
9		1	1	1	1	$A$	$A$	$A$	$A$	1	1	1	1	$A$	$A$	$A$	$A$
10		1	1	1	1	$A$	$A$	$A$	$A$	1	1	1	1	$A$	$A$	$A$	$A$
11		1	1	1	1	$A$	$A$	$A$	$A$	1	1	1	1	$A$	$A$	$A$	$A$
12		1	1	1	1	$A$	$A$	$A$	$A$	1	1	1	1	$A$	$A$	$A$	$A$
13		1	1	1	1	$A$	$A$	$A$	$A$	$A$	$A$	$A$	$A$	1	1	1	1
14		1	1	1	1	$A$	$A$	$A$	$A$	$A$	$A$	$A$	$A$	1	1	1	1
15		1	1	1	1	$A$	$A$	$A$	$A$	$A$	$A$	$A$	$A$	1	1	1	1
16		1	1	1	1	$A$	$A$	$A$	$A$	$A$	$A$	$A$	$A$	1	1	1	1

**Definition 4.2.** A subloop  $X$  of a loop  $P$  is a *normal subgroup* ([7]) of  $P$  if it is a normal subloop which is in the middle nucleus i.e.

- (i)  $\delta_{a,x} = 1$  for all  $x \in X$  and  $a \in P$ ;
- (ii)  $\delta_{a,b}(X) \subseteq X$  for all  $a, b \in P$ ;
- (iii)  $a \odot X = X \odot a$  for all  $a \in P$ .

*Note.* If  $X$  is a normal subgroup of a loop  $P$ , then  $X$  is a group with group operation given by the restriction of  $\odot$  to  $X$ .

*Note.* Since a normal subgroup  $X$  of  $P$  is a normal subloop,  $P/X$  forms a factor loop.

**Theorem 4.5.** *If  $(P, \odot)$  is a  $A_l$ -Bol-loop, then  $P$  has a normal subgroup  $\Xi$  such that  $P/\Xi$  is a Bruck-loop.*

PROOF: ([7, Theorem 4.11]). □

**Example 4.6.** The  $4 \times 4$  upper left corner of Table I forms a multiplication table of a group,  $H$ . The group  $H$  is a normal subgroup of  $K_{16}$ . The quotient  $K_{16}/H$  turns out to be an abelian group. Hence, we have in hand an example of an extension of a group by another group that gives a non-associative structure (that is, the  $A_l$ -Bol-loop  $K_{16}$ ). It is an extension which is far from being trivial since  $H$  and  $K_{16}/H$  are groups while  $K_{16}$  is a non-Bruck-loop  $A_l$ -Bol-loop.

## §5. Open questions

**Question 5.1.** Are there any nontrivial  $AP$ -transversals that are loops in finite simple groups (The answer is positive in the infinite case [12], but I suspect that it is negative in the finite case)?

**Question 5.2.** For which groups is the associated left gyrogroup an  $AP$ -loop?

## REFERENCES

- [1] Bruck R.H., *A Survey of Binary Systems*, Springer-Verlag, 1966.
- [2] Cannon J., Playoust C., *An introduction to MAGMA*, University of Sydney, Sydney, 1993.
- [3] Conway J.H., *A simple construction for the Fischer-Griess monster group*, *Invent. Math.* **79** (1985), no. 3, 513–540.
- [4] Foguel T., *Groups with all cyclic subgroups conjugate-permutable groups*, *J. Group Theory* **2** (1999), no. 1, 47–51.
- [5] Foguel T., *Conjugate-permutable subgroups*, *J. Algebra* **191** (1997), no. 1, 235–239.
- [6] Foguel T., Ungar A.A., *Gyrogroups and the decomposition of groups into twisted subgroups and subgroups*, *Pacific J. Math.*, to appear.
- [7] Foguel T., Ungar A.A., *Involutory decomposition of groups into twisted subgroups and subgroups*, *J. Group Theory* **3** (2000), no. 1, 27–46.
- [8] Griess R.L., Jr., *Code loops*, *J. Algebra* **100** (1986), no. 1, 224–234.
- [9] Karzel H., *Raum-Zeit-Welt und hyperbolische Geometrie*, *Beiträge zur Geometrie und Algebra* **29** (1994), Technische Universität München, Mathematisches Institut, Munich.
- [10] Kepka T., Niemenmaa M., *On multiplication groups of loops*, *J. Algebra* **135** (1990), 112–122.
- [11] Kepka T., Phillips J.D., *Connected transversals to subnormal subgroups*, *Comment. Math. Univ. Carolinae* **38** (1997), 223–230.
- [12] Kiechle H., *K-loops from classical groups over ordered fields*, *J. Geom.* **61** (1998), no. 1–2, 105–127.
- [13] Kinyon M.K., Jones O., *Loops and semidirect products*, *Comm. Algebra*, submitted.
- [14] Robinson D.J.S., *A Course in the Theory of Groups*, Springer, New York, 1995.
- [15] Sabinin L.V., Sabinina L.L., Sbitneva L.V., *On the notion of gyrogroup*, *Aequationes Math.* **56** (1998), 11–17.



- [16] Sabinin L.V., *On the equivalence of categories of loop and homogeneous spaces*, Soviet Math. Dokl. **13** (1972), no. 4, 970–974.
- [17] Ungar A.A., *Axiomatic approach to the nonassociative group of relativistic velocities*, Found. Phys. Lett. **2** (1989), 199–203.

DEPARTMENT OF MATHEMATICS, NORTH DAKOTA STATE UNIVERSITY, FARGO, ND 58105,  
USA

*E-mail:* foguel@prairie.Nodak.edu

(Received September 7, 1999, revised January 10, 2000)