

## Sequentially compact sets in a class of generalized Orlicz spaces

JINCAI WANG

*Abstract.* In this paper, we will characterize sequentially compact sets in a class of generalized Orlicz spaces.

*Keywords:* generalized Orlicz space  $L^{(M^{-1})}$ , sequentially compact set,  $\Delta_2$ -condition

*Classification:* 46B20, 46E30

### §1. Introduction and basic results

**Definition 1.1.**  $M : \mathbb{R} \rightarrow \mathbb{R}$  is called an *Orlicz function* if it has the following properties:

- (1)  $M$  is even, continuous, convex on  $(0, \infty)$  and  $M(0) = 0$ ,
- (2)  $M(u) > 0$  for all  $u \neq 0$ ,
- (3)  $\lim_{u \rightarrow 0} \frac{M(u)}{u} = 0$  and  $\lim_{u \rightarrow +\infty} \frac{M(u)}{u} = +\infty$ .

**Definition 1.2.** A function  $\phi : [0, \infty) \rightarrow [0, \infty)$  is called a  *$\varphi$ -function*, if  $\phi$  satisfies

- (i)  $\phi(0) = 0, \phi(u) > 0, u > 0$ ;
- (ii)  $\phi(u)$  is increasing, continuous;
- (iii)  $\lim_{x \rightarrow \infty} \phi(x) = \infty$ .

A  $\varphi$ -function  $\phi(u)$  is said to satisfy the  $\Delta_2$ -condition for small  $u$  (for all  $u \geq 0$  or for large  $u$ ), in symbol  $\phi \in \Delta_2(0)$  ( $\phi \in \Delta_2$  or  $\phi \in \Delta_2(\infty)$ ), if there exists  $u_0 > 0$  and  $c > 0$  such that  $\phi(2u) \leq c\phi(u)$  for  $0 \leq u \leq u_0$  (for all  $u \geq 0$  or for  $u \geq u_0$ ). The generalized Orlicz class, generalized Orlicz space and subspace, respectively, generated by a  $\varphi$ -function  $\phi$  are defined as follows:

$$\begin{aligned} \tilde{L}^\phi(G) &= \{x(t) : x(t) \text{ is measurable on a Lebesgue measurable set } G \text{ and} \\ &\quad \rho_\phi(x) = \int_G \phi(|x(t)|) dt < \infty\}, \\ L^\phi(G) &= \{x(t) : x(t) \text{ is measurable on a Lebesgue measurable set } G \text{ and} \\ &\quad \rho_\phi(\lambda x) < \infty \text{ for some } \lambda > 0\}, \\ E^\phi(G) &= \{x(t) : x(t) \text{ is measurable on a Lebesgue measurable set } G \text{ and} \\ &\quad \rho_\phi(\lambda x) < \infty \text{ for all } \lambda > 0\}. \end{aligned}$$

Now we denote by  $M^{-1}$  the inverse function to an Orlicz function  $M$  on  $[0, \infty)$ . It is obvious that  $M^{-1}$  is a  $\varphi$ -function and it is a concave function. Since  $M^{-1}(2u) < 2M^{-1}(u)$  for all  $u \geq 0$ , we have  $M^{-1} \in \Delta_2$ . It follows that  $\tilde{L}^{M^{-1}}(G) = L^{M^{-1}}(G) = E^{M^{-1}}(G)$  (see [1]). So we only study one of them. The theory of generalized Orlicz spaces can be found in [2]. For convenience, we give the following definition.

**Definition 1.3.** Let  $M^{-1}$  be the inverse function to an Orlicz function  $M$  on  $[0, \infty)$ , let  $(\Omega, \Sigma, \mu)$  be a measure space. We define a class of generalized Orlicz spaces as follows:

$$(1) \quad L^{M^{-1}}(\Omega, \Sigma, \mu) = \{x(t) : x(t) \text{ is } \mu \text{ measurable and } \rho_{M^{-1}}(x) < \infty\}.$$

Let  $X$  be a vector space over  $\mathbb{K}$ .  $\|\cdot\| : X \rightarrow \mathbb{R}$  is called an F-norm if

- (i) for every  $x \in X$ ,  $\|x\| \geq 0$ , and  $\|x\| = 0 \iff x = 0$ ,
- (ii)  $\|-x\| = \|x\|$ ,
- (iii)  $\|x + y\| \leq \|x\| + \|y\|$ ,
- (iv)  $\|x_n\| \rightarrow 0 \implies \|\alpha x_n\| \rightarrow 0$  for  $\alpha \in \mathbb{K}$  and  $\alpha_n \rightarrow 0 \implies \|\alpha_n x\| \rightarrow 0$  for  $x \in X$ .

As in a generalized Orlicz space, we can define the normal F-norm in  $L^{M^{-1}}$  as follows:

$$(2) \quad \|x\|_{M^{-1}} = \inf \left\{ c > 0 : \rho_{M^{-1}} \left( \frac{x}{c} \right) \leq c \right\}.$$

But we can define a more simple F-norm in  $L^{M^{-1}}$  as in Theorem 1.1 below. The new F-norm is equivalent to the normal F-norm (2). Using the new simple F-norm, we can investigate  $L^{M^{-1}}$  which we can call a *generalized Orlicz space generated by the Orlicz function  $M$* . These generalized Orlicz spaces include  $L^p$  ( $0 < p < 1$ ).

**Theorem 1.1.**  $\|x\|_{(M^{-1})} = \rho_{M^{-1}}(x)$ ,  $x \in L^{M^{-1}}$ , is an F-norm in  $L^{M^{-1}}$ , and  $L^{(M^{-1})} = (L^{M^{-1}}, \|\cdot\|_{(M^{-1})})$  is a linear complete space with F-norm  $\|\cdot\|_{(M^{-1})}$ . So it is a Fréchet space.

PROOF: By the basic properties of an Orlicz function  $M$  (see [1]), we have

$$(3) \quad M^{-1}(|u + v|) \leq M^{-1}(|u| + |v|) \leq M^{-1}(|u|) + M^{-1}(|v|),$$

$$(4) \quad M^{-1}(\lambda|u|) \leq \lambda M^{-1}(|u|), \quad \lambda \geq 1,$$

$$(5) \quad M^{-1}(\lambda|u|) \leq M^{-1}(|u|), \quad 0 < \lambda < 1.$$

These estimates imply that  $L^{(M^{-1})}$  is a linear space and  $\|\cdot\|_{(M^{-1})}$  is an F-norm. □

Some properties of the generalized space  $L^{(M^{-1})}$  have been established in [3] by M.M. Rao and Z.D. Ren. In this paper, we will discuss the criteria for a set to be sequentially compact in  $L^{(M^{-1})}$ .

The three criteria for a set to be sequentially compact in  $L^p[a, b]$  ( $1 < p < \infty$ ) were introduced by F. Riesz [4], Kolomogorov [5] and Krasnoselskii [6], respectively. The three criteria were generalized to  $E^M[a, b]$ , a closed subset of Orlicz space  $L^M[a, b]$ , by Takahashi [7], Griбанov [8] and Krasnoselskii [6], respectively. In 1951, Tsuji [9] generalized the F. Riesz criterion to the space  $L^p$  ( $0 < p < 1$ ).

Now we generalize the F. Riesz criterion and Krasnoselskii criterion for sequential compactness to the *generalized Orlicz space generated by the Orlicz function*  $L^{(M^{-1})}(G)$ .

### §2. The Riesz criterion for sequential compactness in $L^{(M^{-1})}$

A set  $A \subset L^{(M^{-1})}[a, b]$  is said to be *sequentially compact* if, for any  $\{x_n(t)\}_{n=1}^\infty \subset A$ , there exists a subsequence  $\{x_{n_i}(t)\}_{i=1}^\infty$  and  $x_0(t) \in L^{(M^{-1})}[a, b]$  such that  $\lim_{i \rightarrow \infty} \|x_{n_i} - x_0\|_{(M^{-1})} = 0$ .

**Lemma 2.1.** Let  $M(u) = \int_0^{|u|} p(t) dt$  be an Orlicz function. Then for  $y > x > 0$ ,

$$(6) \quad \frac{y - x}{p[M^{-1}(y)]} \leq M^{-1}(y) - M^{-1}(x) \leq \frac{y - x}{p[M^{-1}(x)]}.$$

PROOF: Let  $0 < t_1 < t_2$ . Since

$$M(t_2) - M(t_1) = \int_{t_1}^{t_2} p(t) dt$$

and  $p(t_1) \leq p(t) \leq p(t_2)$  ( $t_1 \leq t \leq t_2$ ), we have

$$p(t_1)(t_2 - t_1) \leq M(t_2) - M(t_1) \leq p(t_2)(t_2 - t_1)$$

or equivalently,

$$\frac{1}{p(t_2)}(M(t_2) - M(t_1)) \leq t_2 - t_1 \leq \frac{1}{p(t_1)}(M(t_2) - M(t_1)).$$

Since  $M(t)$  is strictly increasing,  $M(0) = 0$  and  $M(\infty) = \infty$ , we can set  $x = M(t_1)$ ,  $y = M(t_2)$ . Thus we get (6).  $\square$

**Lemma 2.2.** *Let  $M(u) = \int_0^{|u|} p(t) dt$  be an Orlicz function and let  $M \in \Delta_2$ , then*

(i) *for nonnegative real numbers  $x \neq y$ ,*

$$(7) \quad M^{-1}(|y - x|) \leq \frac{\overline{B}_M \max\{p(M^{-1}(x)), p(M^{-1}(y))\}}{p[M^{-1}(|y - x|)]} |M^{-1}(|y|) - M^{-1}(|x|)|,$$

where

$$(8) \quad \overline{B}_M = \sup_{t>0} \frac{tp(t)}{M(t)}.$$

(ii) *Let  $a > 1$  and  $0 < y < x < ay$ , then*

$$(9) \quad p[M^{-1}(x)] < a\overline{B}_M p[M^{-1}(y)].$$

PROOF: (i). Since  $M \in \Delta_2$ , we have  $\overline{B}_M < \infty$  (see [1]). So, by (8), for nonnegative real numbers  $x \neq y$ , we have

$$\frac{M^{-1}(|y - x|)p[M^{-1}(|y - x|)]}{|y - x|} \leq \overline{B}_M$$

or equivalently

$$|y - x| \geq \frac{1}{\overline{B}_M} \{M^{-1}(|y - x|)p[M^{-1}(|y - x|)]\}.$$

By Lemma 2.1, we get

$$\begin{aligned} |M^{-1}(|y|) - M^{-1}(|x|)| &\geq \frac{1}{\max\{p(M^{-1}(|x|)), p(M^{-1}(|y|))\}} |y - x| \\ &\geq \frac{M^{-1}(|y - x|)p[M^{-1}(|y - x|)]}{\overline{B}_M \max\{p(M^{-1}(|x|)), p(M^{-1}(|y|))\}}, \end{aligned}$$

i.e.,

$$M^{-1}(|y - x|) \leq \frac{\overline{B}_M \max\{p(M^{-1}(|x|)), p(M^{-1}(|y|))\}}{p[M^{-1}(|y - x|)]} |M^{-1}(|y|) - M^{-1}(|x|)|.$$

(ii) Since  $\frac{M^{-1}(y)p[M^{-1}(y)]}{y} > 1$ ,  $y > 0$ , by (8), we have

$$p[M^{-1}(ay)] \leq \overline{B}_M \frac{ay}{M^{-1}(ay)} < a\overline{B}_M \frac{y}{M^{-1}(y)} < a\overline{B}_M p[M^{-1}(y)].$$

So  $p[M^{-1}(x)] \leq p[M^{-1}(ay)] \leq a\overline{B}_M p[M^{-1}(y)]$ .  $\square$

For the proof of the main theorem, we need the following result.

**Lemma 2.3** (Riesz).  $F \subset L^p[a, b]$  ( $1 \leq p < \infty$ ) is a sequentially compact set if and only if

- (1)  $F$  is bounded:  $\exists c > 0$ , such that  $\sup_{x \in F} \|x(t)\|_p \leq c < \infty$ ;
- (2)  $F$  is equicontinuous:  $\forall \varepsilon > 0, \exists \delta > 0$ , such that for  $|h| < \delta$ ,

$$\sup_{x \in F} \|x(t+h) - x(t)\|_p < \varepsilon.$$

The main result in this section is

**Theorem 1.1.** Let  $M \in \Delta_2$  be an Orlicz function. Then a set  $F \subset L^{(M^{-1})}[a, b]$  is sequentially compact if and only if

- (1)  $F$  is bounded:  $\exists c > 0$ , such that

$$(10) \quad \sup_{x \in F} \int_a^b M^{-1}(|x(t)|) dt \leq c.$$

- (2)  $F$  is equicontinuous:  $\forall \varepsilon > 0, \exists \delta > 0$  such that for  $|h| < \delta$ ,

$$(11) \quad \sup_{x \in F} \int_a^b M^{-1}(|x(t+h) - x(t)|) dt < \varepsilon.$$

PROOF: *Sufficiency.* If  $x(t) \in L^{(M^{-1})}[a, b]$ , then  $M^{-1}(|x(t)|) \in L^1[a, b]$ . Now we consider the function family  $F' = \{M^{-1}(|x(t)|) : x(t) \in F\} \subset L^1[a, b]$ . By condition (1), we know  $F'$  is bounded in  $L^1[a, b]$ , i.e.  $\|M^{-1}(|x(t)|)\|_1 \leq c$  for all  $M^{-1}(|x(t)|) \in F'$ . For every Orlicz function  $M(u)$  and  $x(t) \in L^{(M^{-1})}[a, b]$ , by basic properties of Orlicz functions (see [1]), we have

$$|M^{-1}(|x(t+h)|) - M^{-1}(|x(t)|)| \leq M^{-1}(|x(t+h) - x(t)|).$$

So by (2),  $\forall \varepsilon > 0, \exists \delta > 0$ , such that for  $|h| < \delta$ , we have

$$\sup_{M^{-1}(|x(t)|) \in F'} \int_G |M^{-1}(|x(t+h)|) - M^{-1}(|x(t)|)| dt < \varepsilon.$$

By Lemma 2.3 we know that  $F' \subset L^1[a, b]$  is sequentially compact in  $L^1[a, b]$ . So for any  $\{x_n\}_{n=1}^\infty \subset F$  there exists a subsequence (denoted without risk of confusion still by  $\{x_n\}_{n=1}^\infty$ ) and  $x_0 \in L^1[a, b]$ , such that  $\{M^{-1}(|x_n|)\}_{n=1}^\infty \xrightarrow{\|\cdot\|_1} x_0$ .

Since  $M \in \Delta_2$ ,  $\forall 0 < \varepsilon < 6c$  (where  $c$  is introduced in condition (1)),  $\exists K_\varepsilon > 0$ , such that for  $u > 0$ , we have  $M(\frac{u}{K_\varepsilon}) \leq K_\varepsilon M(u)$ , or equivalently,

$$(12) \quad M^{-1}\left(\frac{v}{K_\varepsilon}\right) \leq \frac{\varepsilon}{6c} M^{-1}(v), \quad v > 0.$$

Choose  $\varepsilon' < \varepsilon$ , such that  $\sqrt{\varepsilon'} < \frac{1}{K_\varepsilon}$ ,  $\sqrt{\varepsilon'}\mu([a, b]) < \frac{\varepsilon}{3}$  and  $\overline{B}_M^2\sqrt{\varepsilon'} < \frac{\varepsilon}{3}$ . Thus for any  $n \in \mathbb{N}$ , by (12) we get

$$(13) \quad \begin{aligned} \int_{[a, b]} M^{-1}(\sqrt{\varepsilon'}|x_n(t)|) dt &< \int_{[a, b]} M^{-1}\left(\frac{|x_n(t)|}{K_\varepsilon}\right) dt \\ &\leq \frac{\varepsilon}{6c} \int_{[a, b]} M^{-1}(|x_n(t)|) dt \leq \frac{\varepsilon}{6c} \cdot c = \frac{\varepsilon}{6}. \end{aligned}$$

Since  $\{M^{-1}(|x_n|)\}_{n=1}^\infty \xrightarrow{\|\cdot\|_1} x_0 \in L^1[a, b]$ , for  $\varepsilon' > 0$ , there exists  $N(\varepsilon) > 0$  such that for any  $n, m > N(\varepsilon)$ , we have

$$(14) \quad \int_{[a, b]} |M^{-1}(|x_n(t)|) - M^{-1}(|x_m(t)|)| dt < \varepsilon'.$$

So for any  $n, m > N(\varepsilon)$ , we set

$$\begin{aligned} G_1 &= \{t \in [a, b] : M^{-1}(|x_n(t) - x_m(t)|) < \sqrt{\varepsilon'}\}, \\ G_2 &= \left\{t \in [a, b] : M^{-1}(|x_n(t) - x_m(t)|) \geq \sqrt{\varepsilon'} \text{ and } \frac{|x_n(t)| + |x_m(t)|}{|x_n(t) - x_m(t)|} < \frac{1}{\sqrt{\varepsilon'}}\right\}, \\ G_3 &= \left\{t \in [a, b] : M^{-1}[|x_n(t) - x_m(t)|] \geq \sqrt{\varepsilon'} \text{ and } \frac{|x_n(t)| + |x_m(t)|}{|x_n(t) - x_m(t)|} \geq \frac{1}{\sqrt{\varepsilon'}}\right\}. \end{aligned}$$

Obviously,  $G_1, G_2, G_3$  are pairwise disjoint, and  $[a, b] = G_1 \cup G_2 \cup G_3$ . Thus

$$\begin{aligned} \int_{[a, b]} M^{-1}[|x_n(t) - x_m(t)|] dt &= \left(\int_{G_1} + \int_{G_2} + \int_{G_3}\right) M^{-1}[|x_n(t) - x_m(t)|] dt, \\ \int_{G_1} M^{-1}[|x_n(t) - x_m(t)|] dt &\leq \int_{G_1} \sqrt{\varepsilon'} dt \leq \sqrt{\varepsilon'}\mu([a, b]) < \frac{\varepsilon}{3}. \end{aligned}$$

For  $G_2$ ,

$$|x_n(t) - x_m(t)| \leq |x_n(t)| + |x_m(t)| < \frac{1}{\sqrt{\varepsilon'}} |x_n(t) - x_m(t)|.$$

By Lemma 2.2 and (14), we get

$$\begin{aligned} & \int_{G_2} M^{-1}[|x_n(t) - x_m(t)|] dt \\ & \leq \int_{G_2} \frac{\overline{B}_M \max\{p(M^{-1}(|x_n(t)|)), p(M^{-1}|x_m(t)|)\}}{p(M^{-1}(|x_n(t) - x_m(t)|))} \cdot \\ & \quad \cdot |M^{-1}(|x_n(t)|) - M^{-1}(|x_m(t)|)| dt \\ & \leq \int_{G_2} \frac{\overline{B}_M p[M^{-1}(|x_n(t)| + |x_m(t)|)]}{p[M^{-1}(|x_n(t) - x_m(t)|)]} |M^{-1}(|x_n(t)|) - M^{-1}(|x_m(t)|)| dt \\ & \leq \int_{G_2} \overline{B}_M \frac{1}{\sqrt{\varepsilon'}} \overline{B}_M |M^{-1}(|x_n(t)|) - M^{-1}(|x_m(t)|)| dt \\ & \leq \frac{\overline{B}_M^2}{\sqrt{\varepsilon'}} \cdot \varepsilon' = \overline{B}_M^2 \sqrt{\varepsilon'} < \frac{\varepsilon}{3}. \end{aligned}$$

For  $G_3$ ,  $|x_n(t) - x_m(t)| \leq \sqrt{\varepsilon'}(|x_n(t)| + |x_m(t)|)$  and we get by (13)

$$\begin{aligned} & \int_{G_3} M^{-1}(|x_n(t) - x_m(t)|) dt \\ & \leq \int_{G_3} M^{-1}[\sqrt{\varepsilon'}(|x_n(t)| + |x_m(t)|)] dt \\ & = \int_{G_3} M^{-1}(\sqrt{\varepsilon'}|x_n(t)|) dt + \int_{G_3} M^{-1}(\sqrt{\varepsilon'}|x_m(t)|) dt < \frac{\varepsilon}{6} + \frac{\varepsilon}{6} = \frac{\varepsilon}{3}. \end{aligned}$$

So for  $\varepsilon > 0$ , there exists  $N(\varepsilon) > 0$  such that for  $n, m > N(\varepsilon)$ , we have

$$\int_G M^{-1}[|x_n(t) - x_m(t)|] dt < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon,$$

i.e.,  $\{x_n\}_{n=1}^\infty$  is a Cauchy sequence in  $L^{(M^{-1})}[a, b]$ . Since  $L^{(M^{-1})}[a, b]$  is complete, there exists  $y_0 \in L^{(M^{-1})}[a, b]$ , such that  $\lim_{n \rightarrow \infty} \int_{[a,b]} M^{-1}[|x_n(t) - y_0(t)|] dt = 0$ .

*Necessity.* Denote  $G = [a, b]$ . We assume that  $F$  is a sequentially compact set and condition (1) does not hold. Then there exists  $\{x_n(t)\}_{n=1}^\infty \subset F$ , such that

$$\int_G M^{-1}(|x_n(t)|) dt > n,$$

and there exists a subsequence of  $\{x_n(t)\}_1^\infty$  (we still denote it by  $\{x_n(t)\}_1^\infty$ ) which converges to  $x_0(t) \in L^{(M^{-1})}(G)$  in  $\|\cdot\|_{(M^{-1})}$ . Thus for  $\varepsilon_0 = 1$ , there exists  $N > 0$  such that, for  $n > N$ ,

$$\int_G M^{-1}[|x_n(t) - x_0(t)|] dt < 1.$$

On the other hand, by  $M^{-1}(|x_0|) \geq M^{-1}(|x_n|) - M^{-1}(|x_n - x_0|)$ , we have

$$\begin{aligned} \int_G M^{-1}(|x_0(t)|) dt &\geq \int_G M^{-1}(|x_n(t)|) dt - \int_G M^{-1}(|x_n(t) - x_0(t)|) dt \\ &> n - 1 \rightarrow \infty \quad (n \rightarrow \infty), \end{aligned}$$

so  $x_0 \notin L^{(M^{-1})}(G)$ . This is impossible. The proof of the necessity of (1) is complete.

Since the family of continuous function  $C(G)$  is dense in  $L^{(M^{-1})}(G)$ , and a continuous function is uniformly continuous in a bounded closed set  $G$ , for any  $x(t) \in F$  and any  $\varepsilon > 0$ , there exists  $x_c(t) \in C(G)$  such that

$$\int_G M^{-1}[x(t) - x_c(t)] dt < \frac{\varepsilon}{3},$$

and there exists  $\delta(\varepsilon) > 0$  such that

$$\int_G M^{-1}[x_c(t+h) - x_c(t)] dt < \frac{\varepsilon}{3} \quad \text{for } |h| < \delta(\varepsilon).$$

Thus for  $|h| < \delta(\varepsilon)$ ,

$$\begin{aligned} &\int_G M^{-1}[|x(t+h) - x(t)|] dt \\ &\leq \int_G M^{-1}[|x(t+h) - x_c(t+h)|] dt + \int_G M^{-1}[|x_c(t+h) - x_c(t)|] dt \\ &\quad + \int_G M^{-1}[|x_c(t) - x(t)|] dt < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon. \end{aligned}$$

The necessity of (2) is proved. □

**Corollary 2.1.** For  $L^p[a, b]$  ( $0 < p < 1$ ) with  $F$ -norm  $\|x\|_p = \int_a^b |x(t)|^p dt$ ,  $A \subset L^p[a, b]$  is sequentially compact if and only if

- (i)  $\exists c < \infty$ , such that  $\sup_{x \in A} \|x\|_p \leq c$ ;
- (ii)  $\forall \varepsilon > 0, \exists \delta > 0$ , such that for  $|h| < \delta$ ,

$$\sup_{x \in A} \|x(t+h) - x(t)\|_p < \varepsilon.$$



**Theorem 2.2.** *Let  $M$  be an Orlicz function such that  $M \in \Delta_2$ . Then  $F \subset L^{(M^{-1})}(-\infty, +\infty)$  is sequentially compact if and only if*

- (i) *there exists  $c > 0$  such that  $\sup_{x \in F} \int_{-\infty}^{+\infty} M^{-1}(|x(t)|) dt \leq c$ ,*
- (ii)  *$\lim_{N \rightarrow \infty} \int_{|t| \geq N} M^{-1}(|x(t)|) dt = 0$  uniformly for  $x(t) \in F$ ,*
- (iii) *for any  $\varepsilon > 0$  there exists  $\delta > 0$  such that for  $|h| < \delta$ ,*

$$\sup_{x \in F} \int_{(-\infty, +\infty)} M^{-1}[|x(t+h) - x(t)|] dt < \varepsilon.$$

**PROOF:** *Sufficiency.* For  $[-1, 1]$ , by (i), (iii) and Theorem 2.1, we know that the set  $\{f|_{[-1,1]} : f \in F\}$  is sequentially compact. So there exists a convergent subsequence  $\{x_{n_k}^{(1)}\}_{k=1}^\infty \subset F$  in  $L^{(M^{-1})}[-1, 1]$ .

For  $[-2, 2]$ , by (i) and (iii), the set  $\{x_{n_k}^{(1)}|_{[-2,2]}\}_{k=1}^\infty$  is sequentially compact. So there exists a convergent subsequence  $\{x_{n_k}^{(2)}\}_{k=1}^\infty \subset \{x_{n_k}^{(1)}\}_{k=1}^\infty \subset F$  in  $L^{(M^{-1})}[-2, 2]$ .

Going on in this way, we get

- $l_1 : x_{n_1}^{(1)}, x_{n_2}^{(1)}, \dots, x_{n_k}^{(1)}, \dots$  convergence in  $L^{(M^{-1})}[-1, 1]$ ,
- $l_2 : x_{n_1}^{(2)}, x_{n_2}^{(2)}, \dots, x_{n_k}^{(2)}, \dots$  convergence in  $L^{(M^{-1})}[-2, 2]$ ,
- ... ..
- $l_m : x_{n_1}^{(m)}, x_{n_2}^{(m)}, \dots, x_{n_k}^{(m)}, \dots$  convergence in  $L^{(M^{-1})}[-m, m]$ ,
- ... ..

satisfying  $l_1 \supset l_2 \supset \dots \supset l_m \supset \dots$ .

Choosing diagonal elements  $x_{n_1}^{(1)}, x_{n_2}^{(2)}, \dots, x_{n_k}^{(k)}, \dots$ , we get a Cauchy sequence in  $L^{(M^{-1})}(-\infty, +\infty)$ . In fact, by (ii), for  $\varepsilon > 0$ , there exists an  $N \in \mathbb{N}$  such that

$$\int_{|t| \geq N} M^{-1}(|x_{n_k}^{(k)}(t)|) dt < \frac{\varepsilon}{4}, \quad k = 1, 2, \dots.$$

For  $[-N, N]$ ,  $\{x_k^{(k)}\}_{k=N}^\infty$  converges in  $L^{(M^{-1})}[-N, N]$  (assume it converges to  $x_0 \in L^{(M^{-1})}[-N, N]$ ). Thus for  $\varepsilon > 0$ , there exists a  $K_0 \in \mathbb{N}$  such that for

$k, l > K_0$ , we have

$$\begin{aligned}
 & \int_{[-N, N]} M^{-1}[|x_{n_k^{(k)}}(t) - x_{n_l^{(l)}}(t)|] dt \\
 & \leq \int_{[-N, N]} M^{-1}[|x_{n_k^{(k)}}(t) - x_0(t)| + |x_0(t) - x_{n_l^{(l)}}(t)|] dt \\
 & \leq \int_{[-N, N]} M^{-1}[|x_{n_k^{(k)}}(t) - x_0(t)|] dt + \int_{[-N, N]} M^{-1}[|x_0(t) - x_{n_l^{(l)}}(t)|] dt \\
 & \leq \frac{\varepsilon}{4} + \frac{\varepsilon}{4} = \frac{\varepsilon}{2}.
 \end{aligned}$$

So for  $\varepsilon > 0$  and for all  $k, l > K_0$ , we have

$$\begin{aligned}
 & \int_{-\infty}^{+\infty} M^{-1}[|x_{n_k^{(k)}}(t) - x_{n_l^{(l)}}(t)|] dt \\
 & \leq \int_{[-N, N]} M^{-1}[|x_{n_k^{(k)}}(t) - x_{n_l^{(l)}}(t)|] dt + \int_{|t| \geq N} M^{-1}[|x_{n_k^{(k)}}(t) - x_{n_l^{(l)}}(t)|] dt \\
 & \leq \frac{\varepsilon}{2} + \int_{|t| \geq N} M^{-1}(|x_{n_k^{(k)}}(t)|) dt + \int_{|t| \geq N} M^{-1}[|x_{n_l^{(l)}}(t)|] dt \\
 & \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{4} + \frac{\varepsilon}{4} = \varepsilon.
 \end{aligned}$$

This means that  $\{x_{n_k^{(k)}}\}_{k=1}^{\infty}$  is a Cauchy sequence in  $L^{(M^{-1})}(-\infty, +\infty)$  and, since  $L^{(M^{-1})}(-\infty, +\infty)$  is complete, we know the sufficiency is true.

*Necessity.* The proof of (i) is similar to that of Theorem 2.1(1), now we prove (ii) and (iii). If (ii) does not hold, then there exists an  $\varepsilon_0 > 0$  such that for any  $n_k > 0$  there is  $x_{n_k} \in F$  with

$$\int_{|t| > n_k} M^{-1}(|x_{n_k}(t)|) dt > \varepsilon_0.$$

Choose  $n_1 = 1$ . Then  $\exists x_1 \in F$  such that

$$\int_{|t| > 1} M^{-1}(|x_1(t)|) dt > \varepsilon_0.$$

Noting that

$$\int_{(-\infty, +\infty)} M^{-1}(|x_1(t)|) dt < \infty,$$

$\exists n_2 > 0$  such that

$$\int_{|t| > n_2} M^{-1}(|x_1(t)|) dt < \frac{\varepsilon_0}{2}.$$

For  $n_2$ , there exists  $x_2 \in F$  such that  $\int_{|t|>n_2} M^{-1}(|x_2(t)|) dt > \varepsilon_0$ . Thus  $\exists n_3 > 0$  such that

$$\int_{|t|>n_3} M^{-1}(|x_2(t)|) dt < \frac{\varepsilon_0}{2}.$$

Following these steps, we get  $\{x_1, x_2, \dots, x_k, \dots\} \subset F$  and

$$\begin{aligned} \int_{|t|>n_k} M^{-1}(|x_k(t)|) dt &> \varepsilon_0, \\ \int_{|t|>n_{k+1}} M^{-1}(|x_k(t)|) dt &< \frac{\varepsilon_0}{2}. \end{aligned}$$

So for any  $k_1, k_2$  (we assume  $k_1 > k_2$  without loss of generality),

$$\begin{aligned} &\int_{-\infty}^{+\infty} M^{-1}(|x_{k_1}(t) - x_{k_2}(t)|) dt \\ &\geq \int_{-\infty}^{+\infty} |M^{-1}(|x_{k_1}(t)|) - M^{-1}(|x_{k_2}(t)|)| dt \\ &\geq \int_{|t|>n_{k_1}} |M^{-1}(|x_{k_1}(t)|) - M^{-1}(|x_{k_2}(t)|)| dt \\ &\geq \int_{|t|>n_{k_1}} M^{-1}(|x_{k_1}(t)|) dt - \int_{|t|>n_{k_1}} M^{-1}(|x_{k_2}(t)|) dt \\ &> \varepsilon_0 - \frac{\varepsilon_0}{2} = \frac{\varepsilon_0}{2}, \end{aligned}$$

i.e.,  $\|x_{k_1} - x_{k_2}\|_{(M^{-1})} > \frac{\varepsilon_0}{2}$ , so  $\{x_k\}_1^\infty \subset F$  has no convergent sequence in  $L^{(M^{-1})}(-\infty, +\infty)$ , This contradicts the sequential compactness of  $F$ .

(iii)  $\forall \varepsilon > 0$ , by (ii),  $\exists N \in \mathbb{N}$ , such that

$$\int_{|t|\geq N} M^{-1}(|x(t)|) dt < \frac{\varepsilon}{4}$$

for every  $x(t) \in F$ . So for all  $x(t) \in F$  and  $|h| < 1$ , we have

$$\begin{aligned} &\int_{|t|\geq N+1} M^{-1}(|x(t+h) - x(t)|) dt \\ &\leq \int_{|t|\geq N+1} M^{-1}(|x(t+h)| + |x(t)|) dt \\ &\leq \int_{|t|\geq N+1} M^{-1}(|x(t+h)|) dt + \int_{|t|\geq N+1} M^{-1}(|x(t)|) dt \\ &< \frac{\varepsilon}{4} + \frac{\varepsilon}{4} = \frac{\varepsilon}{2}. \end{aligned}$$

For  $\int_{|t| \leq N+1} M^{-1}[|x(t+h) - x(t)|] dt$ , by Theorem 2.1, we know that there exists  $\delta_1 > 0$  such that for  $|h| < \delta_1$ ,

$$\sup_{x \in F} \int_{|t| \leq N+1} M^{-1}[|x(t+h) - x(t)|] dt < \frac{\varepsilon}{2}.$$

Choose  $\delta = \min\{1, \delta_1\}$ ; then for all  $x \in F$  and  $|h| < \delta$ ,

$$\begin{aligned} & \int_{|t| < \infty} M^{-1}[|x(t+h) - x(t)|] dt \\ &= \int_{|t| \leq N+1} M^{-1}[|x(t+h) - x(t)|] dt + \int_{|t| > N+1} M^{-1}[|x(t+h) - x(t)|] dt \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon, \end{aligned}$$

i.e.,  $\sup_{x \in F} \int_{(-\infty, +\infty)} M^{-1}[|x(t+h) - x(t)|] dt < \varepsilon$ . □

A natural consequence is given by:

**Corollary 2.2** (Tsuji [9]). *For  $L^p(-\infty, +\infty)$  ( $0 < p < 1$ ) with  $F$ -norm  $\|x\|_p = \int_a^b |x(t)|^p dt$ ,  $A \subset L^p(-\infty, +\infty)$  is sequentially compact if and only if*

- (i)  $\exists c < \infty$ , such that  $\sup_{x \in A} \int_{-\infty}^{+\infty} |x(t)|^p dt \leq c$ ,
- (ii)  $\lim_{N \rightarrow \infty} [\sup_{x \in A} \int_{|t| \geq N} |x(t)|^p dt] = 0$ ,
- (iii)  $\forall \varepsilon > 0, \exists \delta > 0$  such that for  $|h| < \delta$ ,

$$\sup_{x \in A} \int_{-\infty}^{+\infty} |x(t+h) - x(t)|^p dt < \varepsilon.$$

### §3. Krasnoselskii criterion for sequential compactness in $L^{(M^{-1})}$

**Definition 3.1.**  $S \subset L^{(M^{-1})}(G)$  is said to be *sequentially compact in measure*, if for any  $\{f_n\}_{n=1}^{\infty} \subset S$  there exists a sequence  $\{f_{n_i}\}_{i=1}^{\infty}$  and  $f_0 \in L^{(M^{-1})}(G)$  such that  $f_{n_i}(x) \xrightarrow{m} f_0(x)$ .

**Definition 3.2.**  $S \subset L^{(M^{-1})}(G)$  is said to be *equi-absolutely continuous in  $F$ -norm*, if  $\forall \varepsilon > 0, \exists \delta > 0$ , such that for  $e \subset G$  and  $\mu(e) < \delta$ ,

$$\sup_{f \in S} \|f \cdot \chi_e\|_{(M^{-1})} < \varepsilon.$$

**Theorem 3.1.** *Let  $M$  be an Orlicz function and let  $\mu(G) < \infty$ . Then  $S \subset L^{(M^{-1})}(G)$  is sequentially compact iff*

- (i)  $S$  is sequentially compact in measure;
- (ii)  $S$  has an equi-absolutely continuous  $F$ -norm.

**PROOF:** *Sufficiency.* For any  $\{f_n\}_{n=1}^\infty \subset S$ , by (i) we know there exists a subsequence, we still denote it by  $\{f_n\}_{n=1}^\infty$ , and  $f_0 \in L^{(M^{-1})}(G)$ , such that  $f_n \xrightarrow{m} f_0$ . Next we prove that the subsequence  $\{f_n\}_{n=1}^\infty$  is a Cauchy sequence.

By (ii),  $\forall \varepsilon > 0, \exists \delta > 0$ , for  $\mu(e) < \delta$ ,

$$\sup_{n \geq 1} \|f_n \chi_e\|_{(M^{-1})} < \frac{\varepsilon}{4}.$$

Choose  $\varepsilon' < \varepsilon$ , such that  $M^{-1}(\varepsilon')\mu(G) < \frac{\varepsilon}{2}$ . Denote  $G_{n,m} = \{x \in G : |f_n(x) - f_m(x)| > \varepsilon'\}$ . By  $f_n \xrightarrow{m} f_0$ , we know  $\lim_{n,m \rightarrow \infty} \mu(G_{n,m}) = 0$ . So for  $\delta > 0, \exists n_0 \in \mathbb{N}$ , for all  $n, m > n_0, \mu(G_{n,m}) < \delta$ . Therefore

$$\begin{aligned} & \|f_n - f_m\|_{(M^{-1})} \\ &= \int_G M^{-1}(|f_n(t) - f_m(t)|) dt \\ &= \int_G M^{-1}(|(f_n - f_m)(t)\chi_{G_{n,m}}(t) + (f_n - f_m)(t)\chi_{G \setminus G_{n,m}}(t)|) dt \\ &\leq \int_G M^{-1}(|(f_n - f_m)(t)\chi_{G_{n,m}}(t)|) dt + \\ &\quad + \int_G M^{-1}(|(f_n - f_m)(t)\chi_{G \setminus G_{n,m}}(t)|) dt \\ &= \|(f_n - f_m)\chi_{G_{n,m}}\|_{(M^{-1})} + \int_{G \setminus G_{n,m}} M^{-1}(|(f_n - f_m)(t)|) dt \\ &< \frac{\varepsilon}{2} + M^{-1}(\varepsilon')\mu(G) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

By the completeness of  $L^{(M^{-1})}(G)$ , we know  $\{f_n\}_{n=1}^\infty$  converges in  $L^{(M^{-1})}(G)$ .

*Necessity.* (i). Let  $S \subset L^{(M^{-1})}(G)$  be a sequentially compact set. For any  $\{f_n\}_{n=1}^\infty \subset S$  there exists a subsequence, we still denote it by  $\{f_n\}_{n=1}^\infty$ , and  $f_0 \in L^{(M^{-1})}(G)$  such that  $\|f_n - f_0\|_{(M^{-1})} \rightarrow 0$  ( $n \rightarrow \infty$ ), i.e.,  $\int_G M^{-1}(|f_n(t) - f_0(t)|) dt \rightarrow 0$  ( $n \rightarrow \infty$ ). So  $f_n \xrightarrow{m} f_0$ .

(ii). Let  $S$  be a sequentially compact set. Then  $S$  is a complete bounded set. So, for any  $\varepsilon > 0$  there exists a finite  $\frac{\varepsilon}{2}$ -net of  $S$   $\{g_1, g_2, \dots, g_N\}$  such that  $S \subset \bigcup_{i=1}^N B(g_i, \frac{\varepsilon}{2})$ , where  $B(g_i, \frac{\varepsilon}{2}) = \{f \in L^{(M^{-1})}(G) : \|f - g_i\|_{(M^{-1})} < \frac{\varepsilon}{2}\}$ . Clearly,

for  $\frac{\varepsilon}{2} > 0$ , there is  $\delta > 0$  such that for  $\mu(e) < \delta$  we have  $\int_e M^{-1}(|g_i(t)|) dt < \frac{\varepsilon}{2}$ ,  $i = 1, 2, \dots, N$ . For any  $f \in S$  there exists  $g_{i_0}$  such that  $f \in B(g_{i_0}, \frac{\varepsilon}{2})$ . For  $e \subset G$ ,  $\mu(e) < \delta$ , we have

$$\begin{aligned} \|f \cdot \chi_e\|_{(M^{-1})} &= \int_e M^{-1}(|f(t)|) dt \\ &= \int_e M^{-1}(|f(t) - g_{i_0}(t)|) dt + \int_e M^{-1}(|g_{i_0}(t)|) dt \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

Thus for  $e \subset G$  with  $\mu(e) < \delta$  we have  $\sup_{f \in S} \|f \cdot \chi_e\|_{(M^{-1})} < \varepsilon$ . □

**Corollary 3.1.** For  $L^p[a, b]$  ( $0 < p < 1$ ) with  $F$ -norm  $\|x\|_p = \int_a^b |x(t)|^p dt$ ,  $A \subset L^p[a, b]$  is sequentially compact if and only if

- (i)  $A$  is sequentially compact in measure,
- (ii)  $A$  is equi-absolutely continuous in  $\|\cdot\|_p$ .

**Acknowledgment.** Author would like to thank the referees for reading this paper and correcting some errors.

#### REFERENCES

- [1] Rao M.M., Ren Z.D., *Theory of Orlicz Spaces*, Marcel Dekker, New York, 1991.
- [2] Maligranda L., *Orlicz Spaces and Interpolation*, Seminars in Mathematics 5, Univ. Estadual de Campinas, Capinas SP Brasil, 1989.
- [3] Rao M.M., Ren Z.D., *Applications of Orlicz Spaces*, Marcel Dekker, New York, 2001, to appear.
- [4] Riesz F., *Untersuchungen über Systeme integrierbarer Funktionen*, Math. Ann. **69** (1910), 449–497.
- [5] Kantorovich L.V., Akilov G.P., *Functional Analysis*, Pergamon Press, New York, 1982.
- [6] Krasnosel'skii M.A., Rutickii Ya.B., *Convex Functions and Orlicz Space*, Noordhoff Ltd. Groningen, 1961.
- [7] Takahashi T., *On the compactness of the function-set by the convergence in mean of general type*, Studia Math. **5** (1935), 141–150.
- [8] Gribanov Yu., *Nonlinear operators in Orlicz spaces* (in Russian), Uč. zap. kazansk. Unta **115** (7) (1955).
- [9] Tsuji M., *On the compactness of space  $L^p$  ( $p > 0$ ) and its application to integral equation*, Mathematical Institute, Tokyo University, 1951.

DEPARTMENT OF MATHEMATICS, SUZHOU UNIVERSITY, SUZHOU, 215006, P.R. CHINA

(Received February 24, 2001, revised September 15, 2001)