

Relative normality and product spaces

TAKAO HOSHINA, RYOKEN SOKEI

Abstract. Arhangel'skiĭ defines in [Topology Appl. 70 (1996), 87–99], as one of various notions on relative topological properties, strong normality of A in X for a subspace A of a topological space X , and shows that this is equivalent to normality of X_A , where X_A denotes the space obtained from X by making each point of $X \setminus A$ isolated. In this paper we investigate for a space X , its subspace A and a space Y the normality of the product $X_A \times Y$ in connection with the normality of $(X \times Y)_{(A \times Y)}$. The cases for paracompactness, more generally, for γ -paracompactness will also be discussed for $X_A \times Y$. As an application, we prove that for a metric space X with $A \subset X$ and a countably paracompact normal space Y , $X_A \times Y$ is normal if and only if $X_A \times Y$ is countably paracompact.

Keywords: strongly normal in, normal, γ -paracompact, product spaces, weak C -embedding

Classification: Primary 54B10; Secondary 54B05, 54C20, 54C45, 54D15, 54D20

1. Introduction

Throughout this paper all spaces are assumed to be Hausdorff. Let γ denote an infinite cardinal, and \mathbb{N} the set of natural numbers.

Let X be a space and A a subspace of X .

As is known, A is said to be C^* -embedded (respectively C -embedded) in X if every bounded real-valued (respectively real-valued) continuous function on A can be extended to a continuous function over X .

Next we recall some relative topological properties in Arhangel'skiĭ [2]. We say that A is *strongly normal in* X if for every pair E, F of disjoint closed subsets of A there exist disjoint open subsets U and V of X such that $E \subset U$ and $F \subset V$. The subspace A is *weakly C -embedded* in X if for every real-valued continuous function f on A there exists a real-valued function on X which is an extension of f and continuous at each point of Y .

For a space X and a subspace A of X let X_A denote the space obtained from the space X , with the topology generated by $\{U \mid U \text{ is open in } X \text{ or } U \subset X \setminus A\}$. Hence A is a closed subspace of X_A and points in $X \setminus A$ are isolated. As is seen in [2], the space X_A is often useful to describe several relative topological properties. Indeed, the following are shown in [2]: (1) X_A is normal if and only if A is strongly normal in X if and only if A is normal itself, and is weakly C -embedded in X , (2) A is weakly C -embedded in X if and only if A is C^* -embedded in X_A .

On the other hand, in a joint paper [9] of the first author with Yamazaki the notion of weak C -embedding was characterized by extending disjoint cozero-sets of a subspace to disjoint open sets of the whole space. And it was applied there for a space X , a subspace A of X and a space Y to describe weak C -embedding of $A \times Y$ in the product $X_A \times Y$; actually, it was shown that if Y is compact Hausdorff, $A \times Y$ is C^* -embedded in $X_A \times Y$ if and only if $A \times Y$ is C^* -embedded in $(X \times Y)_{(A \times Y)}$, that is, $A \times Y$ is weakly C -embedded in $X \times Y$. Being motivated by this result, our main concern in this paper is to study normality of the product $X_A \times Y$ in relation to normality of $(X \times Y)_{(A \times Y)}$ (or, equivalently, strong normality of $A \times Y$ in $X \times Y$). Namely we prove

Theorem 1.1. *For a space X , a subspace A of X and a space Y , the product $X_A \times Y$ is normal if and only if $(X \times Y)_{(A \times Y)}$ is normal and the following condition (*) holds:*

- (*) *for every closed subset E of $X_A \times Y$ disjoint from $A \times Y$ there exists an open subset U of $X_A \times Y$ such that $E \subset U$ and $\overline{U} \cap (A \times Y) = \emptyset$.*

As a corollary to this result we have that for a space X , a subspace A of X and a compact Hausdorff space Y , $X_A \times Y$ is normal if and only if $(X \times Y)_{(A \times Y)}$ is normal. Moreover, using condition (*) above we prove analogous results for γ -collectionwise normality or γ -paracompactness. In particular, the case $\gamma = \omega$ is applied to obtain further the following theorem; putting $A = X$, we have the well-known theorem due to Morita [14] (for the proof see [10]) and Rudin and Starbird [16].

Theorem 1.2. *Let X be a metric space, A a subspace of X and Y a normal and countably paracompact space. Then $X_A \times Y$ is normal if and only if $X_A \times Y$ is countably paracompact.*

For undefined notation and terminology see Engelking's book [6].

2. Preliminaries

The following theorem due to Arhangel'skiĭ [2] mentioned in the introduction is useful.

Theorem 2.1 ([2]). *For a subspace A of a space X , the following statements are equivalent:*

- (1) X_A is normal,
- (2) A is strongly normal in X ,
- (3) A is normal and A is weakly C -embedded in X .

Weak C -embedding was characterized in [9] as follows.

Theorem 2.2 ([9]). *Let A be a subspace of a space X . Then A is weakly C -embedded in X if and only if for every pair G_0, G_1 of disjoint cozero-sets in A there exist disjoint open subsets H_0, H_1 of X such that $G_i \subset H_i$ ($i = 0, 1$).*

By this result we see that if either A is dense in X or A is z -embedded in X , then A is weakly C -embedded in X ([5], [9]); a subspace A of a space X is said to be z -embedded in X if every zero-set Z of A can be written as $Z = Z' \cap A$ with a zero-set Z' of X . It is known that every cozero-set of a space or a Lindelöf subspace of a Tychonoff space is z -embedded. Also, observe the following implications:

$$C^* \text{-embedding} \Rightarrow z\text{-embedding} \Rightarrow \text{weak } C\text{-embedding}.$$

The next two results show when a subspace $A \times Y$ is weakly C -embedded in $X \times Y$ for a space X , a subspace A of X and a metric space Y . The first one is essentially due to Kodama [11].

Theorem 2.3 ([11]). *Let X be a normal space, A a closed subspace of X and Y a metric space. If $A \times Y$ is normal and countably paracompact, then $A \times Y$ is z -embedded in $X \times Y$, hence, weakly C -embedded in $X \times Y$.*

In case $A \times Y$ is not assumed to be normal, we have the following.

Theorem 2.4. *Let A be an arbitrary subspace of a hereditarily normal space X , and Y a metric space. Then $A \times Y$ is weakly C -embedded in $X \times Y$.*

PROOF: We show that any two disjoint open sets of $A \times Y$ are separated by disjoint open sets of $X \times Y$, which implies weak C -embedding of $A \times Y$ in $X \times Y$ by Theorem 2.2. Let G_0 and G_1 be disjoint open sets of $A \times Y$. Let $\mathcal{B} = \bigcup_{n \in \mathbb{N}} \mathcal{B}_n$ be a σ -locally finite open base for Y , where each \mathcal{B}_n is locally finite. Let $\Lambda_n = \{\mathcal{B}_{n\lambda} \mid \lambda \in \Lambda_n\}$. Define for $n \in \mathbb{N}$ and $\lambda \in \Lambda_n$

$$H_{n\lambda}^0 = \bigcup \{O \mid O \text{ is open in } A, O \times \overline{B_{n\lambda}} \subset G_0\}.$$

Then $H_{n\lambda}^0$ and $p_A((A \times \overline{B_{n\lambda}}) \cap G_1)$ are disjoint open subsets of A . Since X is hereditarily normal, there exists an open set $W_{n\lambda}^0$ of X such that

$$H_{n\lambda}^0 \subset W_{n\lambda}^0, \quad \overline{W_{n\lambda}^0} \cap p_A((A \times \overline{B_{n\lambda}}) \cap G_1) = \emptyset.$$

For each $n \in \mathbb{N}$ let us put $U_n^0 = \bigcup \{W_{n\lambda}^0 \times B_{n\lambda} \mid \lambda \in \Lambda_n\}$. Then U_n^0 is an open set of $X \times Y$ and we have $G_0 \subset \bigcup_{n \in \mathbb{N}} U_n^0$ and $\overline{U_n^0} \cap G_1 = \emptyset$ for every $n \in \mathbb{N}$. Similarly, we can find an open set U_n^1 of $X \times Y$ for each $n \in \mathbb{N}$ so that $G_1 \subset \bigcup_{n \in \mathbb{N}} U_n^1$ and $\overline{U_n^1} \cap G_0 = \emptyset$ for every $n \in \mathbb{N}$. Hence, as is well-known, G_0 and G_1 are separated by open sets of $X \times Y$. This completes the proof. \square

It was shown in [9] that every subspace of a space X is weakly C -embedded in X if and only if X is hereditarily normal.

In connection with Theorems 2.3 and 2.4, let us observe the following two examples.

Example 2.5. (1) (Michael [12]) Let \mathbb{R} , \mathbb{Q} and \mathbb{P} be the real line, the set of rationals and the set of irrationals, respectively. Then $\mathbb{R}_{\mathbb{Q}}$ is known as the Michael line, and it is hereditarily normal. Since $\mathbb{Q} \times \mathbb{P}$ is Lindelöf, it is z -embedded in $\mathbb{R}_{\mathbb{Q}} \times \mathbb{P}$, but is not C^* -embedded as was shown by Morita [15].

(2) (Vaughan [17]) Let $D(\omega_1)$ denote the set ω_1 with the discrete topology. Let $\widehat{D}(\omega_1)$ denote the space obtained from the space $\omega_1 + 1$ with the usual order topology by letting all points except ω_1 be isolated. That is, $\widehat{D}(\omega_1) = (\omega_1 + 1)_{\{\omega_1\}}$.

Let $X = \square_{\omega} \widehat{D}(\omega_1)$ denote the box product of countably many copies of $\widehat{D}(\omega_1)$, and $Y = D(\omega_1)^{\omega}$ denote the usual product of countably many copies of $D(\omega_1)$.

Then X is hereditarily paracompact and Y is metrizable. Put

$$A = X \setminus Y, \quad \Delta(Y) = \{ \langle x, x \rangle \mid x \in Y \}.$$

Then $A \times Y$ and $\Delta(Y)$ are disjoint closed sets of $X \times Y$ and cannot be separated by open sets, which shows $X \times Y$ is not normal ([17]).

By Theorem 2.4 we see that $A \times Y$ is weakly C -embedded in $X \times Y$. Since A contains a closed subset homeomorphic to X , $A \times Y$ is not normal. Hence, in view of Theorem 2.3, it may be of interest to see whether $A \times Y$ is z -embedded in $X \times Y$, but this is unknown to the authors. However, we can show further that $A \times Y$ is not C^* -embedded in $X \times Y$. To prove this, first note that $Y \cong$ (is homeomorphic to) Y^2 . Hence, if we show the fact below, by the same argument of Morita [15] we can conclude that $A \times Y$ is not C^* -embedded in $X \times Y$.

Fact. $\Delta(Y)$ is a zero-set of $X \times Y$.

PROOF: Since the box topology is stronger than the usual topology, it suffices to show that $\Delta(Y)$ is a zero-set of $\widehat{D}(\omega_1)^{\omega} \times D(\omega_1)^{\omega}$.

For each point $\langle x, y \rangle \in \widehat{D}(\omega_1)^{\omega} \times D(\omega_1)^{\omega} \setminus \Delta(Y)$, define

$$n(x, y) = \min \{ k \mid x_k \neq y_k \}.$$

Put

$$H_m = \{ \langle x, y \rangle \in \widehat{D}(\omega_1)^{\omega} \times D(\omega_1)^{\omega} \setminus \Delta(Y) \mid n(x, y) = m \}.$$

Then we have

$$\begin{aligned} \widehat{D}(\omega_1)^{\omega} \times D(\omega_1)^{\omega} \setminus \Delta(Y) &= \bigcup_{m \in \mathbb{N}} H_m, \\ m \neq m' &\Rightarrow H_m \cap H_{m'} = \emptyset. \end{aligned}$$

Claim. H_m is an open and closed subset of $\widehat{D}(\omega_1)^{\omega} \times D(\omega_1)^{\omega}$.

PROOF OF CLAIM: Let $\langle x, y \rangle \in H_m$. Since $n(x, y) = m$, we have $x_1 = y_1, \dots, x_{m-1} = y_{m-1} < \omega_1$.

Case (i). $x_m > y_m$. Put

$$U = \{x_1\} \times \cdots \times \{x_{m-1}\} \times (y_m, \omega_1] \times \widehat{D}(\omega_1) \times \cdots, \\ V = \{y_1\} \times \cdots \times \{y_{m-1}\} \times \{y_m\} \times D(\omega_1) \times \cdots.$$

Then $\langle x, y \rangle \in U \times V \subset H_m$.

Case (ii). $x_m < y_m$. Put

$$U = \{x_1\} \times \cdots \times \{x_{m-1}\} \times \{x_m\} \times \widehat{D}(\omega_1) \times \cdots, \\ V = \{y_1\} \times \cdots \times \{y_{m-1}\} \times \{y_m\} \times D(\omega_1) \times \cdots.$$

Then $\langle x, y \rangle \in U \times V \subset H_m$.

Hence, in either case H_m is open in $\widehat{D}(\omega_1)^\omega \times D(\omega_1)^\omega$.

For each $\langle y, y \rangle \in \Delta(Y)$, put

$$U = \{y_1\} \times \cdots \times \{y_m\} \times \widehat{D}(\omega_1) \times \cdots, \\ V = \{y_1\} \times \cdots \times \{y_m\} \times D(\omega_1) \times \cdots.$$

Then $(U \times V) \cap H_m = \emptyset$. Hence, $\Delta(Y) \cap \overline{H_m} = \emptyset$, which shows that H_m is closed in $X \times Y$.

It follows that H_m is a cozero-set, therefore, $\bigcup_{m \in \mathbb{N}} H_m$ is a cozero-set of $\widehat{D}(\omega_1)^\omega \times D(\omega_1)^\omega$. Hence $\Delta(Y)$ is a zero-set of $X \times Y$. This completes the proof. \square

3. Proof of Theorem 1.1

First we prove

Lemma 3.1. *Let X be a space, A a subspace of X and Y a space. If $X_A \times Y$ is normal, then $(X \times Y)_{(A \times Y)}$ is normal.*

PROOF: Let E and F be disjoint closed subsets of $A \times Y$. Then they are closed also in $X_A \times Y$ and disjoint. Hence, there exist disjoint open subsets U and V of $X_A \times Y$ such that $E \subset U$ and $F \subset V$. Define $U' = \text{Int}_{(X \times Y)} U$ and $V' = \text{Int}_{(X \times Y)} V$, where $\text{Int}_Z W$ denotes the interior of W in the space Z . Then U' and V' are disjoint open in $X \times Y$ and so in $(X \times Y)_{(A \times Y)}$, and we have $E \subset U'$ and $F \subset V'$. Hence, $A \times Y$ is strongly normal in $X \times Y$. Hence by Theorem 2.1 $(X \times Y)_{(A \times Y)}$ is normal. This completes the proof. \square

Remark. $(\mathbb{R} \times \mathbb{P})_{(\mathbb{Q} \times \mathbb{P})}$ is normal, but $\mathbb{R}_{\mathbb{Q}} \times \mathbb{P}$ is not normal. The converse of the lemma, therefore, need not hold.

PROOF OF THEOREM 1.1: From Lemma 3.1 the “only if” part easily follows. To prove the “if” part, assume that $(X \times Y)_{(A \times Y)}$ is normal and condition (*) holds. Let E, F be a pair of disjoint closed subsets of $X_A \times Y$. Since $A \times Y$ is strongly normal in $X \times Y$ by Theorem 2.1, there exist disjoint open subsets U and V of $X \times Y$ such that $E \cap (A \times Y) \subset U$ and $F \cap (A \times Y) \subset V$. Put $D = (E \setminus U) \cup (F \setminus V)$. Then D is a closed subset of $X \times Y$ and $D \cap (A \times Y) = \emptyset$. Then by (*), there exists an open subset W of $X_A \times Y$ such that $A \times Y \subset W$ and $\overline{W} \cap D = \emptyset$.

Put $U_1 = U \cap W$ and $V_1 = V \cap W$. Then we have

$$(A \times Y) \cap E \subset U_1, \overline{U_1} \cap F = \emptyset, \text{ and } (A \times Y) \cap F \subset V_1, \overline{V_1} \cap E = \emptyset.$$

Then $E \setminus U_1$ and $F \setminus V_1$ are disjoint closed subsets of $(X_A \setminus A) \times Y$. Since $(X_A \setminus A) \times Y$ is normal, there exist disjoint open subsets U_2 and V_2 of $(X_A \setminus A) \times Y$ such that $E \setminus U_1 \subset U_2$ and $F \setminus V_1 \subset V_2$. Therefore, $U_1 \cup (U_2 \setminus \overline{V_1})$ and $V_1 \cup (V_2 \setminus \overline{U_1})$ are disjoint open subsets of $X_A \times Y$, which satisfy $E \subset U_1 \cup (U_2 \setminus \overline{V_1})$ and $F \subset V_1 \cup (V_2 \setminus \overline{U_1})$. Hence $X_A \times Y$ is normal. This completes the proof. \square

The following is proved in Burke and Pol [4].

Theorem 3.2 ([4]). *Let A and X be subsets of \mathbb{R} with $A \subset X$ and let Y be a metric space. Then $X_A \times Y$ is normal if and only if condition (*) holds.*

Since $X \times Y$ is a metric space, $(X \times Y)_{(A \times Y)}$ is normal. Therefore, this theorem immediately follows from Theorem 1.1.

The following result was formulated in [9] without proof.

Theorem 3.3 ([9]). *Let A be a subset of a space X and Y be a compact Hausdorff space. Then $X_A \times Y$ is normal if and only if $(X \times Y)_{(A \times Y)}$ is normal.*

PROOF: Since the projection $p_{X_A}: X_A \times Y \rightarrow X_A$ is a closed map, condition (*) in Theorem 1.1 is easily satisfied. Hence the theorem follows. \square

Recall that a space X is γ -collectionwise normal if for every discrete collection $\{E_\alpha \mid \alpha < \gamma\}$ of closed subsets there exists a disjoint collection $\{G_\alpha \mid \alpha < \gamma\}$ of open subsets such that $E_\alpha \subset G_\alpha$ for each $\alpha < \gamma$.

A subspace A of a space X is said to be strongly γ -collectionwise normal in X if for every discrete collection $\{E_\alpha \mid \alpha < \gamma\}$ of closed subsets of A there is a disjoint collection $\{U_\alpha \mid \alpha < \gamma\}$ of open subsets of X such that $E_\alpha \subset U_\alpha$ for each $\alpha < \gamma$ ([9]).

It was proved in [9] that X_A is γ -collectionwise normal if and only if A is strongly γ -collectionwise normal in X . With this result similarly to Theorem 1.1 we can prove the following.

Theorem 3.4. For a space X , a subspace A of X and a space Y , $X_A \times Y$ is γ -collectionwise normal if and only if $(X \times Y)_{(A \times Y)}$ is γ -collectionwise normal and condition $(*)$ in Theorem 1.1 holds.

A space X is γ -paracompact if every open cover of X of cardinality not greater than γ has a locally finite open refinement.

Theorem 3.5. If $X_A \times Y$ is γ -paracompact, then $(X \times Y)_{(A \times Y)}$ is γ -paracompact. Furthermore, if $X_A \times Y$ satisfies condition $(*)$ in Theorem 1.1, then the converse holds.

PROOF: Assume $X_A \times Y$ is γ -paracompact. Let \mathcal{U} be an open cover of $(X \times Y)_{(A \times Y)}$ of cardinality not greater than γ . Put

$$\mathcal{U}' = \{U \in \mathcal{U} \mid U \cap (A \times Y) \neq \emptyset\}.$$

Then $\bigcup\{\text{Int}_{(X \times Y)} U \mid U \in \mathcal{U}'\} \supset A \times Y$. Hence $\{X_A \times Y \setminus A \times Y\} \cup \mathcal{U}'$ is an open cover of $X_A \times Y$ of cardinality not greater than γ . Since $X_A \times Y$ is γ -paracompact, there exists a locally finite open cover \mathcal{V} of $X_A \times Y$ which refines \mathcal{U} . Put $\mathcal{V}' = \{V \in \mathcal{V} \mid V \cap (A \times Y) \neq \emptyset\}$. Then the collection

$$\mathcal{V}' \cup \{\{ \langle x, y \rangle \} \mid \langle x, y \rangle \notin \bigcup \mathcal{V}'\}$$

is a locally finite open cover of $(X \times Y)_{(A \times Y)}$ and refines \mathcal{U} . Hence $(X \times Y)_{(A \times Y)}$ is γ -paracompact.

To prove the converse under $(*)$, assume that $(X \times Y)_{(A \times Y)}$ is γ -paracompact and $(*)$ holds. Let \mathcal{U} be an open cover of $X_A \times Y$ of cardinality not greater than γ . Then \mathcal{U} is an open cover of $(X \times Y)_{(A \times Y)}$ as well. By assumption there exists a locally finite open cover \mathcal{V} of $(X \times Y)_{(A \times Y)}$ refining \mathcal{U} . Put

$$G = \{ \langle x, y \rangle \in X \times Y \mid \mathcal{V} \text{ is locally finite at } \langle x, y \rangle \text{ in the product } X \times Y \}.$$

Then G is open in $X \times Y$ and $G \supset A \times Y$. Put $\mathcal{V}' = \{G \cap \text{Int}_{(X \times Y)} V \mid V \in \mathcal{V}\}$. Then we have $\bigcup \mathcal{V}' \supset A \times Y$, and \mathcal{V}' refines \mathcal{U} and is locally finite at each $\langle x, y \rangle \in \bigcup \mathcal{V}'$ in $X \times Y$. By $(*)$ there exist open subsets O_1 and O_2 in $X_A \times Y$ such that

$$A \times Y \subset O_1 \subset \overline{O_1} \subset O_2 \subset \overline{O_2} \subset \bigcup \mathcal{V}'.$$

For every $x \in X \setminus A$, let \mathcal{P}_x be a locally finite open cover of Y such that the collection $\{\{x\} \times P \mid P \in \mathcal{P}_x\}$ refines \mathcal{U} . Then the collection

$$\{(\{x\} \times P) \setminus \overline{O_1} \mid x \in X \setminus Y, P \in \mathcal{P}_x\} \cup \{V \cap O_2 \mid V \in \mathcal{V}'\}$$

is a locally finite open cover of $X_A \times Y$ which refines \mathcal{U} . Thus, $X_A \times Y$ is γ -paracompact. This completes the proof. □

4. Proof of Theorem 1.2

First we prove

Theorem 4.1. *Let A be a subset of a space X and Y a space. Suppose that the product $A \times Y$ is γ -paracompact. If $X_A \times Y$ is normal, then $X_A \times Y$ is γ -paracompact.*

PROOF: Assume that $X_A \times Y$ is normal. Then $(X \times Y)_{(A \times Y)}$ is normal by Theorem 1.1. Hence $A \times Y$ is normal and weakly C -embedded in $X \times Y$ by Theorem 2.1. Since $A \times Y$ is γ -paracompact, by [9, Lemma 4.6] $(X \times Y)_{(A \times Y)}$ is γ -paracompact. Since $X_A \times Y$ satisfies $(*)$, $X_A \times Y$ is γ -paracompact by Theorem 3.5. This completes the proof. \square

Corollary 4.2. *Let A be a subset of a space X and Y a space. Suppose that the product $A \times Y$ is countably paracompact. If $X_A \times Y$ is normal, then $X_A \times Y$ is countably paracompact.*

PROOF OF THEOREM 1.2: Let A be a subspace of a metric space X , and Y a normal and countably paracompact space. To prove the “only if” part, assume $X_A \times Y$ is normal. Since $A \times Y$ is closed in $X_A \times Y$, $A \times Y$ is also normal. Hence by Morita, Rudin-Starbird’s theorem ([14], [16]), $A \times Y$ is countably paracompact. Hence $X_A \times Y$ is countably paracompact by Corollary 4.2.

To prove the converse, assume that $X_A \times Y$ is countably paracompact. Then similarly to above we have that $A \times Y$ is countably paracompact and normal. Then by [11] $A \times Y$ is z -embedded in $A \times \beta Y$, where βY is the Čech-Stone compactification of Y . Since $X_A \times \beta Y$ is paracompact, $A \times \beta Y$ is C -embedded in $X_A \times \beta Y$. It follows that $A \times Y$ is z -embedded in $X_A \times Y$, and hence it is weakly C -embedded in $X_A \times Y$. This easily implies that $A \times Y$ is weakly C -embedded in $X \times Y$. Hence $(X \times Y)_{(A \times Y)}$ is normal.

We next show that property $(*)$ in Theorem 1.1 is satisfied. Let $\{\mathcal{B}_n\}$ be a sequence of locally finite open covers of X such that $\{\text{St}(x, \mathcal{B}_n) \mid n \in \mathbb{N}\}$ is a neighborhood base at each point x in X . Let $\mathcal{B}_n = \{B_{n\alpha} \mid \alpha \in \Omega_n\}$. Let us put

$$W(\alpha_1, \dots, \alpha_n) = \bigcap \{B_{i\alpha_i} \mid i = 1, \dots, n\}, \quad \text{for } \alpha_i \in \Omega_i; \quad i = 1, \dots, n.$$

To prove $(*)$, let E be a closed subset of $X_A \times Y$ such that $E \cap (A \times Y) = \emptyset$. Put

$$G(\alpha_1, \dots, \alpha_n) = \bigcup \{O \mid O \text{ is open in } Y, (W(\alpha_1, \dots, \alpha_n) \times O) \cap E = \emptyset\}.$$

Then we have

$$G(\alpha_1, \dots, \alpha_n) \subset G(\alpha_1, \dots, \alpha_n, \alpha_{n+1})$$

for $\alpha_i \in \Omega_i, i = 1, \dots, n, n + 1$, and

$$\{(W(\alpha_1, \dots, \alpha_n) \cap A) \times G(\alpha_1, \dots, \alpha_n) \mid \alpha_i \in \Omega_i, i = 1, \dots, n; n \in \mathbb{N}\}$$

covers $A \times Y$. Since $A \times Y$ is normal and countably paracompact, by Morita [13] (see [8]) there exists a cozero-set $U(\alpha_1, \dots, \alpha_n)$ of Y such that

$$U(\alpha_1, \dots, \alpha_n) \subset G(\alpha_1, \dots, \alpha_n)$$

and

$$\{(W(\alpha_1, \dots, \alpha_n) \cap A) \times U(\alpha_1, \dots, \alpha_n) \mid \alpha_i \in \Omega_i, i = 1, \dots, n; n \in \mathbb{N}\}$$

covers $A \times Y$. Put

$$L = \bigcup \{W(\alpha_1, \dots, \alpha_n) \times U(\alpha_1, \dots, \alpha_n) \mid \alpha_i \in \Omega_i, i = 1, \dots, n; n \in \mathbb{N}\}.$$

Then L is a cozero-set of $X \times Y$ and we have $L \supset A \times Y, L \cap E = \emptyset$. Since $X_A \times Y$ is countably paracompact, by [7] there exists an open subset H of $X \times Y$ such that $A \times Y \subset H \subset \overline{H} \subset L$. Hence $A \times Y$ and E are separated by open sets of $X_A \times Y$. This completes the proof of the theorem. \square

The proof of the “if” part of Theorem 1.1 yields further the following result which seems of interest in itself.

Theorem 4.3. *Let A be a subset of a metric space X and Y a normal and γ -paracompact space. Then $(X \times Y)_{(A \times Y)}$ is γ -paracompact if and only if $A \times Y$ is normal.*

PROOF: To prove the “if” part, assume that $A \times Y$ is normal. Since Y is normal and γ -paracompact, so is $A \times Y$. Hence $(A \times Y) \times I^\gamma$ is normal γ -paracompact, that is, $A \times I^\gamma \times Y$ is normal, where $I = [0, 1]$. Hence, as is shown in the proof of Theorem 1.2, $(X \times (I^\gamma \times Y))_{(A \times (I^\gamma \times Y))}$ is normal. Since $(X \times (I^\gamma \times Y))_{(A \times (I^\gamma \times Y))} \cong ((X \times Y) \times I^\gamma)_{((A \times Y) \times I^\gamma)}, ((X \times Y) \times I^\gamma)_{((A \times Y) \times I^\gamma)}$ is normal. Thus, by Theorem 3.3 $(X \times Y)_{(A \times Y)} \times I^\gamma$ is normal. Therefore, as is well-known, $(X \times Y)_{(A \times Y)}$ is γ -paracompact (see [6]). This completes the proof. \square

Example 4.4. The condition “ X is metric” cannot be excluded from Theorem 1.2. In fact, there exist compact spaces X and Y , and a subset A of X such that $A \times Y$ is normal and countably paracompact and $X_A \times Y$ is countably paracompact, but not normal. We use Bing’s example G [3]. Let $\mathcal{P}(\omega_1)$ be the power set of ω_1 and

$$X = \{f \mid f : \mathcal{P}(\omega_1) \longrightarrow \{0, 1\}\}.$$

For every $\alpha \in \omega_1$, let us define a function $f_\alpha : \mathcal{P} \longrightarrow \{0, 1\}$ for $P \in \mathcal{P}(\omega_1)$ by

$$f_\alpha(P) = \begin{cases} 1 & \text{if } \alpha \in P, \\ 0 & \text{if } \alpha \notin P. \end{cases}$$

Put $A = \{f_\alpha \mid \alpha < \omega_1\}$. Then Bing's example G is precisely the space X_A . It is well-known that X_A is normal and countably paracompact, but it is not ω_1 -collectionwise normal. Let Y be the one-point compactification of the discrete space of Card A . Since X_A is countably paracompact, $A \times Y$ is countably paracompact. Since A is $w(Y)$ -paracompact, $A \times Y$ is normal. However, since X_A is not ω_1 -collectionwise normal, by Alas [1] $X_A \times Y$ is not normal.

REFERENCES

- [1] Alas O.T., *On a characterization of collectionwise normality*, Canad. Math. Bull. **14** (1971), 13–15.
- [2] Arhangel'skiĭ A.V., *Relative topological properties and relative topological spaces*, Topology Appl. **70** (1996), 87–99.
- [3] Bing R.H., *Metrization of topological spaces*, Canad. Math. J. **5** (1951), 175–186.
- [4] Burke D.K., Pol R., *Products of Michael spaces and completely metrizable spaces*, Proc. Amer. Math. Soc. **129** (2000), 1535–1544.
- [5] Costantini C., Marccone A., *Extensions of functions which preserve the continuity on the original domain*, Topology Appl. **103** (2000), 131–153.
- [6] Engelking R., *General Topology*, Heldermann Verlag, Berlin, 1989.
- [7] Hoshina T., *Products of normal spaces with Lašnev spaces*, Fund. Math. **124** (1984), 143–153.
- [8] Hoshina T., *Normality of product spaces, II*, in K. Morita and J. Nagata, eds., Topics in General Topology, North-Holland, Amsterdam, 1989, pp. 121–160.
- [9] Hoshina T., Yamazaki K., *Weak C -embedding and P -embedding, and product spaces*, Topology Appl. **125** (2002), 233–247.
- [10] Ishii T., *On product spaces and product mappings*, J. Math. Soc. Japan **18** (1966), 166–181.
- [11] Kodama Y., *On subset theorems and the dimension of products*, Amer. J. Math. **106** (1969), 486–498.
- [12] Michael E., *The product of a normal space and a metric space need not be normal*, Bull. Amer. Math. Soc. **69** (1963), 375–376.
- [13] Morita K., *Products of normal spaces with metric spaces*, rm II, Sci. Rep. Tokyo Kyoiku Daigaku, Sect. A, **8** (1962), 87–92.
- [14] Morita K., *Note on products of normal spaces with metric spaces*, unpublished.
- [15] Morita K., *On the dimension of the product of topological spaces*, Tsukuba J. Math. **1** (1977), 1–6.
- [16] Rudin M.E., Starbird M., *Products with a metric factor*, General Topology Appl. **5** (1975), 235–348.
- [17] Vaughan J.E., *Non-normal products of ω_μ -metrizable spaces*, Proc. Amer. Math. Soc. **51** (1975), 203–208.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF TSUKUBA, TSUKUBA, IBARAKI 305-8571, JAPAN

E-mail: takaohsn@math.tsukuba.ac.jp

sokei@math.tsukuba.ac.jp

(Received March 3, 2003)