

On weighted spaces of functions harmonic in \mathbb{R}^n

A.I. PETROSYAN

Abstract. The paper establishes integral representation formulas in arbitrarily wide Banach spaces $b_\omega^p(\mathbb{R}^n)$ of functions harmonic in the whole \mathbb{R}^n .

Keywords: weighted spaces, harmonic functions, integral representation, isometry

Classification: 30H05, 46E15

1. Introduction

1.1. This paper extends the results of [1] related to arbitrarily wide spaces of functions harmonic in the unit ball B of \mathbb{R}^n to similar spaces of functions harmonic in the whole \mathbb{R}^n . Namely, the integral representation formulas in spaces $b_\omega^p(\mathbb{R}^n)$, which have natural definition, are obtained by exhausting \mathbb{R}^n by enlarging balls. Also, a representation connected with the natural isometry between $b_\omega^2(\mathbb{R}^n)$ and the space $L^2(S)$ is obtained, which is of an explicit form of integral operator along with its inversion.

The results of this paper are foregoing multidimensional similarities of the early results of M.M. Djrbashian [3], [4] (1945–1948) which in essence gave rise to the theory of $H^p(\alpha)$ spaces in the unit disc: the applied analytic apparatus allows to extend the results of [5] related to the one-dimensional case and holomorphic functions to functions harmonic in \mathbb{R}^n .

Note that in [2], the case of weighted spaces of functions analytical in the unit ball of \mathbb{C}^n is investigated.

1.2. We start with some notation which we use all over the paper.

$B = \{x \in \mathbb{R}^n: |x| < 1\}$ is the open unit ball in \mathbb{R}^n and $S = \{x \in \mathbb{R}^n: |x| = 1\}$ is its boundary, i.e. S is the unit sphere in \mathbb{R}^n ;

σ is the normalized surface-area measure over S , i.e. $\sigma(S) = 1$;

$\mathcal{H}_k(\mathbb{R}^n)$ is the set of all complex-valued homogeneous harmonic polynomials of degree k in \mathbb{R}^n ;

$\mathcal{H}_k(S)$ is the set of all spherical harmonics of degree k , i.e. all restrictions of functions from $\mathcal{H}(\mathbb{R}^n)$ to the sphere S ;

$Z_k(\eta, \zeta)$ is the zonal harmonic of degree k , i.e. $Z_k(\cdot, \zeta) \in \mathcal{H}_k(S)$ and $p(\zeta) = \int_S p(\eta) Z_k(\eta, \zeta) d\sigma(\eta)$ for all $p \in \mathcal{H}_k(S)$;

$P[f]$ is used for the Poisson integral of f :

$$(1) \quad P[f](x) = \int_S P(x, \zeta) f(\zeta) d\sigma(\zeta), \quad \text{where } P(x, \zeta) = \frac{1 - |x|^2}{|\zeta - x|^n}.$$

2. The case of the ball

We shall use the following definitions and statements from [1] related to the weighted spaces $b_\omega^p(B)$ in the unit ball.

As in [5], by Ω we denote the class of all functions $\omega(t)$ defined on $[0, 1]$ and such that $\omega(1) = \omega(1 - 0)$ and

- (i) $0 < \sqrt[\delta]{\omega} < \infty$ for any $\delta \in [0, 1]$;
- (ii) $\Delta_k \equiv \Delta_k(\omega) = - \int_0^1 t^k d\omega(t) \neq 0, \infty, \quad k = 0, 1, \dots$;
- (iii) $\liminf_{k \rightarrow \infty} \sqrt[k]{|\Delta_k|} \geq 1$.

Further, for a given $\omega \in \Omega$, we denote

$$d\mu_\omega(x) = -d\omega(r^2) d\sigma(\zeta),$$

where $x = r\zeta$ is the polar form of x , (i.e. $r = |x|, \zeta \in S$) and define $L_\omega^p(B)$ as the set of all $d\mu_\omega$ -measurable functions in B for which

$$\|u\|_{p,\omega} = \left\{ \int_B |u(x)|^p |d\mu_\omega(x)| \right\}^{1/p} < +\infty, \quad 1 \leq p < \infty.$$

By $b_\omega^p(B)$ we denote the subset of harmonic functions from $L_\omega^p(B)$.

Further, for a given $\omega \in \Omega$ we use the ω -kernel of the form

$$R_\omega(x, y) = \sum_{k=0}^{\infty} \Delta_k^{-1} Z_k(x, y),$$

where $Z_k(x, y)$ is the harmonic extension of the zonal harmonic Z_k by its both arguments. As it is proved in [1], for any function $u \in b_\omega^p(B)$ the following integral representation is true:

$$(2) \quad u(x) = \int_B u(y) R_\omega(x, y) d\mu_\omega(y), \quad x \in B.$$

3. The integral representation in \mathbb{R}^n

3.1. Let Ω^∞ denote the set of parameter-functions $\omega(t)$, which strictly decrease on the whole half-axis $[0, +\infty)$ and are such that $\omega(0) = 1$ and

$$\Delta_k^\infty(\omega) = - \int_0^{+\infty} t^k d\omega(t) < +\infty \text{ for any } k = 0, 1, \dots .$$

For a given $\omega \in \Omega^\infty$ we introduce the space $b_\omega^p(\mathbb{R}^n)$ as the set of all functions which are harmonic in \mathbb{R}^n and such that

$$\|u\|_{p,\omega} = \left\{ \int_{\mathbb{R}^n} |u(y)|^p d\mu_\omega(y) \right\}^{1/p} < +\infty, \quad 1 \leq p < +\infty,$$

where $d\mu_\omega(r\zeta) = -d\omega(r^2)d\sigma(\zeta)$. Let $L_\omega^p(\mathbb{R}^n)$ be the corresponding Lebesgue spaces. We shall deal with the following ω -kernel in \mathbb{R}^n :

$$(3) \quad R_\omega^\infty(x, y) = \sum_{k=0}^\infty \frac{Z_k(x, y)}{\Delta_k^\infty(\omega)} .$$

Lemma 1. *The right-hand side series in (3) is absolutely and uniformly convergent on any compact subset of $\mathbb{R}^n \times \mathbb{R}^n$, and hence $R_\omega^\infty(x, y)$ is harmonic in each of its variables in \mathbb{R}^n .*

PROOF: Let $x = r\zeta, y = \rho\eta$, where $\zeta, \eta \in S$. As the function $Z_k(x, y)$ is homogeneous in its both variables, we get

$$(4) \quad |Z_k(x, y)| = r^k \rho^k |Z_k(\zeta, \eta)| \leq r^k \rho^k h_k,$$

where h_k is the dimension of $\mathcal{H}_k(S)$. Now observe that under the above conditions

$$(5) \quad \lim_{k \rightarrow \infty} \sqrt[k]{\Delta_k^r(\omega)} = r^2 \text{ for } \Delta_k^r(\omega) = - \int_0^{r^2} t^k d\omega(t) \text{ and } \forall r \in (0, +\infty].$$

Indeed, it is obvious that $\Delta_k^r(\omega) \leq r^{2k}(1 - \omega(r^2))$ and hence

$$(6) \quad \limsup_{k \rightarrow \infty} \sqrt[k]{\Delta_k^r(\omega)} \leq r^2 .$$

On the other hand,

$$\Delta_k^r(\omega) \geq - \int_\delta^{r^2} t^k d\omega(t) \geq \delta^k (\omega(r^2) - \omega(\delta))$$

for any $\delta \in (0, r^2)$. Therefore

$$\liminf_{k \rightarrow \infty} \sqrt[k]{\Delta_k^r(\omega)} \geq \delta \lim_{k \rightarrow \infty} \sqrt[k]{(\omega(r^2) - \omega(\delta))} = \delta,$$

and hence by (6) the passage $\delta \rightarrow r^2$ gives (5). Further, note that $\Delta_k^r(\omega) \uparrow r$. Therefore by (5)

$$\liminf_{k \rightarrow \infty} \sqrt[k]{\Delta_k^\infty(\omega)} \geq r^2$$

for any $r > 0$, and consequently

$$(7) \quad \lim_{k \rightarrow \infty} \sqrt[k]{\Delta_k^\infty(\omega)} = +\infty.$$

The desired convergence follows from (4) and (7) in view of the estimate

$$(8) \quad h_k \leq Ck^{n-2}$$

(see, e.g. [7]). □

3.2. The following statement is the main theorem of this section.

Theorem 1. *Let $u \in b_\omega^2(\mathbb{R}^n)$, where $\omega \in \Omega^\infty$. Then*

$$(9) \quad u(x) = \int_{\mathbb{R}^n} u(y)R_\omega^\infty(x, y) d\mu_\omega(y), \quad x \in \mathbb{R}^n.$$

PROOF: The idea of the proof is the following. For any $r > 0$ we introduce a kernel

$$R_\omega^r(x, y) \stackrel{\text{def}}{=} \sum_{k=0}^\infty \frac{Z_k(x, y)}{\Delta_k^r(\omega)},$$

where $\Delta_k^r(\omega)$ is defined in (5). This kernel plays for a ball $|x| < r$ the same role, as $R_\omega(x, y)$ for a unit ball, after dilation we obtain the integral representation (10) from (2). And passing to limits as $r \rightarrow \infty$ we get (9) from (10), which is expected, because the coefficients $\Delta_k^r(\omega)$ of the expansion of the kernel $R_\omega^r(x, y)$ tend to the coefficients $\Delta_k^\infty(\omega)$ of $R_\omega^\infty(x, y)$.

Consider the function $\omega_r(t) = \omega(r^2t)$, $0 \leq t \leq 1$. Then obviously $\Delta_k(\omega_r) = r^{-2k} \Delta_k^r(\omega)$. Therefore by (5)

$$\lim_{k \rightarrow \infty} \sqrt[k]{\Delta_k(\omega_r)} = 1,$$

and hence $\omega_r \in \Omega$. On the other hand, $u(rx) \in b_{\omega_r}^2(\mathbf{B}^n)$. Thus, the representation (2) is valid for $u(rx)$. Now observe that for $|x| < r, |y| < r$

$$R_{\omega_r} \left(\frac{x}{r}, \frac{y}{r} \right) \equiv \sum_{k=0}^{\infty} \frac{Z_k(x, y)}{r^2 \Delta_k(\omega)} = \sum_{k=0}^{\infty} \frac{Z_k(x, y)}{\Delta_k^r(\omega)} \stackrel{\text{def}}{=} R_{\omega}^r(x, y),$$

and $d\mu_{\omega_r} \left(\frac{y}{r} \right) = d\mu_{\omega}(y)$. Consequently, (2) can be written in the form

$$(10) \quad u(x) = \int_{|y| < r} u(y) R_{\omega}^r(x, y) d\mu_{\omega}(y), \quad |x| < r,$$

and to prove (9) it suffices to show that for any fixed $x \in \mathbb{R}^n$

$$(11) \quad \lim_{r \rightarrow \infty} \int_{|y| < r} u(y) R_{\omega}^r(x, y) d\mu_{\omega}(y) = \int_{\mathbb{R}^n} u(y) R_{\omega}^{\infty}(x, y) d\mu_{\omega}(y).$$

To prove this relation, observe that by Lemma 1 the function $R_{\omega}^{\infty}(x, \cdot)$ is harmonic in \mathbb{R}^n . Therefore, by Hölder's inequality

$$\begin{aligned} \int_{\mathbb{R}^n} |u(y) R_{\omega}^{\infty}(x, y)| d\mu_{\omega}(y) &\leq \|u\|_{2, \omega} \left\{ \int_{\mathbb{R}^n} |R_{\omega}^{\infty}(x, y)|^2 d\mu_{\omega}(y) \right\}^{1/2} \\ &= \|u\|_{2, \omega} \left\{ \sum_{k=0}^{\infty} \frac{1}{(\Delta_k^{\infty})^2} \int_{\mathbb{R}^n} Z_k^2(x, \rho\zeta) d\mu_{\omega}(\rho\zeta) \right\}^{1/2} \\ &= \|u\|_{2, \omega} \left\{ \sum_{k=0}^{\infty} \frac{1}{(\Delta_k^{\infty})^2} \int_0^{\infty} \rho^{2k} |d\omega(\rho^2)| \int_S Z_k^2(x, \zeta) d\sigma(\zeta) \right\}^{1/2}. \end{aligned}$$

Further, it is evident that

$$(12) \quad \int_S Z_k^2(x, \zeta) d\sigma(\zeta) = |x|^{2k} \int_S Z_k^2 \left(\frac{x}{|x|}, \zeta \right) d\sigma(\zeta) = |x|^{2k} h_k.$$

Consequently,

$$(13) \quad \int_{\mathbb{R}^n} |u(y) R_{\omega}^{\infty}(x, y)| d\mu_{\omega}(y) \leq \|u\|_{2, \omega} \left\{ \sum_{k=0}^{\infty} \frac{|x|^{2k} h_k}{\Delta_k^{\infty}} \right\}^{1/2}.$$

According to (7) and (8), the right-hand side a series of this estimate is convergent. Hence we conclude that the relation (11) is equivalent to

$$\lim_{r \rightarrow \infty} \int_{|y| < r} u(y) [R_{\omega}^r(x, y) - R_{\omega}^{\infty}(x, y)] d\mu_{\omega}(y) = 0.$$

In order to prove the latter relation, assume that $|x| = r_0$, $r_0 + 1 < r_1 < r < +\infty$, and observe that

$$\begin{aligned}
 I(r) &\equiv \left| \int_{|y|<r} u(y)(R_\omega^r(x, y) - R_\omega^\infty(x, y)) d\mu_\omega(y) \right| \\
 &\leq \int_{|y|<r_1} |u(y)(R_\omega^r(x, y) - R_\omega^\infty(x, y))| d\mu_\omega(y) \\
 &\quad + \int_{r_1<|y|<r} |u(y)R_\omega^r(x, y)| d\mu_\omega(y) \\
 &\quad + \int_{r_1<|y|<r} |u(y)R_\omega^\infty(x, y)| d\mu_\omega(y) \equiv I_1(r) + I_2(r) + I_3(r).
 \end{aligned}$$

To estimate the summand $I_2(r)$ we once again apply Hölder’s inequality:

$$\begin{aligned}
 (14) \quad I_2(r) &\leq \left\{ \int_{r_1<|y|<r} |u(y)|^2 d\mu_\omega(y) \int_{r_1<|y|<r} |R_\omega^r(x, y)|^2 d\mu_\omega(y) \right\}^{1/2} \\
 &= \left\{ \int_{r_1<|y|<r} |u(y)|^2 d\mu_\omega(y) \sum_{k=0}^\infty \frac{1}{(\Delta_k^r)^2} \int_{r_1<|y|<r} Z_k^2(x, y) d\mu_\omega(y) \right\}^{1/2}.
 \end{aligned}$$

Further, according to (12)

$$\begin{aligned}
 (15) \quad \int_{r_1<|y|<r} Z_k^2(x, y) d\mu_\omega(y) &= \int_S Z_k^2(x, y) d\sigma(\zeta) \int_{r_1}^r \rho^{2k} |d\omega(\rho^2)| \\
 &= |x|^{2k} h_k \int_{r_1}^r \rho^{2k} |d\omega(\rho^2)|.
 \end{aligned}$$

Therefore

$$\begin{aligned}
 &\sum_{k=0}^\infty \frac{1}{(\Delta_k^r)^2} \int_{r_1<|y|<r} Z_k^2(x, \rho\eta) d\mu_\omega(\rho\eta) \\
 &\leq \sum_{k=0}^\infty \frac{|x|^{2k}}{(\Delta_k^r)^2} \int_{r_1}^r \rho^{2k} |d\omega(\rho^2)| \int_S Z_k^2\left(\frac{x}{|x|}, \eta\right) d\sigma(\eta) \leq \sum_{k=0}^\infty \frac{|x|^{2k} h_k}{\Delta_k^r} \leq \sum_{k=0}^\infty \frac{r_0^{2k} h_k}{\Delta_{r_0+1}^k}.
 \end{aligned}$$

The last series converges in virtue of (5) and (8), therefore $I_2(r) < \varepsilon/3$ for a given $\varepsilon > 0$ and r_1 large enough. On the other hand, from (13) it follows that $I_3(r) < \varepsilon/3$ for r_1 large enough. Besides, for any fixed r_1

$$\begin{aligned}
 (16) \quad I_1(r) &\leq \|u\|_{2,\omega} \left\{ \int_{|y|<r_1} |R_\omega^r(x, y) - R_\omega^\infty(x, y)|^2 d\mu_\omega(y) \right\}^{1/2} \\
 &= \|u\|_{2,\omega} \left\{ \sum_{k=0}^\infty \left(\frac{1}{\Delta_k^r} - \frac{1}{\Delta_k^\infty} \right)^2 \int_{|y|<r_1} Z_k^2(x, y) d\mu_\omega(y) \right\}^{1/2}.
 \end{aligned}$$

Therefore, by (15)

$$I_1(r) \leq \|u\|_{2,\omega} \left\{ \sum_{k=0}^{\infty} \left(\frac{1}{\Delta_k^r} - \frac{1}{\Delta_k^\infty} \right)^2 |x|^{2k} h_k \int_0^{r_1} \rho^{2k} |d\omega(\rho^2)| \right\}^{1/2},$$

and the latter series has a convergent majorant independent of r . Indeed,

$$\begin{aligned} & \sum_{k=0}^{\infty} \left(\frac{1}{\Delta_k^r} - \frac{1}{\Delta_k^\infty} \right)^2 |x|^{2k} h_k \int_0^{r_1} \rho^{2k} |d\omega(\rho^2)| \\ & \leq \sum_{k=0}^{\infty} |x|^{2k} h_k \left(\frac{2}{\Delta_k^r} \right)^2 \Delta_k^{r_1} \leq 4 \sum_{k=0}^{\infty} \frac{r_0^{2k} h_k}{\Delta_k^{r_1}} < +\infty, \end{aligned}$$

where the right-hand side series converges due to (5) and (8), as $r_0 < r_1$. Therefore, the right-hand side of (16) vanishes as $r \rightarrow +\infty$ and hence $I_1(r) < \varepsilon/3$ for r large enough. Thus, we conclude that $I(r) \rightarrow 0$ as $r \rightarrow +\infty$, which implies the desired representation (9). \square

3.3. As $b_\omega^2(\mathbb{R}^n)$ is a closed subspace of the Hilbert space $L_\omega^2(\mathbb{R}^n)$, there is a unique orthogonal projection Q of $L_\omega^2(\mathbb{R}^n)$ onto $b_\omega^2(\mathbb{R}^n)$, which is described by

Theorem 2. *The operator*

$$Q_\omega[u](x) = \int_{\mathbb{R}^n} u(y) R_\omega^\infty(x, y) d\mu_\omega(y), \quad x \in \mathbb{R}^n, \quad u \in L_\omega^2(\mathbb{R}^n),$$

is the orthogonal projection of $L_\omega^2(\mathbb{R}^n)$ onto $b_\omega^2(\mathbb{R}^n)$.

The proof of this theorem as well as of the statements below follows the same lines as the corresponding one in [1] and is thus omitted.

Proposition 1. *Let a function $\tilde{\omega} \in \Omega^\infty$ be continuously differentiable in $[0, +\infty)$ and such that $\tilde{\omega}(+\infty) = 0$, $\tilde{\omega}'(t) < 0$ and is bounded on $[0, +\infty)$ and $\int_0^{+\infty} t^{-1} d\tilde{\omega}(t) > -\infty$. Further, let ω be the Volterra square of $\tilde{\omega}$, i.e.*

$$(17) \quad \omega(x) = - \int_0^\infty \tilde{\omega}\left(\frac{x}{t}\right) d\tilde{\omega}(t), \quad 0 < x < 1.$$

Then $\omega \in \Omega^\infty$ and

$$(18) \quad \Delta_m^\infty(\omega) = [\Delta_m^\infty(\tilde{\omega})]^2, \quad m \geq 0.$$

By $h^p(B)$ we denote the ordinary harmonic Hardy space in B . Besides, we consider the operator

$$L_{\tilde{\omega}}[u](x) = - \int_0^\infty u(tx) d\tilde{\omega}(t).$$

The following two theorems establish an isometry along with its inversion between $b_\omega^2(\mathbb{R}^n)$ and $L^2(S)$.

Theorem 3. *The mapping $f \mapsto R_{\tilde{\omega}}[f]$, where*

$$R_{\tilde{\omega}}[f](x) = \int_S f(\zeta) R_{\tilde{\omega}}^{\infty}(x, \zeta) d\sigma(\zeta)$$

is a linear isometry from $L^2(S)$ to $b_{\tilde{\omega}}^2(\mathbb{R}^n)$.

Theorem 4. *Let $f \in L^2(S)$ and $u = R_{\tilde{\omega}}[f]$. Then*

- (a) $L_{\tilde{\omega}}[u] = P[f]$, where $P[f]$ is the Poisson integral (1);
- (b) the mapping $u \mapsto L_{\tilde{\omega}}[u]$ is a linear isometry of $b_{\tilde{\omega}}^2(\mathbb{R}^n)$ onto $h^2(B)$.

Remark 1. It is well known that for $f \in L^2(S)$ the function $P[f]$ has a nontangential limit $f(\zeta)$ at almost every point $\zeta \in S$. Thus, it is natural to identify f and $P[f]$ and to say that the operators L_{ω} and R_{ω} are mutually inverse.

REFERENCES

- [1] Petrosyan A.I., *On weighted classes of harmonic functions in the unit ball of \mathbf{R}^n* , Complex Var. Theory Appl. **50** (2005), no. 12, 953–966.
- [2] Petrosyan A.I., *On A_{ω}^p spaces in the unit ball of \mathbf{C}^n* , J. Anal. Appl. **3** (2005), no. 1, 47–53.
- [3] Djrbashian M.M., *On canonical representation of functions meromorphic in the unit disc* (in Russian), Dokl. Akad. Nauk of Armenia **3** (1945), no. 1, 3–9.
- [4] Djrbashian M.M., *On the representability problem of analytic functions* (in Russian), Soobsch. Inst. Math. and Mech. AN Armenii **2** (1948), 3–40.
- [5] Jerbashian A.M., *On the theory of weighted classes of area integrable regular functions*, Complex Var. Theory Appl. **50** (2005), no. 3, 155–183.
- [6] Rudin W., *Real and Complex Analysis*, McGraw-Hill, New York, 1987.
- [7] Axler S., Bourdon P., Ramsey W., *Harmonic Function Theory*, Springer, New York, 2001.

YEREVAN STATE UNIVERSITY, ALECK MANOOGIAN STREET 1, YEREVAN 375049, ARMENIA

E-mail: albpet@xter.net

(Received September 21, 2005, revised November 23, 2005)