

Martin boundary associated with a system of PDE

ALLAMI BENYAICHE, SALMA GHIATE

Abstract. In this paper, we study the Martin boundary associated with a harmonic structure given by a coupled partial differential equations system. We give an integral representation for non negative harmonic functions of this structure. In particular, we obtain such results for biharmonic functions (i.e. $\Delta^2\varphi = 0$) and for non negative solutions of the equation $\Delta^2\varphi = \varphi$.

Keywords: Martin boundary, biharmonic functions, coupled partial differential equations

Classification: Primary 31C35; Secondary 31B30, 31B10, 60J50

1. Introduction

Let D be a domain in \mathbb{R}^d , $d \geq 1$, and let L_i , $i = 1, 2$, be two second order elliptic differential operators on D leading to harmonic spaces (D, H_{L_i}) with Green functions G_i (see [18]). Moreover, we assume that every ball $B \subset \bar{B} \subset D$ is an L_i -regular set. Throughout this paper we consider two positive Radon measures μ_1 and μ_2 such that $K_D^{\mu_i} = \int_D G_i(\cdot, y)\mu_i(dy)$ is a bounded continuous real function on D , $i = 1, 2$, and

$$\|K_D^{\mu_1}\|_\infty \|K_D^{\mu_2}\|_\infty < 1.$$

We consider the system:

$$(S) \begin{cases} L_1u = -v\mu_1, \\ L_2v = -u\mu_2. \end{cases}$$

Note that if U is a relatively compact open subset of D , $\mu_1 = \lambda^d$, where λ^d is the Lebesgue measure, $\mu_2 = 0$ and $L_1 = L_2 = \Delta$, then we obtain the classical biharmonic case on U . In the case when $\mu_1 = \mu_2 = \lambda^d$ and $\lambda^d(D) < \infty$, we obtain equations of type $\Delta^2\varphi = \varphi$. In this work, we shall study the Martin boundary associated with the balayage space given by the system (S) (see [7], [14] and [19]), and we shall characterize minimal points of this boundary in order to give an integral representation for non negative solutions of the system (S) .

Let us note that the notion of a balayage space defined by J. Bliedtner and W. Hansen in [7] is more general than that of a P-harmonic space. It covers harmonic structures given by elliptic or parabolic partial differential equations, Riesz potentials, and biharmonic equations (which are a particular case of this

work). In the biharmonic case, a similar study can be done using couples of functions as presented in [3], [5], [8], [9], [21] and [22].

We are also grateful to the referee for his remarks and comments.

2. Notations and preliminaries

For $j = 1, 2$, let $X_j = D \times \{j\}$, and let $X = X_1 \cup X_2$. Moreover, let i_j and π_j be the mappings defined by

$$i_j : \begin{cases} D \longrightarrow X_j \\ x \longmapsto (x, j) \end{cases} \quad \text{and} \quad \pi_j : \begin{cases} X_j \longrightarrow D \\ (x, j) \longmapsto x. \end{cases}$$

Let \mathcal{U}_0 be the set of all balls B such that $B \subset \bar{B} \subset D$, \mathcal{U}_j be the image of \mathcal{U}_0 by i_j , $j = 1, 2$, and $\mathcal{U} = \mathcal{U}_1 \cup \mathcal{U}_2$.

Definition 2.1. Let v be a measurable function on X . For $U \in \mathcal{U}_1$, we define the kernel S_U by

$$S_U v = (H_{\pi_1(U)}^1(v \circ i_1)) \circ \pi_1 + (K_{\pi_1(U)}^{\mu_1}(v \circ i_2)) \circ \pi_1.$$

For $U \in \mathcal{U}_2$, we define the kernel S_U by

$$S_U v = (H_{\pi_2(U)}^2(v \circ i_2)) \circ \pi_2 + (K_{\pi_2(U)}^{\mu_2}(v \circ i_1)) \circ \pi_2,$$

where $H_{\pi_j(U)}^j$, $j = 1, 2$, denote the harmonic kernels associated with (D, H_{L_j}) and

$$K_{\pi_i(U)}^{\mu_i}(w) = \int G_i^{\pi_i(U)}(\cdot, y) w(y) \mu_i(dy) \quad i = 1, 2,$$

where w is a measurable function on D and $G_i^{\pi_i(U)}$ is the Green function associated with the operator L_i on $\pi_i(U)$. Let G_j , $j = 1, 2$, be the Green kernel associated with L_j on D . The family of kernels $(S_U)_{U \in \mathcal{U}}$ yields a balayage space on X as defined in [7] and [14].

Let ${}^*\mathcal{H}(X)$ denote the set of all hyperharmonic functions on X , i.e.

$${}^*\mathcal{H}(X) := \{v \in \mathcal{B}(X) : v \text{ is l.s.c. and } S_U v \leq v \quad \forall U \in \mathcal{U}\},$$

where $\mathcal{B}(X)$ denotes the set of all Borel functions on X . Let $\mathcal{S}(X)$ be the set of all superharmonic functions on X , i.e.

$$\mathcal{S}(X) := \{v \in {}^*\mathcal{H}(X) : (S_U v)|_{U \in C(U)} \quad \forall U \in \mathcal{U}\},$$

and let $\mathcal{H}(X)$ be the set of all harmonic functions on X :

$$\mathcal{H}(X) := \{h \in \mathcal{S}(X) : S_U h = h \quad \forall U \in \mathcal{U}\}.$$

Denoting $\mathcal{W} := {}^*\mathcal{H}^+(X)$, the space (X, \mathcal{W}) is a balayage space (see [7] and [14]).

For every positive numerical function φ on X and for every $U \in \mathcal{U}$, the reduct R_φ^U is defined by

$$R_\varphi^U := \inf\{v \in {}^*\mathcal{H}(X) : v \geq \varphi \text{ on } U\}.$$

Let \widehat{R}_φ^U be the lower semi-continuous regularization of R_φ^U , i.e.

$$\widehat{R}_\varphi^U(x) := \liminf_{y \rightarrow x} R_\varphi^U(y), \quad x \in X.$$

Theorem 2.1. *Let s be a function on X such that*

$$K_D^{\mu_j}(s \circ i_k) < \infty, \quad j \neq k, \quad j, k = 1, 2.$$

The following statements are equivalent.

1. s is a superharmonic function on X .
2. $s_j := s \circ i_j - K_D^{\mu_j}(s \circ i_k)$, $j \neq k$, $j, k \in \{1, 2\}$, are L_j -superharmonic on D .

PROOF: Let s be a superharmonic function on X and let $U \in \mathcal{U}_0$. We have

$$i_1(U) \in \mathcal{U}_1 \quad \text{and} \quad \pi_1(i_1(U)) = U.$$

Since $S_{i_1(U)}s \leq s$, we have

$$H_U^1(s \circ i_1) + K_U^{\mu_1}(s \circ i_2) \leq s \circ i_1.$$

Knowing that

$$K_U^{\mu_1}(s \circ i_2) = K_D^{\mu_1}(s \circ i_2) - H_U^1(K_D^{\mu_1}(s \circ i_2)),$$

we obtain

$$H_U^1(s \circ i_1) + K_D^{\mu_1}(s \circ i_2) - H_U^1(K_D^{\mu_1}(s \circ i_2)) \leq s \circ i_1.$$

Therefore

$$H_U^1(s \circ i_1 - K_D^{\mu_1}(s \circ i_2)) \leq s \circ i_1 - K_D^{\mu_1}(s \circ i_2).$$

So, $s_1 := s \circ i_1 - K_D^{\mu_1}(s \circ i_2)$ is an L_1 -superharmonic function on D . Similarly, we prove that $s_2 := s \circ i_2 - K_D^{\mu_2}(s \circ i_1)$ is L_2 -superharmonic on D . Conversely, we assume that s_i , $i = 1, 2$, are L_i -superharmonic functions. Let $U \in \mathcal{U}_j$, $j = 1, 2$ and $k \neq j$. Since s_j is an L_j -superharmonic function,

$$H_{\pi_j(U)}^j s_j \leq s_j.$$

Hence

$$H_{\pi_j(U)}^j(s \circ i_j - K_D^{\mu_j}(s \circ i_k)) \leq s \circ i_j - K_D^{\mu_j}(s \circ i_k).$$

Therefore

$$H_{\pi_j(U)}^j(s \circ i_j) + K_{\pi_j(U)}^{\mu_j}(s \circ i_k) \leq s \circ i_j.$$

So,

$$S_U s \leq s, \quad \forall U \in \mathcal{U}.$$

Thus s is superharmonic on X . □

Corollary 2.1. *Let v be a function on X such that $K_D^{\mu_j}(v \circ i_k)$, $j \neq k$, $j, k \in \{1, 2\}$, is a finite function. Then the following properties are equivalent.*

1. v is harmonic on X .
2. $v \circ i_1 - K_D^{\mu_1}(v \circ i_2)$ and $v \circ i_2 - K_D^{\mu_2}(v \circ i_1)$ are L_1 -harmonic and L_2 -harmonic function on D , respectively.

Remarks 2.1. (1) Note that if v is a positive harmonic function on X , then $K_D^{\mu_j}(v \circ i_k)$, $j \neq k$, $j, k \in \{1, 2\}$, is a finite function.

(2) If $v \in \mathcal{H}(X)$, then the couple $(v \circ i_1, v \circ i_2)$ is a solution of (S).

Corollary 2.2. *Let v be a positive function defined on X . Then the following properties are equivalent.*

1. v is hyperharmonic on X .
2. The function

$$v_j := \begin{cases} v \circ i_j - K_D^{\mu_j}(v \circ i_k) & \text{if } K_D^{\mu_j}(v \circ i_k) < \infty, \\ +\infty & \text{otherwise} \end{cases}$$

is a positive L_j -hyperharmonic function on D , $j \neq k$, $j, k \in \{1, 2\}$.

If we identify a function s on X with the couple $(s \circ i_1, s \circ i_2)$ defined on D , then we get the following N. Bouleau’s decomposition [9]:

Theorem 2.2. *Any superharmonic function s on X can be written as $s = t + Vs$, where*

$$V = \begin{pmatrix} 0 & K_D^{\mu_1} \\ K_D^{\mu_2} & 0 \end{pmatrix}$$

and t is a function on X defined by

$$t := \begin{cases} s_1 \circ \pi_1 & \text{on } X_1, \\ s_2 \circ \pi_2 & \text{on } X_2, \end{cases}$$

where $s_j := s \circ i_j - K_D^{\mu_j}(s \circ i_k)$, $j \neq k$, $j, k \in \{1, 2\}$.

PROOF: It follows from Theorem 2.1 that s_j , $j = 1, 2$, is L_j -superharmonic on D . Then, if we identify the function s with the couple $(s \circ i_1, s \circ i_2)$ defined on D and the function t with the couple $(t \circ i_1, t \circ i_2) = (s_1, s_2)$ defined on D , we have

$$\begin{pmatrix} 0 & K_D^{\mu_1} \\ K_D^{\mu_2} & 0 \end{pmatrix} \begin{pmatrix} s \circ i_1 \\ s \circ i_2 \end{pmatrix} + \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} = \begin{pmatrix} s \circ i_1 \\ s \circ i_2 \end{pmatrix}.$$

□

Remark 2.1. In the classical biharmonic case, we obtain the N. Bouleau’s decomposition [9]. Indeed, if we identify a function s on X with the couple $(s \circ i_1, s \circ i_2)$ on D , then

$$s \circ i_1 = s_1 + K_D^{\mu_1}(s \circ i_2),$$

with s_1 L_1 -superharmonic on D and the N. Bouleau’s kernel V is given by $V = K_D^{\mu_1}$.

3. Martin boundary associated with (S)

Let us fix $x_0 \in D$ and set for all $x, y \in D$

$$g^1(x, y) := \begin{cases} \frac{G_1(x, y)}{G_1(x_0, y)} & \text{if } x \neq x_0 \text{ or } y \neq x_0, \\ 1 & \text{if } x = y = x_0, \end{cases}$$

and

$$g^2(x, y) := \begin{cases} \frac{G_2(x, y)}{G_2(x_0, y)} & \text{if } x \neq x_0 \text{ or } y \neq x_0, \\ 1 & \text{if } x = y = x_0. \end{cases}$$

Let $\mathcal{A}_1 = \{g^1(x, \cdot), x \in D\}$, $\mathcal{A}_2 = \{g^2(x, \cdot), x \in D\}$ and $\mathcal{A} = \mathcal{A}_1 \cup \mathcal{A}_2$.

As in [10] and [12], we consider the Martin compactification \widehat{D} of D associated with \mathcal{A} . The boundary $\Delta = \widehat{D} \setminus D$ of D is called the Martin boundary of D associated with the system (S).

The function $g^k(x, \cdot)$, $k = 1, 2$, $x \in D$ can be extended, on \widehat{D} , to a continuous function denoted $g^k(x, \cdot)$, $k = 1, 2$, $x \in D$ as well.

In the following, we denote $Q := \sum_{n=0}^{+\infty} (K_D^{\mu_1} K_D^{\mu_2})^n$ (resp. $T := \sum_{n=0}^{+\infty} (K_D^{\mu_2} K_D^{\mu_1})^n$) which coincides with $(I - K_D^{\mu_1} K_D^{\mu_2})^{-1}$ (resp. $(I - K_D^{\mu_2} K_D^{\mu_1})^{-1}$) on $\mathcal{B}_b(D)$, where $(I - K_D^{\mu_1} K_D^{\mu_2})^{-1}$ (resp. $(I - K_D^{\mu_2} K_D^{\mu_1})^{-1}$) is the inverse of the operator $(I - K_D^{\mu_1} K_D^{\mu_2})$ (resp. $(I - K_D^{\mu_2} K_D^{\mu_1})$) on $\mathcal{B}_b(D)$, and $\mathcal{B}_b(D)$ denotes the set of all bounded Borel measurable functions on D . We recall the following equalities

$$\begin{aligned} (K_D^{\mu_1} K_D^{\mu_2})Q &= Q(K_D^{\mu_1} K_D^{\mu_2}), \\ (K_D^{\mu_1} K_D^{\mu_2})Q + I &= Q. \end{aligned}$$

Similarly we have

$$\begin{aligned} (K_D^{\mu_2} K_D^{\mu_1})T &= T(K_D^{\mu_2} K_D^{\mu_1}), \\ (K_D^{\mu_2} K_D^{\mu_1})T + I &= T, \\ K_D^{\mu_2} Q &= T K_D^{\mu_2} \end{aligned}$$

and

$$K_D^{\mu_1} T = Q K_D^{\mu_1}.$$

Remark 3.1. Note that if φ is a finite positive Borel measurable function on D such that $K_D^{\mu_1} K_D^{\mu_2} \varphi$ is bounded, then $Q\varphi < +\infty$.

Theorem 3.1. *Let $t_i, i = 1, 2$, be two L_i -harmonic functions on D such that $K_D^{\mu_j} t_k$ is finite and $K_D^{\mu_k} K_D^{\mu_j} t_k$ is bounded, $j \neq k, j, k \in \{1, 2\}$, on D . Then the functions v and w defined on X by*

$$v := \begin{cases} (Qt_1) \circ \pi_1 & \text{on } X_1, \\ (K_D^{\mu_2} Qt_1) \circ \pi_2 & \text{on } X_2, \end{cases}$$

and

$$w := \begin{cases} (QK_D^{\mu_1} t_2) \circ \pi_1 & \text{on } X_1, \\ (Tt_2) \circ \pi_2 & \text{on } X_2 \end{cases}$$

are harmonic on X .

Remark 3.2. In the biharmonic case, if we assume that $K_D^{\lambda^d} t_2 < \infty$, then $(t_1, 0)$ and $(K_D^{\lambda^d} t_2, t_2)$ are biharmonic.

PROOF: Let us prove first that v and w are finite.

(i) We have

$$(Qt_1) \circ \pi_1 = (QK_D^{\mu_1} K_D^{\mu_2} t_1) \circ \pi_1 + t_1 \circ \pi_1.$$

Since $K_D^{\mu_1} K_D^{\mu_2} t_1$ is bounded and t_1 is finite,

$$(Qt_1) \circ \pi_1 < \infty.$$

(ii) We have also

$$(K_D^{\mu_2} Qt_1) \circ \pi_2 = (TK_D^{\mu_2} t_1) \circ \pi_2,$$

hence

$$(K_D^{\mu_2} Qt_1) \circ \pi_2 = (TK_D^{\mu_2} K_D^{\mu_1} K_D^{\mu_2} t_1) \circ \pi_2 + (K_D^{\mu_2} t_1) \circ \pi_2.$$

Since $K_D^{\mu_1} K_D^{\mu_2} t_1$ is bounded and $K_D^{\mu_2} t_1$ is finite,

$$(K_D^{\mu_2} Qt_1) \circ \pi_2 < \infty.$$

(iii) We have

$$(QK_D^{\mu_1} t_2) \circ \pi_1 = (QK_D^{\mu_1} K_D^{\mu_2} K_D^{\mu_1} t_2) \circ \pi_1 + (K_D^{\mu_1} t_2) \circ \pi_1.$$

Knowing that $K_D^{\mu_2} K_D^{\mu_1} t_2$ is bounded and $K_D^{\mu_1} t_2$ is finite, we have

$$(QK_D^{\mu_1} t_2) \circ \pi_1 < \infty.$$

(iv) We have

$$(Tt_2) \circ \pi_2 = (TK_D^{\mu_2} K_D^{\mu_1} t_2) \circ \pi_2 + t_2 \circ \pi_2.$$

Since $K_D^{\mu_2} K_D^{\mu_1} t_2$ is bounded and t_2 is finite,

$$(Tt_2) \circ \pi_2 < \infty.$$

Let us show now that v and w are harmonic. From Corollary 2.1, it suffices to show that $v \circ i_j - K_D^{\mu_j}(v \circ i_k)$ and $w \circ i_j - K_D^{\mu_j}(w \circ i_k)$, $j \neq k$, $j, k \in \{1, 2\}$, are L_j -harmonic functions on D .

(v) On the one hand,

$$v \circ i_1 - K_D^{\mu_1}(v \circ i_2) = Qt_1 - (K_D^{\mu_1} K_D^{\mu_2})Qt_1.$$

As

$$Qt_1 = (K_D^{\mu_1} K_D^{\mu_2})Qt_1 + t_1,$$

we get

$$v \circ i_1 - K_D^{\mu_1}(v \circ i_2) = t_1.$$

Since t_1 is an L_1 -harmonic function on D , $v \circ i_1 - K_D^{\mu_1}(v \circ i_2)$ is L_1 -harmonic on D .

On the other hand,

$$v \circ i_2 - K_D^{\mu_2}(v \circ i_1) = K_D^{\mu_2}Qt_1 - K_D^{\mu_2}Qt_1 = 0,$$

i.e. $v \circ i_2 - K_D^{\mu_2}(v \circ i_1)$ is L_2 -harmonic on D . Then we conclude that v is harmonic on X .

(vi) Since

$$(*) \quad T = K_D^{\mu_2}QK_D^{\mu_1} + I,$$

we have

$$w \circ i_1 - K_D^{\mu_1}(w \circ i_2) = (QK_D^{\mu_1} - K_D^{\mu_1}K_D^{\mu_2}QK_D^{\mu_1} - K_D^{\mu_1})t_2.$$

As

$$Q = (K_D^{\mu_1}K_D^{\mu_2})Q + I,$$

we obtain

$$w \circ i_1 - K_D^{\mu_1}(w \circ i_2) = 0.$$

Using (*), we have

$$w \circ i_2 - K_D^{\mu_2}(w \circ i_1) = (K_D^{\mu_2}QK_D^{\mu_1} + I - K_D^{\mu_2}QK_D^{\mu_1})t_2 = t_2.$$

Then $w \circ i_j - K_D^{\mu_j}(w \circ i_k)$ is L_j -harmonic on D and therefore, w is a harmonic function on X . □

Corollary 3.1. *Let $t_i, i = 1, 2$, be two positive L_i -hyperharmonic functions on D . Then the functions v and w defined on D by*

$$v := \begin{cases} (Qt_1) \circ \pi_1 & \text{on } X_1, \\ (K_D^{\mu_2}Qt_1) \circ \pi_2 & \text{on } X_2, \end{cases}$$

and

$$w := \begin{cases} (QK_D^{\mu_1}t_2) \circ \pi_1 & \text{on } X_1, \\ (Tt_2) \circ \pi_2 & \text{on } X_2 \end{cases}$$

are hyperharmonic on X .

Theorem 3.2. *Let ν_1 and ν_2 be two positive Radon measures on Δ such that*

$$\int_{\Delta} K_D^{\mu_j}g^k(\cdot, y) d\nu_k(y) < \infty$$

and

$$\int_{\Delta} K_D^{\mu_j}K_D^{\mu_k}g^j(\cdot, y) d\nu_j(y)$$

is bounded on $D, j \neq k, j, k \in \{1, 2\}$. Then the function v defined on X_1 by

$$v := \int_{\Delta} (Qg^1(\cdot, y)) \circ \pi_1 d\nu_1(y) + \int_{\Delta} (QK_D^{\mu_1}g^2(\cdot, y)) \circ \pi_1 d\nu_2(y)$$

and on X_2 by

$$v := \int_{\Delta} (K_D^{\mu_2}Qg^1(\cdot, y)) \circ \pi_2 d\nu_1(y) + \int_{\Delta} (Tg^2(\cdot, y)) \circ \pi_2 d\nu_2(y)$$

is harmonic on X .

PROOF: It suffices to replace the functions t_j from Theorem 3.1 with the L_j -harmonic functions $\int_{\Delta} g^j(\cdot, y) d\nu_j(y)$. □

Corollary 3.2. *Let ν_1 and ν_2 be two positive Radon measures on Δ such that $\int_{\Delta} K_D^{\mu_1}g^2(\cdot, y) d\nu_2(y) < \infty$. Then*

$$(v, w) = \left(\int_{\Delta} g^1(\cdot, y) d\nu_1(y) + \int_{\Delta} K_D^{\mu_1}g^2(\cdot, y) d\nu_2(y), \int_{\Delta} g^2(\cdot, y) d\nu_2(y) \right)$$

is a biharmonic couple in the classical sense.

Theorem 3.3. *Let v be a positive harmonic function on X such that $K_D^{\mu_j} K_D^{\mu_k} (v \circ i_j)$ is bounded on D , $j, k \in \{1, 2\}$, $j \neq k$. Then there exist two positive Radon measures ν_1 and ν_2 supported by Δ such that v can be represented on X_1 by*

$$v = \int_{\Delta} (Qg^1(\cdot, y)) \circ \pi_1 d\nu_1(y) + \int_{\Delta} (QK_D^{\mu_1} g^2(\cdot, y)) \circ \pi_1 d\nu_2(y)$$

and on X_2 by

$$v = \int_{\Delta} (K_D^{\mu_2} Qg^1(\cdot, y)) \circ \pi_2 d\nu_1(y) + \int_{\Delta} (Tg^2(\cdot, y)) \circ \pi_2 d\nu_2(y).$$

PROOF: Let $(D_n)_n$ be an increasing sequence of relatively compact open subsets of D such that $D = \bigcup D_n$, and let v be a positive harmonic function on X . From Corollary 2.1, the positive functions $v \circ i_1 - K_D^{\mu_1}(v \circ i_2)$ and $v \circ i_2 - K_D^{\mu_2}(v \circ i_1)$ are L_1 -harmonic and L_2 -harmonic on D , respectively. Then for all $n \in \mathbb{N}$, both $\widehat{R}_{v \circ i_1 - K_D^{\mu_1}(v \circ i_2)}^{D_n}$ and $\widehat{R}_{v \circ i_2 - K_D^{\mu_2}(v \circ i_1)}^{D_n}$ are L_1 -potential and L_2 -potential on D , respectively. Therefore, there exist two positive Radon measures μ_n^1 and μ_n^2 on D such that

$$\widehat{R}_{v \circ i_1 - K_D^{\mu_1}(v \circ i_2)}^{D_n} = \int_D G_1(\cdot, y) d\mu_n^1(y)$$

and

$$\widehat{R}_{v \circ i_2 - K_D^{\mu_2}(v \circ i_1)}^{D_n} = \int_D G_2(\cdot, y) d\mu_n^2(y).$$

Then we have

$$\widehat{R}_{v \circ i_1 - K_D^{\mu_1}(v \circ i_2)}^{D_n} = \int_D g^1(\cdot, y) d\nu_n^1(y)$$

and

$$\widehat{R}_{v \circ i_2 - K_D^{\mu_2}(v \circ i_1)}^{D_n} = \int_D g^2(\cdot, y) d\nu_n^2(y)$$

with

$$d\nu_1(y) = G_1(x_0, \cdot) d\mu_n^1(y)$$

and

$$d\nu_2(y) = G_2(x_0, \cdot) d\mu_n^2(y).$$

Since $\widehat{R}_{v \circ i_j - K_D^{\mu_j}(v \circ i_k)}^{D_n}$ is L_j -harmonic on $D \setminus D_n$, $j \neq k$, $j, k \in \{1, 2\}$, ν_n^1 and ν_n^2 are necessarily supported by $D \setminus D_n$.

Because of $\|\nu_n^j\| \leq (v \circ i_j)(x_0) - K_D^{\mu_j}(v \circ i_k)(x_0)$, $j = 1, 2$, we may extract two subsequences $(\nu_{p(n)}^1)$ and $(\nu_{p(n)}^2)$ converging vaguely to two positive Radon measures ν^1 and ν^2 on $\bar{D} = \widehat{D}$. So, ν^1 and ν^2 are supported by Δ . Therefore

$$\begin{cases} v \circ i_1 - K_D^{\mu_1}(v \circ i_2) = \int_{\Delta} g^1(\cdot, y) d\nu^1(y), \\ v \circ i_2 - K_D^{\mu_2}(v \circ i_1) = \int_{\Delta} g^2(\cdot, y) d\nu^2(y). \end{cases}$$

Hence

$$\begin{cases} v \circ i_1 = \int_{\Delta} g^1(\cdot, y) d\nu^1(y) + K_D^{\mu_1} \left(\int_{\Delta} g^2(\cdot, y) d\nu^2(y) + K_D^{\mu_2} (v \circ i_1) \right), \\ v \circ i_2 = \int_{\Delta} g^2(\cdot, y) d\nu^2(y) + K_D^{\mu_2} (v \circ i_1), \end{cases}$$

and

$$\begin{cases} v \circ i_1 = \int_{\Delta} g^1(\cdot, y) d\nu^1(y) + \int_{\Delta} K_D^{\mu_1} g^2(\cdot, y) d\nu^2(y) + K_D^{\mu_1} K_D^{\mu_2} (v \circ i_1), \\ v \circ i_2 = \int_{\Delta} g^2(\cdot, y) d\nu^2(y) + K_D^{\mu_2} (v \circ i_1). \end{cases}$$

Thus,

$$\begin{cases} Q(v \circ i_1) = \int_{\Delta} Qg^1(\cdot, y) d\nu^1(y) + \int_{\Delta} QK_D^{\mu_1} g^2(\cdot, y) d\nu^2(y) \\ \quad + QK_D^{\mu_1} K_D^{\mu_2} (v \circ i_1), \\ v \circ i_2 = \int_{\Delta} g^2(\cdot, y) d\nu^2(y) + K_D^{\mu_2} (v \circ i_1). \end{cases}$$

Since

$$QK_D^{\mu_1} K_D^{\mu_2} + I = Q,$$

we obtain

$$\begin{cases} K_D^{\mu_1} K_D^{\mu_2} Q(v \circ i_1) + v \circ i_1 = \int_{\Delta} Qg^1(\cdot, y) d\nu^1(y) + \int_{\Delta} QK_D^{\mu_1} g^2(\cdot, y) d\nu^2(y) \\ \quad + QK_D^{\mu_1} K_D^{\mu_2} (v \circ i_1), \\ v \circ i_2 = \int_{\Delta} g^2(\cdot, y) d\nu^2(y) + K_D^{\mu_2} (v \circ i_1). \end{cases}$$

Since $K_D^{\mu_1} K_D^{\mu_2} (v \circ i_1)$ is bounded,

$$\begin{cases} v \circ i_1 = \int_{\Delta} Qg^1(\cdot, y) d\nu_1(y) + \int_{\Delta} QK_D^{\mu_1} g^2(\cdot, y) d\nu_2(y), \\ v \circ i_2 = \int_{\Delta} K_D^{\mu_2} Qg^1(\cdot, y) d\nu_1(y) + \int_{\Delta} Tg^2(\cdot, y) d\nu_2(y). \end{cases}$$

So the function v can be written on X_1 as

$$v = \int_{\Delta} (Qg^1(\cdot, y)) \circ \pi_1 d\nu_1(y) + \int_{\Delta} (QK_D^{\mu_1} g^2(\cdot, y)) \circ \pi_1 d\nu_2(y)$$

and on X_2 as

$$v = \int_{\Delta} (K_D^{\mu_2} Qg^1(\cdot, y)) \circ \pi_2 d\nu_1(y) + \int_{\Delta} (Tg^2(\cdot, y)) \circ \pi_2 d\nu_2(y).$$

□

Corollary 3.3 ([5]). *Let (v, w) be a positive biharmonic couple in the classical sense. Then there exist two positive Radon measures μ and ν supported by Δ such that*

$$\begin{cases} v = \int_{\Delta} g^1(\cdot, y) d\mu(y) + \int_{\Delta} K_D^{\mu_1} g^2(\cdot, y) d\nu(y), \\ w = \int_{\Delta} g^2(\cdot, y) d\nu(y). \end{cases}$$

4. Minimal points and uniqueness of the integral representation

Definition 4.1. (1) A positive L_1 -harmonic (resp. L_2 -harmonic) function h on D is called L_1 -minimal (resp. L_2 -minimal) if for any positive L_1 -harmonic (resp. L_2 -harmonic) function u on D , $u \leq h$ implies $u = \alpha h$ with a factor $\alpha > 0$.

(2) A positive harmonic function h on X is called *minimal* if for any positive harmonic function u on X , $u \leq h$ implies $u = \alpha h$ with a factor $\alpha > 0$.

Denote

$$\begin{aligned} \Delta_1 &= \{y \in \Delta : g^1(\cdot, y) \text{ is } L_1\text{-minimal}\}, \\ \Delta_2 &= \{y \in \Delta : g^2(\cdot, y) \text{ is } L_2\text{-minimal}\}. \end{aligned}$$

Note that for all $y \in \Delta$, the function $g^1(\cdot, y)$ (resp. $g^2(\cdot, y)$) is L_1 -harmonic (resp. L_2 -harmonic) on D .

Proposition 4.1. Any positive harmonic function v on X such that $K_D^{\mu_k} K_D^{\mu_j} (v \circ i_k)$ is bounded for all $j \neq k$, $j, k \in \{1, 2\}$, can be written as $v = w + s$, where w and s are defined by

$$w := \begin{cases} (Qv_1) \circ \pi_1 & \text{on } X_1, \\ (K_D^{\mu_2} Qv_1) \circ \pi_2 & \text{on } X_2, \end{cases}$$

and

$$s := \begin{cases} (QK_D^{\mu_1} v_2) \circ \pi_1 & \text{on } X_1, \\ (Tv_2) \circ \pi_2 & \text{on } X_2, \end{cases}$$

with $v_j := v \circ i_j - K_D^{\mu_j} (v \circ i_k)$, $j \neq k$, $j, k \in \{1, 2\}$.

Remark 4.1. (1) Note that if $v = w' + s'$ is another decomposition of v with

$$w' := \begin{cases} (Qt_1) \circ \pi_1 & \text{on } X_1, \\ (K_D^{\mu_2} Qt_1) \circ \pi_2 & \text{on } X_2, \end{cases}$$

and

$$s' := \begin{cases} (QK_D^{\mu_1} t_2) \circ \pi_1 & \text{on } X_1, \\ (Tt_2) \circ \pi_2 & \text{on } X_2, \end{cases}$$

where t_j , $j = 1, 2$, are L_j -harmonic on D , then $t_1 = v_1$ and $t_2 = v_2$.

(2) In the classical case, for any biharmonic couple (h_1, h_2) the following holds:

$$(h_1, h_2) = (t, 0) + (K_D^{\mu_1} h_2, h_2),$$

where t is a harmonic function on D . Note that $(K_D^{\mu_1} h_2, h_2)$ is a pure biharmonic couple (see [3] and [21], [22]).

Corollary 4.1. *Let v be a positive minimal harmonic function on X such that $K_D^{\mu_k} K_D^{\mu_j}(v \circ i_k)$, $j \neq k$, $j, k \in \{1, 2\}$, is bounded. Then $v = \alpha w$ or $v = \beta s$, where α and β are positive constants; w and s are defined as in Proposition 4.1.*

Proposition 4.2. *Let v be a positive function on X such that $K_D^{\mu_j}(v \circ i_k)$ is finite and $K_D^{\mu_k} K_D^{\mu_j}(v \circ i_k)$, $j \neq k$, $j, k \in \{1, 2\}$, is bounded. The following statements are equivalent.*

- (1) v is a minimal harmonic function on X .
- (2) v_1 is a positive minimal L_1 -harmonic function on D , or v_2 is a positive minimal L_2 -harmonic function on D , where $v_j := v \circ i_j - K_D^{\mu_j}(v \circ i_k)$.

PROOF: Let v be a positive minimal harmonic function on X . Then we have $v = \alpha w$ or $v = \beta s$ by Corollary 4.1.

We shall show that if $v = \alpha w$, then v_1 is L_1 -minimal and if $v = \beta s$, then v_2 is L_2 -minimal.

(i) Case $v = \alpha w$:

Suppose that v_1 is not L_1 -minimal. Then there exist two L_1 -harmonic functions u_1 and u_2 such that $v_1 = u_1 + u_2$. So $v = \alpha f_1 + \alpha f_2$, with

$$f_1 = \begin{cases} (Qu_1) \circ \pi_1 & \text{on } X_1, \\ (K_D^{\mu_2} Qu_1) \circ \pi_2 & \text{on } X_2, \end{cases}$$

and

$$f_2 = \begin{cases} (Qu_2) \circ \pi_1 & \text{on } X_1, \\ (K_D^{\mu_2} Qu_2) \circ \pi_2 & \text{on } X_2. \end{cases}$$

It follows from Theorem 3.1 that f_1 and f_2 are harmonic on X . This contradicts that v is minimal.

(ii) Case $v = \beta s$:

Suppose that v_2 is not L_2 -minimal. Then there exist two L_2 -harmonic functions u_1 and u_2 such that $v_2 = u_1 + u_2$. Therefore $v = \beta s_1 + \beta s_2$, with

$$s_1 = \begin{cases} (QK_D^{\mu_1} u_1) \circ \pi_1 & \text{on } X_1, \\ (Tu_1) \circ \pi_2 & \text{on } X_2, \end{cases}$$

and

$$s_2 = \begin{cases} (QK_D^{\mu_1} u_2) \circ \pi_1 & \text{on } X_1, \\ (Tu_2) \circ \pi_2 & \text{on } X_2. \end{cases}$$

It follows from Theorem 3.1 that s_1 and s_2 are harmonic on X . This contradicts that v is minimal.

Conversely, suppose that v_1 is L_1 -minimal and let us show that v is minimal. Assume the contrary and put $v = g_1 + g_2$, where g_1 and g_2 are harmonic functions

on X . Then, from Proposition 4.1, there exist two L_1 -harmonic functions s_1 and s_2 , and two L_2 -harmonic functions w_1 and w_2 such that

$$g_1 = \begin{cases} (Qs_1) \circ \pi_1 + (QK_D^{\mu_1}w_1) \circ \pi_1 & \text{on } X_1, \\ (K_D^{\mu_2}Qs_1) \circ \pi_2 + (Tw_1) \circ \pi_2 & \text{on } X_2, \end{cases}$$

and

$$g_2 = \begin{cases} (Qs_2) \circ \pi_1 + (QK_D^{\mu_1}w_2) \circ \pi_1 & \text{on } X_1, \\ (K_D^{\mu_2}Qs_2) \circ \pi_2 + (Tw_2) \circ \pi_2 & \text{on } X_2. \end{cases}$$

Therefore the function $g_1 + g_2$ is defined on X_1 by

$$g_1 + g_2 := (Q(s_1 + s_2)) \circ \pi_1 + (QK_D^{\mu_1}(w_1 + w_2)) \circ \pi_1$$

and on X_2 by

$$g_1 + g_2 := (K_D^{\mu_2}Q(s_1 + s_2)) \circ \pi_2 + (T(w_1 + w_2)) \circ \pi_2.$$

We deduce, from Proposition 4.1 and Remark 4.1.1, that $v_1 = s_1 + s_2$, which leads to a contradiction because v_1 is L_1 -minimal.

In the same way, we suppose that v_2 is an L_2 -minimal function and we show that v is a minimal function. □

By using the fact that any positive minimal L_j -harmonic function on D is proportional to $g^j(\cdot, y)$, $y \in \Delta_j$ (see [10]), w and s from Corollary 4.1 can be given more precisely.

Corollary 4.2. *Let v be a positive minimal harmonic function defined on X such that the function $K_D^{\mu_k}K_D^{\mu_j}(v \circ i_k)$, $j \neq k$, $j, k \in \{1, 2\}$, is bounded. Then*

$$v = \alpha w \quad \text{or} \quad v = \beta s,$$

with

$$w := \begin{cases} (Qg^1(\cdot, y)) \circ \pi_1 & \text{on } X_1, y \in \Delta_1, \\ (K_D^{\mu_2}Qg^1(\cdot, y)) \circ \pi_2 & \text{on } X_2, y \in \Delta_1, \end{cases}$$

and

$$s := \begin{cases} (QK_D^{\mu_1}g^2(\cdot, y)) \circ \pi_1 & \text{on } X_1, y \in \Delta_2, \\ (Tg^2(\cdot, y)) \circ \pi_2, & \text{on } X_2, y \in \Delta_2. \end{cases}$$

PROOF: This result follows immediately from Proposition 4.2 and Corollary 4.1. □

Remark 4.2. Note that $K_D^{\mu_j}(v \circ i_k) < \infty, j \neq k, j, k \in \{1, 2\}$, because v is a positive harmonic function on X .

Consider the family of mappings on the real vector space $\mathcal{H}(X)$ defined by

$$\varphi_K : \begin{cases} \mathcal{H}(X) & \longrightarrow \mathbb{R}^+, \\ h & \longmapsto \varphi_K(h), \end{cases}$$

where

$$\varphi_K(h) = \sup_{x \in K} (|h \circ i_1(x)| + |h \circ i_2(x)|),$$

and K is a compact subset of D . (φ_K) is a family of semi-norms on $\mathcal{H}(X)$ and these semi-norms define a topology that makes $\mathcal{H}(X)$ a metrizable topological space. It follows that this space is locally convex.

The cone $\mathcal{H}^+(X) = \{h \in \mathcal{H}(X) : h \geq 0\}$ defines on $\mathcal{H}(X)$ an order relation called specific order:

$$h_1 \prec h_2 \iff h_2 = h_1 + g, \quad g \in \mathcal{H}^+(X).$$

Equipped with this order, $\mathcal{H}^+(X)$ is a lattice. The minimal harmonic functions are the points of the extreme generatrices of $\mathcal{H}^+(X)$. We recall that a base of $\mathcal{H}^+(X)$ is the intersection of $\mathcal{H}^+(X)$ with a closed hyperplane.

Let us consider the set

$$B := \{h \in \mathcal{H}^+(X) : (h \circ i_1)(x_o) + (h \circ i_2)(x_o) = 1\}, \quad x_o \in D.$$

B is a compact base of the cone $\mathcal{H}^+(X)$. Indeed, the mapping

$$\phi_{x_o} : \begin{cases} \mathcal{H}^+(X) & \longrightarrow \mathbb{R}, \\ h & \longmapsto (h \circ i_1)(x_o) + (h \circ i_2)(x_o) = 1 \end{cases}$$

is a continuous linear form. Then it defines a closed hyperplane B such that the origin $0 \notin B$. Then, B is equicontinuous at any point $x \in X$. So, we conclude, by Ascoli's theorem, that B is compact. Note that $\mathcal{H}^+(X) = \mathbb{R}^+ B$. Let $\mathcal{E}(B)$ denote the set of all extreme points of $\mathcal{H}^+(X)$ belonging to B (see [11]). Moreover, using Corollary 4.2, we have

$$\mathcal{E}(B) = \mathcal{E}_1(B) \cup \mathcal{E}_2(B),$$

where

$$\mathcal{E}_1(B) = \left\{ h \in \mathcal{E}(B) : \exists \alpha \in \mathbb{R}^+, \exists y \in \Delta_1 : h = \begin{cases} (\alpha Qg^1(\cdot, y)) \circ \pi_1 & \text{on } X_1 \\ (\alpha K_D^{\mu_2} Qg^1(\cdot, y)) \circ \pi_2 & \text{on } X_2 \end{cases} \right\}$$

and

$$\mathcal{E}_2(B) = \left\{ h \in \mathcal{E}(B) : \exists \beta \in \mathbb{R}^+, \exists y \in \Delta_2 : h = \begin{cases} (\beta QK_D^{\mu_1} g^2(\cdot, y)) \circ \pi_1 & \text{on } X_1 \\ (\beta Tg^2(\cdot, y)) \circ \pi_2 & \text{on } X_2 \end{cases} \right\}.$$

We recall the following results which are useful for showing the uniqueness of an integral representation (see [16]).

Definition 4.2 ([16]). Let Γ a closed convex cone. A mapping $\ell : \lambda \mapsto e_\lambda$ of a separated topological space Ω in $\mathcal{E}(\Gamma)$ is called a *parametrization* of $\mathcal{E}(\Gamma)$, if any element $\gamma \in \mathcal{E}(\Gamma)$ is proportional to a unique element e_λ . It is called *admissible* if it is continuous and the inverse mapping $\mathcal{E}(\Gamma) \rightarrow \Omega$ is universally measurable.

Theorem A ([16]). *Let a closed cone convex Γ and an admissible parametrization ℓ of $\mathcal{E}(\Gamma)$ be given. For any $\gamma \in \Gamma$, there exist a positive Radon measure μ on Ω such that*

$$\gamma = \int_{\Omega} e_\lambda d\mu(\lambda).$$

Theorem B ([16]). *The measure μ given by Theorem A is unique for any $\gamma \in \Gamma$, if and only if the cone Γ is a lattice.*

Theorem 4.1. *If $g^1(x, \cdot), x \in D$, separates Δ_1 and $g^2(x, \cdot), x \in D$, separates Δ_2 , then for any positive harmonic function v on X such that the function $K_D^{\mu_k} K_D^{\mu_j} (v \circ i_k), j \neq k, j, k \in \{1, 2\}$, is bounded, there exist two unique measures ν_1 and ν_2 supported respectively by Δ_1 and Δ_2 such that v can be represented on X_1 by*

$$v = \int_{\Delta_1} (Qg^1(\cdot, y)) \circ \pi_1 d\nu_1(y) + \int_{\Delta_2} (QK_D^{\mu_1} g^2(\cdot, y)) \circ \pi_1 d\nu_2(y)$$

and on X_2 by

$$v = \int_{\Delta_1} (K_D^{\mu_2} Qg^1(\cdot, y)) \circ \pi_2 d\nu_1(y) + \int_{\Delta_2} (Tg^2(\cdot, y)) \circ \pi_2 d\nu_2(y).$$

PROOF: If $v = 0$, we have $\nu_1 = \nu_2 = 0$.

If $v \neq 0$, we may assume without loss of generality that $v \in B$. Consider the mapping

$$\Psi : \begin{cases} \Delta_1 \cup \Delta_2 \longrightarrow \mathcal{E}(B) \\ y \longmapsto \Psi(y) \end{cases}$$

where $\Psi(y)$ is defined by

$$\Psi(y) := \begin{cases} (Qg^1(\cdot, y)) \circ \pi_1 & \text{on } X_1 \\ (K_D^{\mu_2} Qg^1(\cdot, y)) \circ \pi_2 & \text{on } X_2 \end{cases}, \quad y \in \Delta_1,$$

$$\Psi(y) := \begin{cases} (QK_D^{\mu_1} g^2(\cdot, y)) \circ \pi_1 & \text{on } X_1 \\ (Tg^2(\cdot, y)) \circ \pi_2 & \text{on } X_2 \end{cases}, \quad y \in \Delta_2.$$

The mapping Ψ is bijective because $g^1(x, \cdot)$ and $g^2(x, \cdot)$ separate Δ_1 and Δ_2 , respectively. Ψ and its inverse Ψ^{-1} are continuous because g^1 and g^2 are continuous on $\Delta \times D$. Then there exists, by Theorem B, a unique measure ν supported by $\Delta_1 \cup \Delta_2$ such that

$$v = \int_{\Delta_1 \cup \Delta_2} \Psi(y) d\nu(y).$$

Let $\nu_j, j = 1, 2$, be the restriction of the measure ν to Δ_j . Then v may be written on X_1 as

$$v = \int_{\Delta_1} (Qg^1(\cdot, y)) \circ \pi_1 d\nu_1(y) + \int_{\Delta_2} (QK_D^{\mu_1}g^2(\cdot, y)) \circ \pi_1 d\nu_2(y)$$

and on X_2 as

$$v = \int_{\Delta_1} (K_D^{\mu_2}Qg^1(\cdot, y)) \circ \pi_2 d\nu_1(y) + \int_{\Delta_2} (Tg^2(\cdot, y)) \circ \pi_2 d\nu_2(y).$$

□

Let $t_i, i = 1, 2$, be two positive L_i -harmonic functions on D such that the function $K_D^{\mu_j}t_k$ is finite and the function $K_D^{\mu_k}K_D^{\mu_j}t_k, j \neq k, j, k \in \{1, 2\}$, is bounded on D . By [10] and [12], there exists a unique measure ν_{t_j} , supported by Δ_j , such that $t_j = \int_{\Delta_j} g^j(\cdot, y) d\nu_{t_j}(y), j = 1, 2$. We consider the harmonic function w from Theorem 3.1 defined on X by

$$w := \begin{cases} (Qt_1 + QK_D^{\mu_1}t_2) \circ \pi_1 & \text{on } X_1, \\ (K_D^{\mu_2}Qt_1 + Tt_2) \circ \pi_2 & \text{on } X_2. \end{cases}$$

Corollary 4.3. *If the functions $g^j(x, \cdot), x \in D$, separate $\Delta_j, j = 1, 2$, then w is written on X_1 by*

$$w = \int_{\Delta_1} (Qg^1(\cdot, y)) \circ \pi_1 d\nu_{t_1}(y) + \int_{\Delta_2} (QK_D^{\mu_1}g^2(\cdot, y)) \circ \pi_1 d\nu_{t_2}(y),$$

and on X_2 by

$$w = \int_{\Delta_1} (K_D^{\mu_2}Qg^1(\cdot, y)) \circ \pi_2 d\nu_{t_1}(y) + \int_{\Delta_2} (Tg^2(\cdot, y)) \circ \pi_2 d\nu_{t_2}(y).$$

PROOF: It suffices to replace $t_j, j = 1, 2$, with their Martin representations in the expression of w , and the result follows from the uniqueness of the measures ν_j in Theorem 4.1. □

Remark 4.3. By Corollary 4.3, we have $\nu_{t_j}(\Delta \setminus \Delta_j) = 0$, thus $\nu_{t_j}(\Delta \setminus (\Delta_1 \cup \Delta_2)) = 0, j = 1, 2$.

5. Dirichlet problem on the Martin boundary associated with (S)

Given a couple of functions (u_1, u_2) defined on Δ , the Dirichlet problem on Δ consists to find a couple of functions (h_1, h_2) solving the system (S) such that

$$\lim_{x \rightarrow y} h_i(x) = u_i(y) \quad \forall y \in \Delta.$$

The couple (u_1, u_2) can be identified with a function f on $\bar{\Delta} := \bigcup_{j=1}^2 \Delta \times \{j\}$ such that $f \circ i_j = u_j$, where $i_j, j = 1, 2$, denote always the mappings of Δ in $\Delta \times \{j\}$ defined by $i_j(z) := (z, j), z \in \Delta$. The Dirichlet problem may be stated as follows: for a given function f defined on $\bar{\Delta}$, determine, if possible, a harmonic function H_f on X such that $H_f(x) \rightarrow f(y)$ as $x \rightarrow y$ for each $y \in \bar{\Delta}$. As in harmonic and biharmonic cases, there are some examples where there is no solution of this problem. In this section, we will discuss the Perron-Wiener-Brelot (PWB) approach to the Dirichlet problem. To this end, we give the following definition.

Definition 5.1. Let h_1 (resp. h_2) be a strictly positive L_1 -harmonic (resp. L_2 -harmonic) function on D , and let h be the function defined on X by

$$h := \begin{cases} h_1 \circ \pi_1 & \text{on } X_1, \\ h_2 \circ \pi_2 & \text{on } X_2. \end{cases}$$

A function v on X is called *h-harmonic* (resp. *h-hyperharmonic*, *h-superharmonic*) on X if and only if the function u defined on X by

$$u := \begin{cases} (h_1(v \circ i_1)) \circ \pi_1 & \text{on } X_1, \\ (h_2(v \circ i_2)) \circ \pi_2 & \text{on } X_2 \end{cases}$$

is harmonic (resp. hyperharmonic, superharmonic) on X .

We also define the upper and lower class associated with a function defined on $\bar{\Delta}$. Let f be a function defined on $\bar{\Delta}$ and let h be a function defined on X as in Definition 5.1. We define:

$$\bar{U}_f := \{v : v \text{ is } h\text{-hyperharmonic and bounded from below on } X \text{ and} \\ \liminf_{x \rightarrow y} v(x) \geq f(y), \forall y \in \bar{\Delta}\}$$

and

$$U_f := \{s : s \text{ is } h\text{-hypoharmonic and bounded from above on } X \text{ and} \\ \limsup_{x \rightarrow y} v(x) \leq f(y), \forall y \in \bar{\Delta}\}.$$

We note that \bar{U}_f and \underline{U}_f are never empty since they contain the constant functions $+\infty$ and $-\infty$ respectively, and that $\bar{U}_f = -\underline{U}_{-f}$. Put

$$\bar{H}_f := \inf \bar{U}_f \quad \text{and} \quad \underline{H}_f := \sup \underline{U}_f.$$

f is called *h-resolutive* if \bar{H}_f and \underline{H}_f are equal and h -harmonic on X . If f is h -resolutive, then we define $H_f^h := \bar{H}_f = \underline{H}_f$ and call H_f^h the *PWB-solution of the Dirichlet problem on X with boundary function f* . If $f \circ i_j$ is h_j -resolutive on Δ , we call $H_{f \circ i_j}^{h_j}$ the *PWB-solution of Dirichlet problem on D associated with $f \circ i_j$, $j = 1, 2$* .

Further properties of PWB solutions.

Let f and g be two functions defined on $\bar{\Delta}$. Then we have

- (i) $\underline{H}_f^h = -\bar{H}_{-f}^h$.
- (ii) $\underline{H}_f^h \leq \bar{H}_f^h$.
- (iii) $\underline{H}_f^h \leq \underline{H}_g^h$ and $\bar{H}_f^h \leq \bar{H}_g^h$ if $f \leq g$.
- (iv) Let f, g be two h -resolutive functions and $\alpha \in \mathbb{R}$. Then $f + g$ and αf are h -resolutive and

$$H_{f+g}^h = H_f^h + H_g^h, \quad H_{\alpha f}^h = \alpha H_f^h.$$

- (v) If $\underline{U}_f \cap (-S(X)) \neq \emptyset$ (resp. $\bar{U}_f \cap S(X) \neq \emptyset$), then the function \bar{H}_f^h (resp. \underline{H}_f^h) is identically ∞ , or h -harmonic on X .

Let f be a positive function on $\bar{\Delta}$ such that $f \circ i_2 = 0$ and w the function defined on X by

$$w := \begin{cases} (\frac{1}{h_1} Q(h_1 \cdot \bar{H}_{f \circ i_1}^{h_1})) \circ \pi_1 & \text{on } X_1, \\ (\frac{1}{h_2} K_D^{\mu_2} Q(h_1 \cdot \bar{H}_{f \circ i_1}^{h_1})) \circ \pi_2 & \text{on } X_2. \end{cases}$$

We have $\bar{H}_f^h \leq w$. Indeed, it follows from Corollary 3.1 that w is a positive h -hyperharmonic function on X and moreover, we have

$$\liminf_{x \rightarrow y} (w \circ i_1)(x) \geq (f \circ i_1)(y), \quad \text{for all } y \in \Delta$$

and

$$\liminf_{x \rightarrow y} (w \circ i_2)(x) \geq 0, \quad \text{for all } y \in \Delta.$$

Hence, $w \in \bar{U}_f$. Thus $\bar{H}_f^h \leq w$ and therefore if $\bar{H}_f^h = +\infty$ then $w = +\infty$. If $\bar{H}_f^h < \infty$, we have

Lemma 5.1. *Let f be a positive function on $\bar{\Delta}$ such that $f \circ i_2 = 0$ and $K_D^{\mu_1} K_D^{\mu_2} (h_1(\bar{H}_f^h \circ i_1))$ is bounded on D . Then we have*

$$\bar{H}_f^h = \begin{cases} (\frac{1}{h_1} Q(h_1 \bar{H}_{f \circ i_1}^{h_1})) \circ \pi_1 & \text{on } X_1, \\ (\frac{1}{h_2} K_D^{\mu_2} Q(h_1 \bar{H}_{f \circ i_1}^{h_1})) \circ \pi_2 & \text{on } X_2. \end{cases}$$

PROOF: It suffices to show that $w \leq \bar{H}_f^h$.

(a) Let us show that $w \circ i_1 \leq \bar{H}_f^h \circ i_1$.

It follows from property (v) of PWB solutions that the function \bar{H}_f^h is h -harmonic on X . Then the function

$$\bar{u} := \begin{cases} (h_1(\bar{H}_f^h \circ i_1)) \circ \pi_1 & \text{on } X_1, \\ (h_2(\bar{H}_f^h \circ i_2)) \circ \pi_2 & \text{on } X_2 \end{cases}$$

is a positive harmonic function on X , and by Corollary 2.1, the functions $\bar{u}_j = h_j(\bar{H}_f^h \circ i_j) - K_D^{\mu_j} (h_k(\bar{H}_f^h \circ i_k))$, $j, k \in \{1, 2\}$, $j \neq k$ are positive and L_j -harmonic on D . Put $v_j := \frac{1}{h_j} \bar{u}_j$. On the one hand, we have

$$K_D^{\mu_2} (h_1(\bar{H}_f^h \circ i_1)) \leq (h_2(\bar{H}_f^h \circ i_2)),$$

hence

$$K_D^{\mu_1} K_D^{\mu_2} (h_1(\bar{H}_f^h \circ i_1)) \leq K_D^{\mu_1} (h_2(\bar{H}_f^h \circ i_2)),$$

i.e.

$$K_D^{\mu_1} K_D^{\mu_2} (h_1(\bar{H}_f^h \circ i_1)) \leq (h_1(\bar{H}_f^h \circ i_1) - h_1.v_1).$$

So,

$$Q(h_1.v_1) + QK_D^{\mu_1} K_D^{\mu_2} (h_1(\bar{H}_f^h \circ i_1)) \leq Q(h_1(\bar{H}_f^h \circ i_1)).$$

Since

$$QK_D^{\mu_1} K_D^{\mu_2} + I = Q,$$

we get

$$Q(h_1.v_1) + QK_D^{\mu_1} K_D^{\mu_2} (h_1(\bar{H}_f^h \circ i_1)) \leq QK_D^{\mu_1} K_D^{\mu_2} (h_1(\bar{H}_f^h \circ i_1)) + h_1(\bar{H}_f^h \circ i_1).$$

Therefore,

$$(5.1.1) \quad Q(h_1.v_1) \leq h_1(\bar{H}_f^h \circ i_1).$$

On the other hand,

$$\begin{aligned} \liminf_{x \rightarrow y} v_1(x) &= \liminf_{x \rightarrow y} (\bar{H}_f^h \circ i_1 - \frac{1}{h_1} K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)))(x) \\ &\geq (f \circ i_1)(y) - \limsup_{x \rightarrow y} (\frac{1}{h_1} K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)))(x) \end{aligned}$$

for all $y \in \Delta$. Since

$$\begin{aligned} \limsup_{x \rightarrow y} (\frac{1}{h_1} K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)))(x) \\ \leq \int_D \limsup_{x \rightarrow y} \frac{1}{h_1(x)} G_1(x, z) h_2(z) (\bar{H}_f^h \circ i_2)(z) d\mu_1(z), \end{aligned}$$

and $\limsup_{x \rightarrow y} \frac{1}{h_1(x)} G_1(x, z) = 0$ ν_{h_1} -a.e. on Δ_1 , where ν_{h_1} is the measure associated with h_1 in the Martin representation ([13, p. 218]), we have, by Remark 4.3, $\nu_{h_1}(\Delta \setminus \Delta_1) = 0$. Hence $\limsup_{x \rightarrow y} \frac{1}{h_1(x)} G_1(x, z) = 0$ ν_{h_1} -a.e. on Δ . Thus $\liminf_{x \rightarrow y} v_1(x) \geq (f \circ i_1)(y)$ ν_{h_1} -a.e. on Δ . Hence v_1 is a positive $h_1 - L_1$ -hyperharmonic function on D and $\liminf_{x \rightarrow y} v_1(x) \geq (f \circ i_1)(y)$ ν_{h_1} -a.e. on Δ . So

$$(5.1.2) \quad v_1 \geq \bar{H}_{f \circ i_1}^{h_1}.$$

Thus, by (5.1.1), we have

$$Q(h_1 \bar{H}_{f \circ i_1}^{h_1}) \leq (h_1(\bar{H}_f^h \circ i_1)).$$

(b) Let us show that $w \circ i_2 \leq (\bar{H}_f^h \circ i_2)$.

It follows from (a) that

$$Q(h_1 \bar{H}_{f \circ i_1}^{h_1}) \leq (h_1(\bar{H}_f^h \circ i_1)).$$

Then,

$$K_D^{\mu_2} Q(h_1 \bar{H}_{f \circ i_1}^{h_1}) \leq K_D^{\mu_2} (h_1(\bar{H}_f^h \circ i_1)) \leq (h_2(\bar{H}_f^h \circ i_2)).$$

This finishes the proof. □

Remark 5.1. The result of Lemma 5.1 is still valid if instead of the assumption $K_D^{\mu_1} K_D^{\mu_2} (h_1(\bar{H}_f^h \circ i_1))$ is bounded, we suppose only that $Q(h_1(\bar{H}_f^h \circ i_1))$ is finite.

Let f be a positive function on $\bar{\Delta}$ such that $f \circ i_1 = 0$ and \tilde{w} the function defined on X by

$$\tilde{w} := \begin{cases} (\frac{1}{h_1} Q K_D^{\mu_1}(h_2 \bar{H}_{f \circ i_2}^{h_2})) \circ \pi_1 & \text{on } X_1, \\ (\frac{1}{h_2} T(h_2 \bar{H}_{f \circ i_2}^{h_2})) \circ \pi_2 & \text{on } X_2. \end{cases}$$

We have $\bar{H}_f^h \leq \tilde{w}$. Therefore if $\bar{H}_f^h = +\infty$, then $\tilde{w} = +\infty$. If $\bar{H}_f^h < \infty$, we have:

Lemma 5.2. *Let f be a positive function on $\bar{\Delta}$ such that $f \circ i_1 = 0$ and $K_D^{\mu_2} K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2))$ is bounded on D . Then*

$$\bar{H}_f^h = \begin{cases} (\frac{1}{h_1} Q K_D^{\mu_1}(h_2 \bar{H}_{f \circ i_2}^{h_2})) \circ \pi_1 & \text{on } X_1, \\ (\frac{1}{h_2} T(h_2 \bar{H}_{f \circ i_2}^{h_2})) \circ \pi_2 & \text{on } X_2. \end{cases}$$

PROOF: It suffices to show that $\tilde{w} \leq \bar{H}_f^h$.

(a) Let us show that $\tilde{w} \circ i_1 \leq \bar{H}_f^h \circ i_1$.

By the property (v) of PWB solutions, the function \bar{H}_f^h is h -harmonic on X . Then the function

$$\bar{u} := \begin{cases} (h_1(\bar{H}_f^h \circ i_1)) \circ \pi_1 & \text{on } X_1, \\ (h_2(\bar{H}_f^h \circ i_2)) \circ \pi_2 & \text{on } X_2 \end{cases}$$

is a positive harmonic function on X and by Corollary 2.1, $\bar{u}_j = h_j(\bar{H}_f^h \circ i_j) - K_D^{\mu_j}(h_k(\bar{H}_f^h \circ i_k))$, $j, k \in \{1, 2\}$, $j \neq k$, are positive and L_j -harmonic functions on D . Put $v_j := \frac{1}{h_j} \bar{u}_j$. On the one hand, we have

$$K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)) \leq (h_1(\bar{H}_f^h \circ i_1)),$$

hence

$$K_D^{\mu_1}(h_2 v_2 + K_D^{\mu_2}(h_1(\bar{H}_f^h \circ i_1))) \leq h_1(\bar{H}_f^h \circ i_1)$$

and

$$Q K_D^{\mu_1}(h_2 v_2) + Q K_D^{\mu_1} K_D^{\mu_2}(h_1(\bar{H}_f^h \circ i_1)) \leq Q(h_1(\bar{H}_f^h \circ i_1)).$$

Since

$$Q K_D^{\mu_1} K_D^{\mu_2} + I = Q,$$

we get

$$Q K_D^{\mu_1}(h_2 \cdot v_2) \leq h_1(\bar{H}_f^h \circ i_1).$$

As in the proof of Lemma 5.1, we show that $\liminf_{x \rightarrow y} v_2(x) \geq (f \circ i_2)(y)$ ν_{h_2} -a.e. on Δ . Since v_2 is a positive h_2 - L_2 -hyperharmonic function and $\liminf_{x \rightarrow y} v_2(x) \geq (f \circ i_2)(y)$, ν_{h_2} -a.e. on Δ , we obtain

$$(5.1.2) \quad v_2 \geq \bar{H}_{f \circ i_2}^{h_2},$$

hence

$$Q K_D^{\mu_1}(h_2 \bar{H}_{f \circ i_2}^{h_2}) \leq (h_1(\bar{H}_f^h \circ i_1)).$$

(b) Let us show that $\tilde{w} \circ i_2 \leq (\bar{H}_f^h \circ i_2)$. We have

$$K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)) \leq h_1(\bar{H}_f^h \circ i_1).$$

So

$$K_D^{\mu_2} K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)) \leq K_D^{\mu_2}(h_1(\bar{H}_f^h \circ i_1)) = h_2(\bar{H}_f^h \circ i_2) - h_2 v_2.$$

Hence

$$T(h_2.v_2) + T K_D^{\mu_2} K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)) \leq T(h_2(\bar{H}_f^h \circ i_2)).$$

Since

$$T K_D^{\mu_2} K_D^{\mu_1} + I = T,$$

we get

$$T(h_2 \bar{H}_{f \circ i_2}^{h_2}) \leq (h_2(\bar{H}_f^h \circ i_2)).$$

□

Remark 5.2. The result of Lemma 5.2 is still valid if instead of the assumption $K_D^{\mu_2} K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2))$ is bounded, we suppose only that $T(h_2(\bar{H}_f^h \circ i_2))$ is finite.

Let f be a positive function on $\bar{\Delta}$ and let w' be the function defined on X by

$$w' := \begin{cases} \frac{1}{h_1}(Q(h_1 \bar{H}_{f \circ i_1}^{h_1}) + Q K_D^{\mu_1}(h_2 \bar{H}_{f \circ i_2}^{h_2})) \circ \pi_1 & \text{on } X_1, \\ \frac{1}{h_2}(K_D^{\mu_2} Q(h_1 \bar{H}_{f \circ i_1}^{h_1}) + T(h_2 \bar{H}_{f \circ i_2}^{h_2})) \circ \pi_2 & \text{on } X_2. \end{cases}$$

We have $\bar{H}_f^h \leq w'$. Therefore, if $\bar{H}_f^h = +\infty$ then $w' = +\infty$. If $\bar{H}_f^h < \infty$, we have

Proposition 5.1. Let f be a positive function on $\bar{\Delta}$ such that $K_D^{\mu_j} K_D^{\mu_k}(h_j(\bar{H}_f^h \circ i_j))$ is bounded on D , $j, k \in \{1, 2\}$, $j \neq k$. Then we have

$$\bar{H}_f^h = \begin{cases} \frac{1}{h_1}(Q(h_1 \bar{H}_{f \circ i_1}^{h_1}) + Q K_D^{\mu_1}(h_2 \bar{H}_{f \circ i_2}^{h_2})) \circ \pi_1 & \text{on } X_1, \\ \frac{1}{h_2}(K_D^{\mu_2} Q(h_1 \bar{H}_{f \circ i_1}^{h_1}) + T(h_2 \bar{H}_{f \circ i_2}^{h_2})) \circ \pi_2 & \text{on } X_2. \end{cases}$$

PROOF: It suffices to show that $w' \leq \bar{H}_f^h$.

(a) Let us show that $w' \circ i_1 \leq \bar{H}_f^h \circ i_1$.

By the property (v) of PWB solutions, the function \bar{H}_f^h is h -harmonic on X . Then the function

$$\bar{u} := \begin{cases} (h_1(\bar{H}_f^h \circ i_1)) \circ \pi_1 & \text{on } X_1, \\ (h_2(\bar{H}_f^h \circ i_2)) \circ \pi_2 & \text{on } X_2 \end{cases}$$

is a positive harmonic on X and by Corollary 2.1, $\bar{u}_j = h_j(\bar{H}_f^h \circ i_j) - K_D^{\mu_j}(h_k(\bar{H}_f^h \circ i_k))$, $j, k \in \{1, 2\}$, $j \neq k$, are positive L_j -harmonic on D . Put $v_j = \frac{1}{h_j}\bar{u}_j$. On the one hand,

$$h_1.v_1 + K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)) = h_1(\bar{H}_f^h \circ i_1)$$

and

$$h_2.v_2 + K_D^{\mu_2}(h_1(\bar{H}_f^h \circ i_1)) = h_2(\bar{H}_f^h \circ i_2).$$

Hence

$$Q(h_1.v_1) + QK_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)) = Q(h_1(\bar{H}_f^h \circ i_1))$$

and

$$QK_D^{\mu_1}(h_2.v_2) + QK_D^{\mu_1}K_D^{\mu_2}(h_1(\bar{H}_f^h \circ i_1)) = QK_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)).$$

Since

$$QK_D^{\mu_1}K_D^{\mu_2} + I = Q,$$

we have

$$Q(h_1.v_1) + QK_D^{\mu_1}(h_2.v_2) = h_1(\bar{H}_f^h \circ i_1).$$

It follows from (5.1.2) and (5.2.1) that

$$Q(h_1\bar{H}_{f \circ i_1}^{h_1}) + QK_D^{\mu_1}(h_2\bar{H}_{f \circ i_2}^{h_2}) \leq h_1(\bar{H}_f^h \circ i_1).$$

Similarly, we show that

$$\frac{1}{h_2}(K_D^{\mu_2}Q(h_1\bar{H}_{f \circ i_1}^{h_1}) + T(h_2\bar{H}_{f \circ i_2}^{h_2})) \leq h_2(\bar{H}_f^h \circ i_2).$$

□

Remark 5.3. The result of Proposition 5.1 is still valid if instead of the assumption $K_D^{\mu_j}K_D^{\mu_k}(h_j(\bar{H}_f^h \circ i_j))$ is bounded on D , $j, k \in \{1, 2\}$, $j \neq k$, we suppose that $Q(h_1(\bar{H}_f^h \circ i_1)) < \infty$ and $T(h_2(\bar{H}_f^h \circ i_2)) < \infty$.

***h*-negligible sets.**

Definition 5.2. Let e be a subset of $\bar{\Delta}$. e is called *h-negligible* if $\bar{H}_{1_e}^h = 0$, where 1_e is the indicator of the set e .

Let \tilde{e} be a subset of Δ . \tilde{e} is called *h_j-negligible* if and only if $\bar{H}_{1_{\tilde{e}}}^{h_j} = 0$, $j = 1, 2$.

Proposition 5.2. Let $e \subset \bar{\Delta} = (\Delta \times \{1\}) \cup (\Delta \times \{2\})$ be such that $e = (e_1 \times \{1\}) \cup (e_2 \times \{2\})$, where $e_j \subset \Delta$, $j = 1, 2$. The following are equivalent:

- (1) e is *h-negligible*;
- (2) e_j is *h_j-negligible*, $j = 1, 2$.

PROOF: Suppose that e is h -negligible; then $\bar{H}_{1_e}^h = 0$. By Proposition 5.1, we have

$$\bar{H}_{1_e}^h = \begin{cases} \frac{1}{h_1}(Q(h_1\bar{H}_{1_e\circ i_1}^{h_1}) + QK_D^{\mu_1}(h_2\bar{H}_{1_e\circ i_2}^{h_2})) \circ \pi_1 & \text{on } X_1, \\ \frac{1}{h_2}(K_D^{\mu_2}Q(h_1\bar{H}_{1_e\circ i_1}^{h_1}) + T(h_2\bar{H}_{1_e\circ i_2}^{h_2})) \circ \pi_2 & \text{on } X_2, \end{cases}$$

hence

$$Q(h_1\bar{H}_{1_e\circ i_1}^{h_1}) = -QK_D^{\mu_1}(h_2\bar{H}_{1_e\circ i_2}^{h_2}), \quad K_D^{\mu_2}Q(h_1\bar{H}_{1_e\circ i_1}^{h_1}) = -T(h_2\bar{H}_{1_e\circ i_2}^{h_2}).$$

Since the functions $h_j\bar{H}_{1_e\circ i_j}^{h_j}$, $j = 1, 2$, are positive, $\bar{H}_{1_e\circ i_j}^{h_j} = 0$, $j = 1, 2$. Since $1_e \circ i_j = 1_{e_j}$, $\bar{H}_{1_{e_j}}^{h_j} = 0$, i.e., the set e_j is h_j -negligible. The converse is obvious. □

Proposition 5.3. *Let f and \tilde{f} be two positive functions defined on $\bar{\Delta}$ such that $e = \{f \neq \tilde{f}\}$ is a h -negligible set. Then $\bar{H}_f^h = \bar{H}_{\tilde{f}}^h$.*

PROOF: We have $e = \{f \neq \tilde{f}\} = (e_1 \times \{1\}) \cup (e_2 \times \{2\})$, where $e_j = \{f \circ i_j \neq \tilde{f} \circ i_j\}$, $j = 1, 2$, and e is h -negligible. Then, by Proposition 5.2, e_j is h_j -negligible. Thus $\bar{H}_{f \circ i_j}^{h_j} = \bar{H}_{\tilde{f} \circ i_j}^{h_j}$, $j = 1, 2$. Therefore, by Proposition 5.1, $\bar{H}_f^h = \bar{H}_{\tilde{f}}^h$. □

Lemma 5.3. *Let f be a positive function on $\bar{\Delta}$ such that $K_D^{\mu_j}K_D^{\mu_k}(h_j(\bar{H}_f^h \circ i_j))$ is bounded on D , $j, k \in \{1, 2\}$, $j \neq k$. Then we have*

$$h_j\bar{H}_{f \circ i_j}^{h_j} = h_j(\bar{H}_f^h \circ i_j) - K_D^{\mu_j}(h_k(\bar{H}_f^h \circ i_k)).$$

PROOF: By Proposition 5.1, we have

$$\begin{cases} \bar{H}_f^h \circ i_1 = \frac{1}{h_1}(Q(h_1\bar{H}_{f \circ i_1}^{h_1}) + QK_D^{\mu_1}(h_2\bar{H}_{f \circ i_2}^{h_2})), \\ \bar{H}_f^h \circ i_2 = \frac{1}{h_2}(K_D^{\mu_2}Q(h_1\bar{H}_{f \circ i_1}^{h_1}) + T(h_2\bar{H}_{f \circ i_2}^{h_2})). \end{cases}$$

Then

$$\begin{cases} h_1\bar{H}_f^h \circ i_1 = (Q(h_1\bar{H}_{f \circ i_1}^{h_1}) + QK_D^{\mu_1}(h_2\bar{H}_{f \circ i_2}^{h_2})), \\ h_2\bar{H}_f^h \circ i_2 = (K_D^{\mu_2}Q(h_1\bar{H}_{f \circ i_1}^{h_1}) + T(h_2\bar{H}_{f \circ i_2}^{h_2})). \end{cases}$$

Hence

$$\begin{cases} K_D^{\mu_2}(h_1.\bar{H}_f^h \circ i_1) = K_D^{\mu_2}(Q(h_1\bar{H}_{f \circ i_1}^{h_1})) + K_D^{\mu_2}(QK_D^{\mu_1}(h_2\bar{H}_{f \circ i_2}^{h_2})), \\ h_2\bar{H}_f^h \circ i_2 = (K_D^{\mu_2}Q(h_1\bar{H}_{f \circ i_1}^{h_1}) + T(h_2\bar{H}_{f \circ i_2}^{h_2})). \end{cases}$$

Since \bar{H}_f^h is h -harmonic on X , $K_D^{\mu_2}(h_1(\bar{H}_f^h \circ i_1)) < \infty$. Thus,

$$h_2(\bar{H}_f^h \circ i_2) - K_D^{\mu_2}(h_1(\bar{H}_f^h \circ i_1)) = T(h_2\bar{H}_{f \circ i_2}^{h_2}) - K_D^{\mu_2}QK_D^{\mu_1}(h_2\bar{H}_{f \circ i_2}^{h_2}).$$

Since

$$T = K_D^{\mu_2}QK_D^{\mu_1} + I,$$

we get

$$h_2(\bar{H}_f^h \circ i_2) - K_D^{\mu_2}(h_1(\bar{H}_f^h \circ i_1)) = h_2\bar{H}_{f \circ i_2}^{h_2}.$$

Similarly, we show that

$$h_1(\bar{H}_f^h \circ i_1) - K_D^{\mu_1}(h_2(\bar{H}_f^h \circ i_2)) = h_1\bar{H}_{f \circ i_1}^{h_1}.$$

□

Theorem 5.1. *Let f be a positive function defined on $\bar{\Delta}$ such that $K_D^{\mu_j}K_D^{\mu_k}(h_j(\bar{H}_f^h \circ i_j))$ is bounded, $j \neq k, j, k \in \{1, 2\}$. The following are equivalent:*

- (a) f is h -resolutive;
- (b) (1) $f \circ i_j$ is h_j -resolutive on $\Delta, j = 1, 2$, and
 (2) $K_D^{\mu_k}(h_jH_{f \circ i_j}^{h_j})$ is finite, $j \neq k, j, k \in \{1, 2\}$.

PROOF: Suppose that (b) holds. Then the function $h_jH_{f \circ i_j}^{h_j}$ is L_j -harmonic, $j = 1, 2$. Moreover, we have

$$h_jH_{f \circ i_j}^{h_j} \leq h_j(\bar{H}_f^h \circ i_j).$$

Since $K_D^{\mu_j}K_D^{\mu_k}(h_j(\bar{H}_f^h \circ i_j))$ is bounded, $j \neq k, j, k \in \{1, 2\}$, $K_D^{\mu_j}K_D^{\mu_k}(h_jH_{f \circ i_j}^{h_j})$ is bounded, $j \neq k, j, k \in \{1, 2\}$. Hence, by Theorem 3.1, the function

$$\bar{H}_f^h = \begin{cases} \frac{1}{h_1}(Q(h_1H_{f \circ i_1}^{h_1}) + QK_D^{\mu_1}(h_2H_{f \circ i_2}^{h_2})) \circ \pi_1 & \text{on } X_1, \\ \frac{1}{h_2}(K_D^{\mu_2}Q(h_1H_{f \circ i_1}^{h_1}) + T(h_2H_{f \circ i_2}^{h_2})) \circ \pi_2 & \text{on } X_2 \end{cases}$$

is h -harmonic on X , moreover $\bar{H}_f^h = \underline{H}_f^h = H_f^h$, therefore f is h -resolutive.

Conversely, suppose that f is h -resolutive. Then $\bar{H}_f^h = \underline{H}_f^h = H_f^h$ and H_f^h is h -harmonic. On the one hand, it follows from Lemma 5.3 that

$$h_j\bar{H}_{f \circ i_j}^{h_j} = h_j(H_f^h \circ i_j) - K_D^{\mu_j}(h_k(H_f^h \circ i_k)),$$

and by Corollary 2.1, the function $H_{f \circ i_j}^{h_j}$ is $h_j - L_j$ -harmonic on D , i.e. $f \circ i_j$ is h_j -resolutive on Δ . On the other hand,

$$K_D^{\mu_k}(h_j H_{f \circ i_j}^{h_j}) \leq K_D^{\mu_k}(h_j(H_f^h \circ i_j)) \leq h_k H_f^h \circ i_k,$$

thus

$$K_D^{\mu_k}(h_j H_{f \circ i_j}^{h_j}) < \infty. \quad \square$$

REFERENCES

- [1] Benyaïche A., *On almost biharmonic functions in \mathbb{R}^n* , Publ. Math., Ec. Norm. Supér, Takaddoum **4** (1988), 47–53 (See Zbl. Math. 678).
- [2] Benyaïche A., *Distributions bi-sousharmoniques sur \mathbb{R}^n ($n \geq 2$)*, Math. Bohem. **119** (1994), no. 1, 1–13.
- [3] Benyaïche A., *Mesures de représentation sur les espaces biharmoniques*, Proc. ICPT 91, Kluwer Acad. Publ., Dordrecht, 1994, pp. 171–178.
- [4] Benyaïche A., Ghiate S., *Propriété de moyenne restreinte associée à un système d'E.D.P.*, Rend. Accad. Naz. Sci. XL Mem. Mat. Appl. (1) **27** (2003), 125–143.
- [5] Benyaïche A., Ghiate S., *Frontière de Martin biharmonique*, preprint, 2000.
- [6] Benyaïche A., El Gourari A., *Caractérisation des ensembles essentiels*, J. Math. Soc. Japan **54** (2002), no. 2, 467–486.
- [7] Bliedtner J., Hansen W., *Potential Theory: An Analytic and Probabilistic Approach to Balayage*, Universitext, Springer, Berlin, 1986.
- [8] Boukricha A., *Espaces biharmoniques*, Théorie du potentiel (Orsay, 1983), Lecture Notes in Math. 1096, Springer, Berlin, 1984, pp. 116–149.
- [9] Bouleau N., *Espaces biharmoniques et couplage de processus de Markov*, J. Math. Pures Appl. **59** (1980), 187–240.
- [10] Brelot M., *On Topologies and Boundaries in Potential Theory*, Lecture Notes in Math. 175, Springer, Berlin, 1971.
- [11] Brelot M., *Éléments de la théorie classique du potentiel*, 4ème édition, Centre de Documentation Universitaire, Paris, 1969.
- [12] Constantinescu C., Cornea A., *Potential Theory on Harmonic Spaces*, Springer, New York-Heidelberg, 1972.
- [13] Doob J.L., *Classical Potential Theory and its Probabilistic Counterpart*, Springer, New York, 1984.
- [14] Hansen W., *Modification of balayage spaces by transitions with application to coupling of PDE's*, Nagoya Math. J. **169** (2003), 77–118.
- [15] Helms L.L., *Introduction to Potential Theory*, Wiley, New York, 1969.
- [16] Phelps R.R., *Lectures on Choquet's Theorem*, Van Nostrand, Princeton-Toronto-London, 1966.
- [17] Hervé R.M., *Recherches axiomatiques sur la théorie des fonctions surharmoniques et du potentiel*, Ann. Inst. Fourier **12** (1962), 415–571.
- [18] Hervé R.M., Hervé M., *Les fonctions surharmoniques associées à un opérateur elliptique du second ordre à coefficients discontinus*, Ann. Inst. Fourier **19** (1969), no. 1, 305–359.
- [19] Meyer M., *Balayage spaces on topological sums*, Potential Theory (Prague, 1987), Plenum, New York, 1988, pp. 237–246.
- [20] Nakai M., *Martin boundary over an isolated singularity of rotation free density*, J. Math. Soc. Japan **26** (1974), 483–507.

- [21] Smyrnelis E.P., *Axiomatique des fonctions biharmoniques, I*, Ann. Inst. Fourier **26** (1976), no. 1, 35–97.
- [22] Smyrnelis E.P., *Axiomatique des fonctions biharmoniques, II*, Ann. Inst. Fourier **26** (1976), no. 3, 1–47.

UNIVERSITÉ IBN TOFAIL, FACULTÉ DE SCIENCES, B.P. 133, KÉNITRA, MOROCCO

(Received April 13, 2004, revised April 7, 2006)