

The fixed point property in a Banach space isomorphic to c_0

COSTAS POULIOS

Abstract. We consider a Banach space, which comes naturally from c_0 and it appears in the literature, and we prove that this space has the fixed point property for non-expansive mappings defined on weakly compact, convex sets.

Keywords: non-expansive mappings; fixed point property; Banach spaces isomorphic to c_0

Classification: Primary 47H10, 47H09, 46B25

1. Introduction

Let K be a weakly compact, convex subset of a Banach space X . A mapping $T : K \rightarrow K$ is called *non-expansive* if $\|Tx - Ty\| \leq \|x - y\|$ for any $x, y \in K$. In the case where every non-expansive map $T : K \rightarrow K$ has a fixed point, we say that K has the *fixed point property*. The space X is said to have the fixed point property if every weakly compact, convex subset of X has the fixed point property.

A lot of Banach spaces are known to enjoy the aforementioned property. The earlier results show that uniformly convex spaces have the fixed point property (see [3]) and this is also true for the wider class of spaces with normal structure (see [7]). The classical Banach spaces ℓ_p , L_p with $1 < p < \infty$ are uniformly convex and hence they have the fixed point property. On the contrary, the space L_1 fails this property (see [1]).

The proofs of many positive results depend on the notion of minimal invariant sets. Suppose that K is a weakly compact, convex set, $T : K \rightarrow K$ is a non-expansive mapping and C is a nonempty, weakly compact, convex subset of K such that $T(C) \subseteq C$. The set C is called *minimal* for T if there is no strictly smaller weakly compact, convex subset of C which is invariant under T . A straightforward application of Zorn's lemma implies that K always contains minimal invariant subsets. So, a standard approach in proving fixed point theorems is to first assume that K itself is minimal for T and then use the geometrical properties of the space to show that K must be a singleton. Therefore, T has a fixed point.

Although a non-expansive map $T : K \rightarrow K$ does not have to have fixed points, it is well-known that T always has an *approximate fixed point sequence*. This means that there is a sequence (x_n) in K such that $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$. For such sequences, the following result holds (see [6]).

Theorem 1.1. *Let K be a weakly compact, convex set in a Banach space, let $T : K \rightarrow K$ be a non-expansive map such that K is T -minimal, and let (x_n) be any approximate fixed point sequence. Then, for all $x \in K$,*

$$\lim_{n \rightarrow \infty} \|x - x_n\| = \text{diam}(K).$$

Although from the beginning of the theory it became clear that the classical spaces $\ell_p, L_p, 1 < p < \infty$ have the fixed point property, the case of c_0 remained unsolved for some period of time. The geometrical properties of this space are not very nice, in the sense that c_0 does not possess normal structure. However, it was finally proved that the geometry of c_0 is still good enough and it does not allow the existence of minimal sets with positive diameter, that is, c_0 has the fixed point property. This was done by B. Maurey [8] (see also [4]) who also proved that every reflexive subspace of L_1 has the fixed point property.

Theorem 1.2. *The space c_0 has the fixed point property.*

The proof of Theorem 1.2 is based on the fact that the set of approximate fixed point sequences is convex in a natural sense. More precisely, we have the following ([8], [4]).

Theorem 1.3. *Let K be a weakly compact, convex subset of a Banach space which is minimal for a non-expansive map $T : K \rightarrow K$. Let (x_n) and (y_n) be approximate fixed point sequences for T such that $\lim_{n \rightarrow \infty} \|x_n - y_n\|$ exists. Then there is an approximate fixed point sequence (z_n) in K such that*

$$\lim_{n \rightarrow \infty} \|x_n - z_n\| = \lim_{n \rightarrow \infty} \|y_n - z_n\| = \frac{1}{2} \lim_{n \rightarrow \infty} \|x_n - y_n\|.$$

In the present paper, we define a Banach space X isomorphic to c_0 and we prove that this space has the fixed point property. Our interest in this space derives from several reasons. Firstly, the space X comes from c_0 in a natural way. In fact, the Schauder basis of X is equivalent to the summing basis of c_0 . Secondly, the space X is close to c_0 in the sense that the Banach-Mazur distance between the two spaces is equal to 2. It is worth mentioning that from the proof of Theorem 1.2 we can conclude that whenever Y is a Banach space isomorphic to c_0 and the Banach-Mazur distance between Y and c_0 is strictly less than 2, then Y has the fixed point property. In our case, the Banach-Mazur distance is equal to 2, that is the space X lies on the boundary of what is already known. This fact should also be compared with the following question in metric fixed point theory: Find a nontrivial class of Banach spaces invariant under isomorphism such that each member of the class has the fixed point property (a trivial example is the class of spaces isomorphic to ℓ_1). We shall see that even for spaces close to c_0 , such as the space X , the situation is quite complicated and this points out the difficulty of the aforementioned question. Finally, the space X has been used in several places in the study of the geometry of Banach spaces (for instance see [5], [2]). More precisely, the well-known Hagler Tree space (HT) [5] contains

a plethora of subspaces isometric to X . Nevertheless, we do not know if HT has the fixed point property.

2. Definition and basic properties

We consider the vector space c_{00} of all real-valued finitely supported sequences. We let $(e_n)_{n \in \mathbb{N}}$ stand for the usual unit vector basis of c_{00} , that is $e_n(i) = 1$ if $i = n$ and $e_n(i) = 0$ if $i \neq n$. If $S \subset \mathbb{N}$ is any interval of integers and $x = (x_i) \in c_{00}$ then we set $S^*(x) = \sum_{i \in S} x_i$. We now define the norm of x as follows

$$\|x\| = \sup |S^*(x)|$$

where the supremum is taken over all finite intervals $S \subset \mathbb{N}$. The space X is the completion of the normed space we have just defined.

It is easily verified that the sequence (e_n) is a normalized monotone Schauder basis for the space X . In the following, $(e_n^*)_{n \in \mathbb{N}}$ denotes the sequence of the biorthogonal functionals and $(P_n)_{n \in \mathbb{N}}$ denotes the sequence of the natural projections associated to the basis (e_n) . That is, for any $x = \sum_{i=1}^{\infty} x_i e_i \in X$ we have $e_n^*(x) = x_n$ and $P_n(x) = \sum_{i=1}^n x_i e_i$.

Furthermore, if $S \subset \mathbb{N}$ is any interval of integers (not necessarily finite), we define the functional $S^* : X \rightarrow \mathbb{R}$ by $S^*(x) = S^*(\sum_{i=1}^{\infty} x_i e_i) = \sum_{i \in S} x_i$. It is easy to see that S^* is a bounded linear functional with $\|S^*\| = 1$. In the special case where $S = \mathbb{N}$, the corresponding functional is denoted by B^* (instead of the confusing \mathbb{N}^*). Therefore, $B^*(x) = \sum_{i=1}^{\infty} x_i$ for any $x = \sum_{i=1}^{\infty} x_i e_i \in X$.

The following proposition provides some useful properties of the space X and demonstrates the relation between X and c_0 . We remind that for any pair E, F of isomorphic normed spaces, the Banach-Mazur distance between E and F is defined as follows

$$d(E, F) = \inf \{ \|T\| \cdot \|T^{-1}\| \mid T : E \rightarrow F \text{ is an isomorphism from } E \text{ onto } F \}.$$

Proposition 2.1. *The following holds.*

- (1) *The space X is isomorphic to c_0 and in particular the basis of X is equivalent to the summing basis of c_0 .*
- (2) *The subspace of X^* generated by the sequence of the biorthogonal functionals has codimension 1. More precisely, $X^* = \overline{\text{span}}\{e_n^*\}_{n \in \mathbb{N}} \oplus \langle B^* \rangle$.*
- (3) *The Banach-Mazur distance $d(X, c_0)$ between X and c_0 is equal to 2.*

PROOF: We define the linear operator

$$\begin{aligned} \Phi : X &\rightarrow c_0 \\ x = (x_i) &\mapsto \left(\sum_{i=1}^{\infty} x_i, \sum_{i=2}^{\infty} x_i, \dots \right). \end{aligned}$$

It is easily verified that Φ is an isomorphism from X onto c_0 with $\|\Phi\| = 1$, $\|\Phi^{-1}\| = 2$ and Φ maps the basis of X to the summing basis of c_0 . This proves

the first assertion. The second assertion is an immediate consequence of the relation between X and c_0 established above.

It remains to show that the Banach-Mazur distance $d = d(X, c_0)$ is equal to 2. Firstly, we observe that the isomorphism Φ defined above implies that $d \leq 2$. In order to prove the reverse inequality we fix a real number $\epsilon > 0$. Then there exists an isomorphism $T : X \rightarrow c_0$ from X onto c_0 such that $\|x\| \leq \|Tx\|_{c_0} \leq (d + \epsilon)\|x\|$ for any $x \in X$. We now consider the normalized sequence (x_n) in X where $x_n = (x_n(i))_{i \in \mathbb{N}}$ is defined by

$$x_n(2n - 1) = -1, \quad x_n(2n) = 1, \quad x_n(i) = 0 \text{ otherwise.}$$

The description of X^* given by the second assertion implies that any bounded sequence $(t_n)_{n \in \mathbb{N}}$ of elements of X converges weakly to 0 if and only if $e_m^*(t_n) \rightarrow 0$ for every $m \in \mathbb{N}$ and $B^*(t_n) \rightarrow 0$. It follows that the sequence $(x_n)_{n \in \mathbb{N}}$ defined above is weakly null. Now we set $y_n = T(x_n)$ for any $n \in \mathbb{N}$ and we have $1 \leq \|y_n\|_{c_0} \leq d + \epsilon$ and $(y_n)_{n \in \mathbb{N}}$ converges weakly to 0. Therefore, we find $k_1 \in \mathbb{N}$ such that the vectors y_1 and y_{k_1} have essentially disjoint supports. More precisely, since $y_1 \in c_0$, there exists $N_1 \in \mathbb{N}$ such that $|y_1(i)| < \epsilon$ for any $i > N_1$. Since $y_n \rightarrow 0$ weakly, we find k_1 so that $|y_{k_1}(i)| < \epsilon$ for any $i \leq N_1$. It follows that $\|y_1 - y_{k_1}\|_{c_0} \leq \max\{\|y_1\|_{c_0}, \|y_{k_1}\|_{c_0}\} + \epsilon \leq d + 2\epsilon$. On the other hand, $\|x_1 - x_{k_1}\| = 2$. Therefore,

$$2 = \|x_1 - x_{k_1}\| \leq \|y_1 - y_{k_1}\|_{c_0} \leq d + 2\epsilon.$$

If ϵ tends to 0, we obtain $2 \leq d$ as we desire. □

3. The fixed point property

This section is entirely devoted to the proof of the fixed point property for the space X . First we need to establish some notation. If $S, S' \subset \mathbb{N}$ are intervals we write $S < S'$ to mean that $\max S < \min S'$. Moreover, if $k \in \mathbb{N}$, we write $k < S$ (resp., $S < k$) to mean $k < \min S$ (resp., $\max S < k$). Finally, for any $x = (x_i) \in X$, $\text{supp}(x) = \{i \in \mathbb{N} \mid x_i \neq 0\}$ denotes the support of x .

Theorem 3.1. *The space X has the fixed point property.*

PROOF: We follow the standard approach. We assume that K is a weakly compact, convex subset of X which is minimal for a non-expansive map $T : K \rightarrow K$. Using the geometry of the space X , we have to show that K is a singleton, that is $\text{diam}(K) = 0$. Let us suppose that $\text{diam}(K) > 0$ and now we have to reach a contradiction. Without loss of generality we may assume that $\text{diam}(K) = 1$.

Let $(x_n)_{n \in \mathbb{N}}$ be an approximate fixed point sequence for the map T in the set K . By passing to a subsequence and then using some translation, we may assume that $0 \in K$ and (x_n) converges weakly to 0. Theorem 1.1 implies that $\lim_n \|x_n\| = \text{diam}(K) = 1$.

Furthermore, using a standard perturbation argument we may assume that (x_n) is a finitely supported approximate fixed point sequence. Indeed, we inductively construct a subsequence (x_{q_n}) of (x_n) and integers $l_0 = 0 < l_1 < l_2 < \dots$ such that for every $n \in \mathbb{N}$, $\|P_{l_{n-1}}(x_{q_n})\| < 1/n$ and $\|x_{q_n} - P_{l_n}(x_{q_n})\| < 1/n$. We start with $x_{q_1} = x_1$ and $l_0 = 0$. Suppose that $q_1 < q_2 < \dots < q_n$ and $l_0 < l_1 < \dots < l_{n-1}$ have been defined. Then there exists $l_n > l_{n-1}$ such that $\|x_{q_n} - P_{l_n}(x_{q_n})\| < 1/n$. Since (x_n) is weakly null, it follows that $P_m(x_n) \rightarrow 0$ for every $m \in \mathbb{N}$. Therefore, there exists $q_{n+1} > q_n$ such that $\|P_{l_n}(x_{q_{n+1}})\| < \frac{1}{n+1}$. The construction of (x_{q_n}) and (l_n) is complete. Consequently, by passing to the subsequence (x_{q_n}) and perturbing (x_{q_n}) , if necessary, we may assume that for the original sequence (x_n) we have $\text{supp}(x_n) \subset (l_{n-1}, l_n]$ for every $n \in \mathbb{N}$, that is, (x_n) consists of finitely supported vectors.

We next consider the subsequences $(z_n) = (x_{2n-1})$ and $(y_n) = (x_{2n})$ and we also set $l_{2n-1} = k_n, l_{2n} = m_n$ for every $n \in \mathbb{N}$ and $m_0 = l_0$. The properties of the sequence (x_n) imply that the following holds.

- (1) (z_n) and (y_n) are approximate fixed point sequences for the map T and $\lim \|z_n\| = \lim \|y_n\| = 1$.
- (2) (z_n) and (y_n) converge weakly to 0.
- (3) $\text{supp}(z_n) \subset (m_{n-1}, k_n]$ and $\text{supp}(y_n) \subset (k_n, m_n]$ for every $n \in \mathbb{N}$.
- (4) $\lim \|z_n - y_n\| = 1$.

In order to justify the fourth conclusion, we first observe that $\limsup \|z_n - y_n\| \leq \text{diam}(K) = 1$. On the other hand, by the definition of the norm of the space X , for every $n \in \mathbb{N}$ there exists a finite interval $E_n \subset \mathbb{N}$ such that $\|z_n\| = |E_n^*(z_n)|$. Clearly we may assume that $E_n \subset (m_{n-1}, k_n]$. Then $\|z_n - y_n\| \geq |E_n^*(z_n - y_n)| = \|z_n\|$. Since $\lim \|z_n\| = 1$, it emerges that $\liminf \|z_n - y_n\| \geq 1$ and finally $\lim \|z_n - y_n\| = 1$.

We are ready now to apply Maurey's theorem (Theorem 1.3). To this end, we fix a positive integer $N \geq 4$, we set $\epsilon = 2^{-N}$ and we iteratively use Theorem 1.3 as follows. Firstly, we consider the sequences (z_n) and (y_n) . Applying Theorem 1.3 we obtain an approximate fixed point sequence $(v_n^1)_{n \in \mathbb{N}}$ in the set K such that $\lim \|v_n^1 - y_n\| = \frac{1}{2} \lim \|z_n - y_n\| = \frac{1}{2}$ and $\lim \|v_n^1 - z_n\| = \frac{1}{2} \lim \|z_n - y_n\| = \frac{1}{2}$. Assume now that in the i -th step of this procedure we find an approximate fixed point sequence $(v_n^i)_{n \in \mathbb{N}}$ satisfying $\lim \|v_n^i - z_n\| = 2^{-i}$ and $\lim \|v_n^i - y_n\| = 1 - 2^{-i}$. Then, Theorem 1.3 implies that "halfway" between (z_n) and (v_n^i) there exists an approximate fixed point sequence $(v_n^{i+1})_{n \in \mathbb{N}}$, that is, $\lim \|v_n^{i+1} - v_n^i\| = \frac{1}{2} \lim \|v_n^i - z_n\| = 2^{-(i+1)}$ and $\lim \|v_n^{i+1} - z_n\| = \frac{1}{2} \lim \|v_n^i - z_n\| = 2^{-(i+1)}$. Now, we estimate the distance between v_n^{i+1} and y_n . We have

$$\begin{aligned} \|v_n^{i+1} - y_n\| &\leq \|v_n^{i+1} - v_n^i\| + \|v_n^i - y_n\| \text{ and} \\ \|v_n^{i+1} - y_n\| &\geq \|z_n - y_n\| - \|v_n^{i+1} - z_n\|. \end{aligned}$$

Therefore, it follows that $\lim \|v_n^{i+1} - y_n\| = 1 - 2^{-(i+1)}$. After N iterated applications of Theorem 1.3 we find a sequence $(v_n)_{n \in \mathbb{N}} = (v_n^N)_{n \in \mathbb{N}}$ in the set K satisfying

the following: (v_n) is an approximate fixed point sequence for the map T (which implies that $\lim \|v_n\| = 1$) and further $\lim \|v_n - z_n\| = \epsilon$ and $\lim \|v_n - y_n\| = 1 - \epsilon$. Therefore, for all sufficiently large $n \in \mathbb{N}$ the following holds:

- (1) $\|v_n\| > 1 - \frac{\epsilon}{2}$;
- (2) $\|v_n - z_n\| < 3\epsilon/2$ and $\|v_n - y_n\| < 1 - \frac{\epsilon}{2}$;
- (3) $|B^*(z_n)| < \epsilon/2$ (since (z_n) is weakly null).

We also set $S_n = (m_{n-1}, k_n]$ so that we have $S_1 < S_2 < \dots$. Concerning the sequence (v_n) in the set K and the sequence of intervals (S_n) we prove the following two claims.

Claim 1. For all sufficiently large n , the support of v_n is essentially contained in the interval S_n , in the sense that if S is any interval with $S \cap S_n = \emptyset$ then $|S^*(v_n)| < 3\epsilon/2$.

Indeed, we know that $\text{supp}(z_n) \subset (m_{n-1}, k_n] = S_n$. Therefore, if S is any interval with $S \cap S_n = \emptyset$ then $S^*(z_n) = 0$ and hence

$$|S^*(v_n)| = |S^*(v_n - z_n)| \leq \|v_n - z_n\| < \frac{3\epsilon}{2}.$$

Claim 2. For all sufficiently large n , there exist intervals $L_n < R_n$ such that $S_n = L_n \cup R_n$ and $L_n^*(v_n) < -1 + 7\epsilon$, $R_n^*(v_n) > 1 - 2\epsilon$.

We fix a sufficiently large positive integer n . Since $\|v_n\| > 1 - \frac{\epsilon}{2}$, it follows that there exists a finite interval $F_n \subset \mathbb{N}$ such that $|F_n^*(v_n)| > 1 - \frac{\epsilon}{2}$. If $k_n < F_n$, we know by the previous claim that $|F_n^*(v_n)| < 3\epsilon/2$, which is a contradiction. Moreover, if we assume that $F_n \leq k_n$ then $F_n \cap (k_n, m_n] = \emptyset$ and the choice of (y_n) implies $F_n^*(y_n) = 0$. Thus,

$$|F_n^*(v_n)| = |F_n^*(v_n - y_n)| \leq \|v_n - y_n\| < 1 - \frac{\epsilon}{2},$$

which is also a contradiction. By this discussion it is clear that $\min F_n \leq k_n < \max F_n$. Now we set $R_n = F_n \cap [1, k_n]$ and we estimate

$$1 - \frac{\epsilon}{2} < |F_n^*(v_n)| \leq |R_n^*(v_n)| + |(F_n \setminus R_n)^*(v_n)| < |R_n^*(v_n)| + \frac{3\epsilon}{2},$$

where the last inequality follows from Claim 1. Therefore, $|R_n^*(v_n)| > 1 - 2\epsilon$. Passing to a subsequence, we may assume that either $R_n^*(v_n) > 1 - 2\epsilon$ for all sufficiently large n or $R_n^*(v_n) < -1 + 2\epsilon$ for all sufficiently large n . We suppose that the first possibility happens, as the second one is treated similarly (by interchanging the roles of L_n and R_n). Consequently, for the interval R_n we have $\max R_n = k_n$ and $R_n^*(v_n) > 1 - 2\epsilon$.

On the other hand, we observe that

$$|B^*(v_n)| \leq |B^*(v_n - z_n)| + |B^*(z_n)| \leq \|v_n - z_n\| + \frac{\epsilon}{2} < 2\epsilon.$$

We note that the sequence (v_n) is not necessarily weakly null. However, v_n is close to z_n and hence $|B^*(v_n)|$ is very small. We next set $G_n = [1, \min R_n)$ (possibly empty) and $W_n = (k_n, +\infty)$. Then,

$$\begin{aligned} 2\epsilon > |B^*(v_n)| &= |G_n^*(v_n) + R_n^*(v_n) + W_n^*(v_n)| \\ &\geq R_n^*(v_n) - |G_n^*(v_n)| - |W_n^*(v_n)| \\ &> 1 - 2\epsilon - |G_n^*(v_n)| - \frac{3\epsilon}{2}. \end{aligned}$$

Therefore G_n is non-empty and $|G_n^*(v_n)| > 1 - \frac{11\epsilon}{2}$. However, if $G_n^*(v_n) > 1 - \frac{11\epsilon}{2}$, then it would follow

$$|B^*(v_n)| \geq R_n^*(v_n) + G_n^*(v_n) - |W_n^*(v_n)| \geq 2 - 9\epsilon,$$

which is a contradiction. Hence, $G_n^*(v_n) < -1 + \frac{11\epsilon}{2}$. Further, we observe that we cannot have $G_n < S_n$, since in this case it would follow $|G_n^*(v_n)| < \frac{3\epsilon}{2}$. Consequently, $\max G_n > m_{n-1}$ which clearly implies $\min R_n > m_{n-1} + 1$. Finally, we set $L_n = G_n \cap (m_{n-1}, k_n]$ and we estimate

$$-1 + \frac{11\epsilon}{2} > G_n^*(v_n) = L_n^*(v_n) + (G_n \setminus L_n)^*(v_n) \geq L_n^*(v_n) - \frac{3\epsilon}{2}.$$

We deduce that $L_n^*(v_n) < -1 + 7\epsilon$. Therefore, the intervals $L_n < R_n$ satisfy the following: $S_n = L_n \cup R_n$, $R_n^*(v_n) > 1 - 2\epsilon$ and $L_n^*(v_n) < -1 + 7\epsilon$. The proof of the claim is now complete.

Using the construction and the properties of the sequences (v_n) and (S_n) , we can reach the final contradiction and finish the proof of the theorem. Our goal is to show that for all sufficiently large $n \in \mathbb{N}$, $\|v_n - v_{n+1}\| \geq 5/4 > 1$, contradicting the assumption $\text{diam}(K) = 1$. Indeed, we fix a sufficiently large $n \in \mathbb{N}$ and we consider the intervals $D = (k_n, m_n]$ and $S = R_n \cup D \cup L_{n+1}$. Then, using Claim 1 and Claim 2 we have

$$\begin{aligned} S^*(v_n) &= R_n^*(v_n) + (D \cup L_{n+1})^*(v_n) > 1 - 2\epsilon - \frac{3\epsilon}{2} = 1 - \frac{7\epsilon}{2} \\ S^*(v_{n+1}) &= (R_n \cup D)^*(v_{n+1}) + L_{n+1}^*(v_{n+1}) < \frac{3\epsilon}{2} - 1 + 7\epsilon = -1 + \frac{17\epsilon}{2}. \end{aligned}$$

Therefore,

$$\|v_n - v_{n+1}\| \geq |S^*(v_n - v_{n+1})| = |S^*(v_n) - S^*(v_{n+1})| \geq 2 - 12\epsilon.$$

The choice of ϵ implies that $\|v_n - v_{n+1}\| \geq 5/4 > 1$ for all sufficiently large $n \in \mathbb{N}$, hence we obtain the desired contradiction. \square

Acknowledgments. The author would like to thank the referee for his/her valuable remarks.

REFERENCES

- [1] Alspach D., *A fixed point free nonexpansive map*, Proc. Amer. Math. Soc. **82** (1981), 423–424.
- [2] Argyros S.A., Deliyanni I., Toliás A.G., *Hereditarily indecomposable Banach Algebras of diagonal operators*, Israel J. Math. **181** (2011), 65–110.
- [3] Browder F.E., *Nonexpansive nonlinear operators in Banach spaces*, Proc. Nat. Acad. Sci. U.S.A. **54** (1965), 1041–1044.
- [4] Elton J., Lin P., Odell E., Szarek S., *Remarks on the fixed point problem for nonexpansive maps*, Contemporary Math. **18** (1983), 87–119.
- [5] Hagler J., *A counterexample to several questions about Banach spaces*, Studia Math. **60** (1977), 289–308.
- [6] Karlovitz L.A., *Existence of fixed points for nonexpansive mappings in a space without normal structure*, Pacific J. Math. **66** (1976), 153–159.
- [7] Kirk W.A., *A fixed point theorem for mappings which do not increase distances*, Amer. Math. Monthly **72** (1965), 1004–1006.
- [8] Maurey B., *Points fixes des contractions sur un convexe forme de L^1* , Seminaire d'Analyse Fonctionnelle 80–81, Ecole Polytechnique, Palaiseau, 1981.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ATHENS, 15784 ATHENS, GREECE

E-mail: k-poullos@math.uoa.gr

(Received March 4, 2013, revised December 4, 2013)