

A continuum X such that $C(X)$ is not continuously homogeneous

ALEJANDRO ILLANES

Abstract. A metric continuum X is said to be continuously homogeneous provided that for every two points $p, q \in X$ there exists a continuous surjective function $f : X \rightarrow X$ such that $f(p) = q$. Answering a question by W.J. Charatonik and Z. Garncarek, in this paper we show a continuum X such that the hyperspace of subcontinua of X , $C(X)$, is not continuously homogeneous.

Keywords: continuum; continuously homogeneous; hyperspace

Classification: Primary 54B20; Secondary 54F15

1. Introduction

A *compactum* is a compact metric space with more than one point. A *continuum* is a connected compactum. A *mapping* is a continuous function. A continuum X is said to be *continuously homogeneous* if for every two points $p, q \in X$, there exists a surjective mapping $f : X \rightarrow X$ such that $f(p) = q$.

For the continuum X , we consider its hyperspaces:

$$2^X = \{A \subset X : A \text{ is closed and nonempty}\}, \text{ and}$$

$$C(X) = \{A \in 2^X : A \text{ is connected}\}.$$

The hyperspace 2^X is endowed with the Hausdorff metric H [5, Definition 2.1].

In [2] W.J. Charatonik and Z. Garncarek studied conditions for a hyperspace being continuously homogeneous. They showed that the hyperspace 2^X is continuously homogeneous for an arbitrary continuum X , and the hyperspace $C(X)$ is such if either X is locally connected or X contains an open subset with uncountably many components. They also asked [2, Question 2] the question: Is the hyperspace $C(X)$ continuously homogeneous for every continuum X ?

In this paper we answer the question by Charatonik and Garncarek in the negative by showing a continuum Z such $C(Z)$ is not continuously homogeneous.

DOI 10.14712/1213-7243.2015.146

This paper was partially supported by the project “Hiperespacios topológicos (0128584)” of Consejo Nacional de Ciencia y Tecnología (CONACYT), 2009; and the project “Teoría de Continuos, Hiperespacios y Sistemas Dinámicos” (IN104613) of PAPIIT, DGAPA, UNAM.

2. The example

Given a connected subset S of a continuum X we denote by $C(S)$ the set of subcontinua of X that are contained in S . A compactum X is *connected im kleinen* at a point $p \in X$ provided that for each open subset U of X such that $p \in U$, there exists a subcontinuum M of X such that p belongs to the interior of M and $M \subset U$. The set of points p in X such that X is not connected im kleinen at p is denoted by $N(X)$. We will use the following well known lemma [4, p. 28].

Lemma 1 ([4, p. 28]). *Let $f : X \rightarrow Y$ be a surjective mapping between compacta. Then $N(Y) \subset f(N(X))$.*

The following lemma can be easily proven by using Theorems 2 and 3 of [4].

Lemma 2. *Let X be a compactification of the ray $[0, \infty)$ with remainder R and $S = X - R$. Then:*

- (a) $N(C(X)) \subset C(R)$,
- (b) if $C(R) \subset \text{cl}_{C(X)}(C(S))$, then $N(C(X)) = C(R) - \{R\}$.

In order to construct the continuum Z , we use ideas in the paper [1]. We construct a sequence of continua Z_1, Z_2, \dots in the Hilbert cube $Q = [-1, 1]^\infty$ in such a way that the complexity of the set of points of non connectedness im kleinen of Z_{n+1} is bigger than the one of Z_n . In this way, we obtain that there is not a mapping from $C(Z_n)$ onto $C(Z_{n+1})$. Each Z_{n+1} is a compactification of the ray $[0, \infty)$ with remainder Z_n and Z_n is a retract of Z_{n+1} . The continua Z_n can also be defined in the Euclidean plane but the description is more complicated.

We start by defining $Z_1 = \{(0, t, 0, 0, \dots) \in Q : t \in [-1, 1]\}$ and $Z_2 = Z_1 \cup S_1$, where $S_1 = \{(t, \sin(\frac{1}{t}), 0, 0, \dots) \in Q : t \in (0, 1]\}$.

Define $g_1 : [0, \infty) \rightarrow [0, 1]^2$ by $g_1(t) = (\frac{1}{1+t}, \sin(1+t))$.

Inductively, suppose that $n \geq 2$, $Z_n \subset [-1, 1]^n \times \{0\} \times \{0\} \times \dots$ has been constructed, $Z_n = Z_{n-1} \cup S_{n-1}$, $Z_{n-1} \cap S_{n-1} = \emptyset$, Z_n is a compactification of $[0, \infty)$ with remainder Z_{n-1} , $h_{n-1} : [0, \infty) \rightarrow S_{n-1}$ is a homeomorphism, $h_{n-1} = g_{n-1} \times \{0\} \times \{0\} \times \dots$, where $g_{n-1} : [0, \infty) \rightarrow [-1, 1]^n$ is an embedding. Define $k : [0, \infty) \rightarrow [0, \infty)$ by

$$k(t) = \begin{cases} (2m+1)(t-2m), & \text{if } t \in [2m, 2m+1] \text{ for some } m \geq 0, \\ (2m+1)(2+2m-t) & \text{if } t \in [2m+1, 2m+2] \text{ for some } m \geq 0. \end{cases}$$

Define $g_n : [0, \infty) \rightarrow [-1, 1]^{n+1}$ by $g_n(t) = (g_{n-1}(k(t)), \frac{1}{t+1})$ and define $h_n : [0, \infty) \rightarrow Q$ by

$$h_n = g_n \times \{0\} \times \{0\} \times \dots$$

Clearly, h_n is an embedding. Let $S_n = \text{Im } h_n$ and $Z_{n+1} = Z_n \cup S_n$. Then Z_{n+1} is a compactification of $[0, \infty)$ with remainder Z_n and the natural projection $r_n : Z_{n+1} \rightarrow Z_n$, defined on S_n by $(g_{n-1}(k(t)), \frac{1}{t+1}, 0, \dots) \mapsto (g_{n-1}(k(t)), 0, \dots)$, is a retraction.

Claim 1. For each $n \in \mathbb{N}$, $C(Z_n) \subset \text{cl}_{C(Z_{n+1})}(C(S_n))$.

We prove Claim 1. Clearly, $C(Z_1) \subset \text{cl}_{C(Z_2)}(C(S_1))$. Take $n \geq 2$ and take a subarc J of S_{n-1} , let L be a subinterval of $[0, \infty)$ such that $h_{n-1}(L) = J$. Let $M \in \mathbb{N}$ be such that $L \subset [0, 2M + 1]$. Then for each $m > M$, there exists a subinterval J_m of $[2m, 2m + 1]$ such that $k(J_m) = L$. Thus, the sequence $\{h_n(J_m)\}_{m=M+1}^\infty$ is a sequence in S_n such that $\lim h_n(J_m) = J$. We have shown that $C(S_{n-1}) \subset \text{cl}_{C(Z_{n+1})}(C(S_n))$ for every $n \geq 2$.

Notice that $C(Z_2) \subset \text{cl}_{C(Z_2)}(C(S_1)) \subset \text{cl}_{C(Z_3)}(C(S_2))$, so

$$C(Z_2) \subset \text{cl}_{C(Z_3)}(C(S_2)).$$

Since $Z_3 = Z_2 \cup S_2$ is a compactification of the ray $[0, \infty)$ with remainder Z_2 , $C(Z_3) = C(Z_2) \cup C(S_2) \cup \{A \in C(Z_3) : Z_2 \subset A\}$. It is easy to prove that $C(S_2) \cup \{A \in C(Z_3) : Z_2 \subset A\} \subset \text{cl}_{C(Z_3)}(C(S_2))$. Since $C(Z_2) \subset \text{cl}_{C(Z_3)}(C(S_2))$, we conclude that $C(Z_3) \subset \text{cl}_{C(Z_3)}(C(S_2))$. By the fact we prove two paragraphs above, $\text{cl}_{C(Z_3)}(C(S_2)) \subset \text{cl}_{C(Z_4)}(C(S_3))$. Hence, $C(Z_3) \subset \text{cl}_{C(Z_4)}(C(S_3))$.

With a similar procedure as the one in the previous paragraph, Claim 1 can be proved for each $n \geq 4$.

The following claim is a consequence of Claim 1 and Lemma 2.

Claim 2. For each $n \in \mathbb{N}$, $N(C(Z_{n+1})) = C(Z_n) - \{Z_n\}$.

Given a subset A of Q and $n \in \mathbb{N}$ let $A(n) = \{(\frac{1}{2^n} + \frac{a}{8^n}, 0) \in Q \times [-1, 1] : a \in A\}$.

Now, we construct the continuum Z as a subspace of the space $Q \times [-1, 1]$. For each $n \geq 2$, let $X_n = Z_n(n)$. Then X_n is a compactification of the ray $[0, \infty)$ with remainder $Z_{n-1}(n)$ and $X_n - Z_{n-1}(n) = S_{n-1}(n)$. Let p_n be the end point of the ray $S_{n-1}(n)$. Choose an arc $L_n \subset Q \times [0, 1]$ with end points p_n and p_{n+1} such that $L_n - \{p_n, p_{n+1}\} \subset Q \times (0, 1]$, $\lim L_n = \{\theta\}$, where $\theta = (0, 0, \dots)$, and $L_n \cap (\bigcup\{L_m : m \geq 2 \text{ and } m \neq n\}) \subset \{p_n, p_{n+1}\}$. Let $\sigma_n : Z_n \rightarrow X_n$ be the homeomorphism given by $\sigma_n(z) = (\frac{1}{2^n} + \frac{z}{8^n}, 0)$.

Finally, define $Z = \{\theta\} \cup (\bigcup\{X_n : n \geq 2\}) \cup (\bigcup\{L_n : n \geq 2\})$. Then Z is a continuum.

The following claim follows from Theorems 2 and 3 of [4].

Claim 3. $N(C(Z)) = \bigcup\{N(C(X_n)) : n \geq 2\}$.

We are going to show that $C(Z)$ is not continuously homogeneous. Suppose the contrary. Then there exists a continuous surjective mapping $f : C(Z) \rightarrow C(Z)$ such that $f(\{\theta\}) = \{p_2\}$. Let $\mathcal{U} = C(L_2 \cup X_2)$. Since $L_2 \cup X_2$ contains p_2 in its interior, \mathcal{U} is a neighborhood of $\{p_2\}$ in $C(Z)$. Since $\lim(X_n \cup L_n) = \{\theta\}$, there exists $M \geq 3$ such that

$$f(C(\{\theta\} \cup (\bigcup\{X_n : n > M\}) \cup (\bigcup\{L_n : n > M\}))) \subset \mathcal{U}.$$

Let $Y = (\bigcup\{X_n : n \in \{2, \dots, M\}\}) \cup (\bigcup\{L_n : n \in \{2, \dots, M\}\})$. Notice that $N(C(Y)) = \bigcup\{N(C(X_n)) : n \in \{2, \dots, M\}\}$.

By Claim 3 and Lemma 1,

$$N(C(X_{M+1})) \subset N(C(Z)) \subset f(N(C(Z))) = \bigcup \{f(N(C(X_n))) : n \geq 2\}.$$

Since $\bigcup \{f(C(X_n)) : n > M\} \subset C(L_2 \cup X_2)$ and $C(L_2 \cup X_2) \cap C(X_{M+1}) = \emptyset$, we have that

$$N(C(X_{M+1})) \subset \bigcup \{f(N(C(X_n))) : n \in \{2, \dots, M\}\}.$$

Since $\{p_{M+1}\} = \text{Bd}_Z(X_{M+1})$, we have that the function $t : Z \rightarrow X_{M+1}$ defined by

$$t(z) = \begin{cases} z, & \text{if } z \in X_{M+1}, \\ p_{M+1}, & \text{if } z \notin X_{M+1}, \end{cases}$$

is a retraction. Then the mapping $T : C(Z) \rightarrow C(X_{M+1})$ given by $T(A) = t(A)$ (the image of A under t) is a retraction. Then the inclusion $N(C(X_{M+1})) \subset \bigcup \{f(N(C(X_n))) : n \in \{2, \dots, M\}\}$ implies that

$$N(C(X_{M+1})) \subset \bigcup \{T(f(N(C(X_n)))) : n \in \{2, \dots, M\}\}.$$

Consider the homeomorphism $\sigma_{M+1}^{-1} : C(X_{M+1}) \rightarrow C(Z_{M+1})$ that sends each $A \in C(X_{M+1})$ to $\sigma_{M+1}^{-1}(A)$ (the image of A under σ_{M+1}^{-1}). Then

$$\sigma_{M+1}^{-1}(N(C(X_{M+1}))) = N(C(Z_{M+1})) = C(Z_M) - \{Z_M\}.$$

Since $r_M : Z_{M+1} \rightarrow Z_M$ is a retraction, the mapping $r : C(Z_{M+1}) \rightarrow C(Z_M)$ given by $r(A) = r_M(A)$ (the image of A under r_M) is a retraction.

Thus, the function $g = \sigma_{M+1}^{-1} \circ T \circ f : C(Z) \rightarrow C(Z_{M+1})$ is a surjective mapping such that

$$N(C(Z_{M+1})) = \sigma_{M+1}^{-1}(N(C(X_{M+1}))) \subset g(\bigcup \{N(C(X_n)) : n \in \{2, \dots, M\}\}).$$

Hence, $C(Z_M) - \{Z_M\} \subset g(\bigcup \{N(C(X_n)) : n \in \{2, \dots, M\}\})$. Thus,

$$\begin{aligned} C(Z_M) - \{Z_M\} &\subset r(g(\bigcup \{N(C(X_n)) : n \in \{2, \dots, M\}\})) \\ &= r(g(\bigcup \{\sigma_n(N(C(Z_n))) : n \in \{2, \dots, M\}\})) \\ &= r(g(\bigcup \{\sigma_n(C(Z_{n-1})) - \{Z_{n-1}\} : n \in \{2, \dots, M\}\})) \\ &\subset r(g(\bigcup \{\sigma_n(C(Z_{n-1})) : n \in \{2, \dots, M\}\})) \subset C(Z_M). \end{aligned}$$

Since, $r(g(\bigcup \{\sigma_n(C(Z_{n-1})) : n \in \{2, \dots, M\}\}))$ is compact, we obtain that

$$C(Z_M) = r(g(\bigcup \{\sigma_n(C(Z_{n-1})) : n \in \{2, \dots, M\}\})).$$

Consider the compactum $W = C(Z_1) \oplus \dots \oplus C(Z_{M-1})$, which is a disjoint union of the spaces $C(Z_1), \dots, C(Z_{M-1})$ with the sum topology. Since the subsets $\sigma_2(C(Z_1)), \dots, \sigma_M(C(Z_{M-1}))$ of Z are compact and pairwise disjoint, $\bigcup\{\sigma_n(C(Z_{n-1})) : n \in \{2, \dots, M\}\}$ is homeomorphic to W .

We have shown that $C(Z_M)$ is the image under a continuous function φ of the compactum W .

By Lemma 1, $N(C(Z_M)) \subset \varphi(N(W))$. By Claim 2, this implies that

$$\begin{aligned} C(Z_{M-1}) - \{Z_{M-1}\} &\subset \varphi(\emptyset \oplus (C(Z_1) - \{Z_1\}) \oplus \dots \oplus (C(Z_{M-2}) - \{Z_{M-2}\})) \\ &= \varphi((C(Z_1) - \{Z_1\}) \oplus \dots \oplus (C(Z_{M-2}) - \{Z_{M-2}\})) \\ &\subset \varphi(C(Z_1) \oplus \dots \oplus C(Z_{M-2})). \end{aligned}$$

By the compactness of $\varphi(C(Z_1) \oplus \dots \oplus C(Z_{M-2}))$, we have that

$$C(Z_{M-1}) \subset \varphi(C(Z_1) \oplus \dots \oplus C(Z_{M-2})).$$

Since the retraction $r_{M-1} : Z_M \rightarrow Z_{M-1}$ induces a retraction $R_{M-1} : C(Z_M) \rightarrow C(Z_{M-1})$, we obtain that

$$C(Z_{M-1}) = R_{M-1}(\varphi(C(Z_1) \oplus \dots \oplus C(Z_{M-2}))).$$

This shows that $C(Z_{M-1})$ is the image under a continuous function φ_1 of the compactum $C(Z_1) \oplus \dots \oplus C(Z_{M-2})$.

Repeating this argument, we conclude that $C(Z_2)$ is a continuous image of the continuum $C(Z_1)$. This is a contradiction since $C(Z_1)$ is locally connected and $C(Z_2)$ is not locally connected. Therefore, $C(Z)$ is not continuously homogeneous.

REFERENCES

- [1] Charatonik J.J., Charatonik W.J., *A degree of nonlocal connectedness*, Rocky Mountain J. Math. **31** (2001), 1205–1236.
- [2] Charatonik W.J., Garncarek Z., *Some remarks on continuously homogeneous continua*, Bull. Polish. Acad. Sci. Math. **32** (1984), 339–342.
- [3] Engelking R., Lelek A., *Cartesian products and continuous images*, Colloq. Math. **8** (1961), 27–29.
- [4] Goodykoontz J.T., Jr., *More on connectedness in kleinen and local connectedness in $C(X)$* , Proc. Amer. Math. Soc. **65** (1977), 357–364.
- [5] Illanes A., Nadler S.B., Jr., *Hyperspaces: Fundamentals and Recent Advances*, Monographs and Textbooks in Pure and Applied Mathematics, 216, Marcel Dekker, Inc., New York and Basel, 1999.

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO,
CIRCUITO EXTERIOR, CD. UNIVERSITARIA, MÉXICO, 04510, D.F.

E-mail: illanes@matem.unam.mx

(Received February 12, 2015, revised April 18, 2015)