

# The (dis)connectedness of products of Hausdorff spaces in the box topology

VITALIJ A. CHATYRKO

*Abstract.* In this paper the following two propositions are proved:

(a) If  $X_\alpha$ ,  $\alpha \in A$ , is an infinite system of connected spaces such that infinitely many of them are nondegenerated completely Hausdorff topological spaces then the box product  $\prod_{\alpha \in A} X_\alpha$  can be decomposed into continuum many disjoint nonempty open subsets, in particular, it is disconnected.

(b) If  $X_\alpha$ ,  $\alpha \in A$ , is an infinite system of Brown Hausdorff topological spaces then the box product  $\prod_{\alpha \in A} X_\alpha$  is also Brown Hausdorff, and hence, it is connected.

A space is Brown if for every pair of its open nonempty subsets there exists a point common to their closures. There are many examples of countable Brown Hausdorff spaces in literature.

*Keywords:* box topology; connectedness; completely Hausdorff space; Urysohn space; Brown space

*Classification:* 54B10, 54D05, 54D10

## 1. Introduction

All considered spaces are nondegenerated.

It is well known (cf. [15]) that the topological product  $\prod_{\alpha \in A} X_\alpha$  of an indexed family  $\{X_\alpha\}_{\alpha \in A}$  of connected spaces is connected. Let us endow the product with the box topology and denote the new space as  $\prod_{\alpha \in A} X_\alpha$ .

**Theorem 1.1** ([11]). *Let  $X_\alpha$ ,  $\alpha \in A$ , be an infinite system of connected regular  $T_1$ -spaces. Then the space  $\prod_{\alpha \in A} X_\alpha$  is disconnected.*

This result motivated

**Question 1.2** ([3]). *Let  $X_\alpha$ ,  $\alpha \in A$ , be an infinite system of connected spaces. Under what conditions on the system is the space  $\prod_{\alpha \in A} X_\alpha$  (dis)connected?*

In [3] the authors suggested two independent sufficient conditions on topological spaces  $X_\alpha$ ,  $\alpha \in A$ , which imply disconnectedness of the product  $\prod_{\alpha \in A} X_\alpha$ .

In view of Theorem 1.1 these conditions are meaningful for connected spaces with axioms lower than  $T_3$ .

Thus a condition called a *decreasing open filtration property*, in short, *DOF-property*, is valid for  $T_1$ -spaces with at least one regular non-isolated point (and not only for such spaces). Other condition, called an *increasing open filtration property*, shortly, *IOF-property*, holds for example for the Khalimsky line  $\mathbb{K}$ , see [6], a topological space  $(\mathbb{Z}, \tau)$ , where  $\mathbb{Z}$  is the set of all integers and  $\tau$  is the topology on  $\mathbb{Z}$  generated by the base  $\mathcal{B} = \{\{2k+1\}, \{2k-1, 2k, 2k+1\} : k \in \mathbb{Z}\}$ . It is easy to see that  $\mathbb{K}$  is a connected  $T_0$ -space (but not  $T_1$ ). Let us note, see [12], that the Khalimsky line  $\mathbb{K}$  and the spaces  $\mathbb{K}^n$ ,  $n \geq 2$ , are basic objects of digital topology.

In [4] the authors observed that if each space  $X_\alpha$ ,  $\alpha \in A$ , is *hyperconnected* (i.e. every pair of nonempty open subsets of such a space has nonempty intersection) then the product  $\prod_{\alpha \in A} X_\alpha$  is also hyperconnected, in particular, connected. Let us note that hyperconnected spaces are not Hausdorff, and there are examples of hyperconnected  $T_1$ -spaces (some of them can even be found in [4]).

In this article I consider Question 1.2 in the realm of Hausdorff spaces.

In particular, I will show the following:

- (a) If  $X_\alpha$ ,  $\alpha \in A$ , is an infinite system of connected spaces such that infinitely many of them are completely Hausdorff topological spaces then the product  $\prod_{\alpha \in A} X_\alpha$  can be decomposed into continuum many disjoint nonempty open subsets, in particular, it is disconnected.
- (b) If  $X_\alpha$ ,  $\alpha \in A$ , is an infinite system of Brown Hausdorff topological spaces then the product  $\prod_{\alpha \in A} X_\alpha$  is also Brown Hausdorff, and hence, it is connected.

## 2. Auxiliary notions and facts

Recall that a topological space  $X$  is called *completely Hausdorff* or *functionally Hausdorff* if for any distinct points  $x, y \in X$  there exists a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $f(x) \neq f(y)$ .

It is easy to see that every connected completely Hausdorff space has cardinality greater than or equal to  $\mathfrak{c}$ .

**Example 2.1.** Let  $X$  be the topological space  $(\mathbb{R}, \tau)$ , where  $\mathbb{R}$  is the set of reals, and  $\tau$  is the topology on  $\mathbb{R}$  generated by the family  $\{(a, b) \setminus M : a < b, M \subset \mathbb{R}, |M| \leq \aleph_0\}$ . Let us note that  $X$  is a connected homogeneous completely Hausdorff topological space without regular points. (A point  $p$  of a topological space  $T$  is called a *regular point* of  $T$  if for any closed subset  $F$  of  $T$  such that

$p \notin F$  there are two disjoint open subsets  $U$  and  $V$  of  $T$  such that  $p \in U$  and  $F \subseteq V$ .) Observe that the identity mapping from  $X$  to  $\mathbb{R}$  is continuous.

A topological space  $X$  is called a *Urysohn space* if for any distinct points  $x, y \in X$  there are open neighborhoods  $U$  and  $V$  of  $x$  and  $y$ , respectively, such that  $U$  and  $V$  have disjoint closures.

It is evident that each completely Hausdorff space is Urysohn, and there are different examples of countable connected Urysohn spaces (which cannot be completely Hausdorff because of their cardinality), see for examples [7], [8], [9], [14], [17], [19]. The examples from [17], [19] have at least one regular point.

A topological space  $X$  is called a *Brown space* if every pair of open nonempty subsets of  $X$  has nonempty intersection of their closures (cf. [1]).

It is evident that every hyperconnected space is Brown, and every Brown space is connected. There are different examples of connected countable Hausdorff Brown spaces (which can neither have regular points nor be Urysohn by its definition), see for examples [2], [5], [10], [13], [16].

Recall [3] that a space  $X$  possesses

- (1) a *DOF-property*, if there exists a countable sequence  $\{O_i\}_{i=1,2,\dots}$  of open subsets of  $X$  such that  $\text{Cl}_X O_{i+1} \subsetneq O_i$  for each  $i = 1, 2, \dots$  and  $\bigcap_{i \in \mathbb{N}} O_i \neq \emptyset$ ;
- (2) an *IOF-property*, if there exists a countable sequence  $\{O_i\}_{i=1,2,\dots}$  of open subsets of  $X$  such that  $\emptyset \neq O_1 \subset \text{Cl}_X O_i \subsetneq O_{i+1}$  for each  $i = 1, 2, \dots$  and  $\bigcup_{i \in \mathbb{N}} O_i = X$ .

**Proposition 2.2** ([3]). *Let  $X_\alpha, \alpha \in A$ , be an infinite system of connected spaces such that infinitely many of them possess the DOF-property (or IOF-property). Then the product  $\prod_{\alpha \in A} X_\alpha$  can be decomposed into continuum many disjoint nonempty open subsets.*

### 3. Disconnectedness of products of completely Hausdorff spaces with the box topology

**Proposition 3.1.** *Let  $f: X \rightarrow Y$  be a continuous mapping of a topological space  $X$  onto a topological space  $Y$  with the DOF-property. Then  $X$  possesses the DOF-property.*

PROOF: Let  $\{O_i\}_{i \in \mathbb{N}}$  be a countable sequence of open subsets of  $Y$  such that  $\text{Cl}_Y O_{i+1} \subsetneq O_i$  for each  $i = 1, 2, \dots$  and  $\bigcap_{i=1,2,\dots} O_i \neq \emptyset$ . Put  $V_i = f^{-1}O_i$  for each  $i \geq 1$ . Note that  $\{V_i\}_{i \in \mathbb{N}}$  is a countable sequence of open subsets of  $X$  such that  $\bigcap_{i \in \mathbb{N}} V_i = f^{-1}(\bigcap_{i \in \mathbb{N}} O_i) \neq \emptyset$ ,  $\text{Cl}_X V_{i+1} \subset f^{-1}\text{Cl}_Y O_{i+1} \subset V_i$  and  $\text{Cl}_X V_{i+1} \neq V_i$

(since  $V_i \setminus \text{Cl}_X V_{i+1} \supseteq f^{-1}O_i \setminus f^{-1}\text{Cl}_Y O_{i+1} = f^{-1}(O_i \setminus \text{Cl}_Y O_{i+1}) \neq \emptyset$ ) for each  $i = 1, 2, \dots$ . Hence  $X$  possesses the DOF-property.  $\square$

**Corollary 3.2.** *Every connected completely Hausdorff topological space  $X$  possesses DOF-property.*

PROOF: Consider two distinct points  $x, y \in X$  and a continuous mapping  $f: X \rightarrow \mathbb{R}$  of  $X$  to the real line such that  $f(x) \neq f(y)$ . Let us note that the image  $Y = f(X)$  of  $X$  is a nondegenerated connected subset of the real line. Since  $Y$  is a  $T_1$  space with at least one regular non-isolated point, the space  $Y$  possesses the DOF-property. By Proposition 3.1 the space  $X$  also possesses the DOF-property.  $\square$

**Remark 3.3.** Every infinite completely Hausdorff space can be continuously mapped onto a subspace of the real line with at least one non-isolated point. This implies that every such a space has the DOF-property.

**Theorem 3.4.** *Let  $X_\alpha, \alpha \in A$ , be an infinite system of connected spaces such that infinitely many of them are completely Hausdorff topological spaces. Then the product  $\prod_{\alpha \in A} X_\alpha$  can be decomposed into continuum many disjoint nonempty open subsets.*

PROOF: Apply Corollary 3.2 and Proposition 2.2 (the case of DOF-property).  $\square$

We get immediately

**Corollary 3.5.** *The product  $X^{\aleph_0}$ , where  $X$  is the space from Example 2.1, with the box topology can be decomposed into continuum many disjoint nonempty open subsets.*

#### 4. Connectedness of products of Brown Hausdorff spaces with the box topology

**Proposition 4.1.** *Let  $X_\alpha, \alpha \in A$ , be any system of Brown spaces. Then the product  $\prod_{\alpha \in A} X_\alpha$  is also Brown.*

PROOF: Let  $U$  and  $V$  be two nonempty open subsets of  $\prod_{\alpha \in A} X_\alpha$ . Consider a point  $x \in U$  and a point  $y \in V$ . Let  $B = \prod_{\alpha \in A} B_\alpha$  and  $C = \prod_{\alpha \in A} C_\alpha$  be basic open neighborhoods of  $x$  and  $y$ , respectively, such that  $B \subset U$  and  $C \subset V$ . Note that for each  $\alpha \in A$  the space  $X_\alpha$  is Brown. So  $\text{Cl}_{X_\alpha} B_\alpha \cap \text{Cl}_{X_\alpha} C_\alpha \neq \emptyset$ . Then put  $X = \prod_{\alpha \in A} X_\alpha$  and observe that  $\text{Cl}_X B \cap \text{Cl}_X C = (\prod_{\alpha \in A} \text{Cl}_{X_\alpha} B_\alpha) \cap (\prod_{\alpha \in A} \text{Cl}_{X_\alpha} C_\alpha) = \prod_{\alpha \in A} ((\text{Cl}_{X_\alpha} B_\alpha) \cap (\text{Cl}_{X_\alpha} C_\alpha)) \neq \emptyset$ . Hence, the space  $\prod_{\alpha \in A} X_\alpha$  is Brown.  $\square$

**Corollary 4.2** ([1]). *Let  $X_\alpha$ ,  $\alpha \in A$ , be an infinite system of Brown spaces. Then the topological product  $\prod_{\alpha \in A} X_\alpha$  is Brown.*

PROOF: Recall, see [1], that a continuous image of a Brown space is also Brown. Since the identity mapping  $\text{id}: \prod_{\alpha \in A} X_\alpha \rightarrow \prod_{\alpha \in A} X_\alpha$  is a continuous bijection, it follows from Proposition 4.1 that the space  $\prod_{\alpha \in A} X_\alpha$  is Brown.  $\square$

**Theorem 4.3.** *Let  $X_\alpha$ ,  $\alpha \in A$ , be an infinite system of Brown Hausdorff topological spaces. Then the product  $\prod_{\alpha \in A} X_\alpha$  is Brown Hausdorff.*

PROOF: Apply Proposition 4.1 and the known fact that the product of Hausdorff spaces with the box topology is also Hausdorff.  $\square$

We get immediately

**Corollary 4.4.** *Any product  $X^\alpha$ , where  $X$  is the Bing space, see [2], the Golomb space, see [5], the Kirch space, see [10], the Lawrence space, see [13], or the Ritter space, see [16], with the box topology is connected. (Recall that all these spaces  $X$  are Hausdorff and Brown.)*

## 5. Box products of countable connected Urysohn spaces

**Remark 5.1.** Since the Roy space  $R$ , see [17], and the Vought space  $V$ , see [19], two countable connected Urysohn spaces, have regular non-isolated points, the products  $R^{\aleph_0}$  and  $V^{\aleph_0}$  with the box topology can be decomposed into continuum many disjoint nonempty open subsets by Proposition 2.2 (the case of DOF-property).

The following proposition is evident.

**Proposition 5.2.** *Each countable connected homogeneous Hausdorff space has no regular points.*

In [18] (cf. [8]) it was suggested a way to produce countable connected Urysohn homogeneous spaces, and hence (by Proposition 5.2) without regular points.

**Example 5.3.** Let  $X$  be any connected countable Urysohn space, see [7], [8], [9], [14], [17], [19]. For each positive integer  $i$  consider a copy  $X_i$  of  $X$  with two distinct points  $p_i, q_i$  in  $X_i$ . In the topological sum  $\bigoplus_{i \in \mathbb{N}} X_i$  identify points  $q_i$  and  $p_{i+1}$  for every  $i \geq 1$ . Denote the quotient space by  $X_\omega$  and the quotient mapping by  $pr$ . Note that the space  $X_\omega$  is countable connected and Urysohn. Moreover, if  $X$  does not have regular points then neither does  $X_\omega$ .

**Proposition 5.4.** *The space  $X_\omega$  possesses the IOF-property.*

PROOF: Let  $U_i$  and  $V_i$  be open neighborhoods of  $p_i$  and  $q_i$  in  $X_i$  with disjoint closures. Put  $O_1 = pr(X_1 \setminus Cl_{X_1} V_1)$ . Then for each positive integer  $i \geq 2$  put

$$O_i = pr((X_1 \oplus \dots \oplus X_i) \setminus Cl_{X_i} V_i).$$

Note that all  $O_i$  are open subsets of  $X_\omega$  such that  $\emptyset \neq O_1 \subset Cl_X O_i \subsetneq O_{i+1}$  for each  $i = 1, 2, \dots$  and  $\bigcup_{i \in \mathbb{N}} O_i = X_\omega$ . Indeed,  $pr(p_1) \in O_1$ , hence  $O_1 \neq \emptyset$ . Then for each integer  $i \geq 2$  we have

$$Cl_{X_\omega} O_i \subseteq pr((X_1 \oplus \dots \oplus X_i) \setminus V_i) \subseteq O_{i+1}.$$

Moreover,  $pr(q_i) \in O_{i+1} \setminus Cl_{X_\omega} O_i$  and  $\bigcup_{i \in \mathbb{N}} O_i \supseteq \bigcup_{i \in \mathbb{N}} pr(X_1 \oplus \dots \oplus X_i) = X_\omega$ . Hence, the space  $X_\omega$  possesses the IOF-property.  $\square$

**Example 5.5.** Let us extend the space  $X_\omega$  by a point  $p$  as follows. In the union  $X_\omega \cup \{p\}$  a base at the point  $p$  consists of the sets  $\{p\} \cup (X_\omega \setminus \bigcup_{i \leq k} pr X_i)$ ,  $k = 1, 2, \dots$ . For the points of the part  $X_\omega$  we take any base from the space  $X_\omega$ . Denote the new space by  $X_\omega^+$ . It is easy to see that the space  $X_\omega^+$  is countable connected Urysohn, and the point  $p$  is a regular point of  $X_\omega^+$ .

**Corollary 5.6.** *The product  $(X_\omega)^{\aleph_0}$  ( $(X_\omega^+)^{\aleph_0}$ , respectively) with the box topology can be decomposed into continuum many disjoint nonempty open subsets.*

PROOF: Apply Proposition 5.4 and Proposition 2.2 (for both cases).  $\square$

In the end one can ask

**Question 5.7.** Are there countable connected Urysohn spaces  $X_i$ ,  $i = 1, 2, \dots$ , such that the product  $\prod_{i=1,2,\dots} X_i$  is connected?

In connection to the question above it would be interesting to precise what examples from [7], [8], [9], [14] do not have regular points.

**Acknowledgement.** The author thanks the referee for his/her valuable remarks.

### REFERENCES

- [1] Acosta G., Madriz-Mendoza M., Dominguez J.D.C.A., *Brown spaces and the Golomb topology*, Open Acc. J. Math. Theor. Phys. **1** (2018), no. 6, 242–247.
- [2] Bing R.H., *A connected countable Hausdorff space*, Proc. Amer. Math. Soc. **4** (1953), 474.
- [3] Chatyrko V. A., Karassev A., *The (dis)connectedness of products in the box topology*, Questions Answers Gen. Topology **31** (2013), no. 1, 11–21.
- [4] Chatyrko V. A., Nyagahakwa V., *Vitali selectors in topological groups and related semi-groups of sets*, Questions Answers Gen. Topology **33** (2015), no. 2, 93–102.

- [5] Golomb S. W., *A connected topology for integers*, Amer. Math. Monthly **66** 8 (1959), no. 8, 663–665.
- [6] Halimskii E. D., *The topologies of generalized segments*, Dokl. Akad. Nauk SSSR **189** (1969), 740–743.
- [7] Hewitt E., *On two problems of Urysohn*, Ann. of Math. **47** (1946), no. 3, 503–509.
- [8] Jones F. B., Stone A. H., *Countable locally connected Urysohn spaces*, Colloq. Math. **22** (1971), 239–244.
- [9] Kannan V., Rajagopalan M., *On countable locally connected spaces*, Colloq. Math. **29** (1974), 93–100, 159.
- [10] Kirch A. M., *A countable connected, locally connected Hausdorff space*, Amer. Math. Monthly **76** (1969), 169–171.
- [11] Knight C. J., *Box topologies*, Quart. J. Math. Oxford Ser. (2) **15** (1964), 41–54.
- [12] Kong T. Y., Kopperman R., Meyer P. R., *A topological approach to digital topology*, Amer. Math. Monthly **98** (1991), no. 10, 901–917.
- [13] Lawrence L. B., *Infinite-dimensional countable connected Hausdorff spaces*, Houston J. Math. **20** (1994), no. 3, 539–546.
- [14] Miller G. G., *Countable connected spaces*, Proc. Amer. Math. Soc. **26** (1970), 355–360.
- [15] Munkres J. R., *Topology*, Prentice Hall, Upper Saddle River, 2000.
- [16] Ritter G. X., *A connected locally connected, countable Hausdorff space*, Amer. Math. Monthly **83** (1976), no. 3, 185–186.
- [17] Roy P., *A countable connected Urysohn space with a dispersion point*, Duke Math. J. **33** (1966), 331–333.
- [18] Shimrat M., *Embedding in homogeneous spaces*, Quart. J. Math. Oxford Ser. (2) **5** (1954), 304–311.
- [19] Vought E. J., *A countable connected Urysohn space with a dispersion point that is regular almost everywhere*, Colloq. Math. **28** (1973), 205–209.

V. A. Chatyrko:

DEPARTMENT OF MATHEMATICS, LINKÖPING UNIVERSITY, UNIVERSITETSVÄGEN,  
581 83 LINKÖPING, SWEDEN

*E-mail:* vitalij.tjatyрко@liu.se

(Received August 6, 2020, revised October 14, 2020)