

On zero-dimensionality of subgroups of locally compact groups

DMITRII B. SHAKHMATOV

Abstract. Improving the recent result of the author we show that $\text{ind } H = 0$ is equivalent to $\dim H = 0$ for every subgroup H of a Hausdorff locally compact group G .

Keywords: zero-dimensionality, covering dimension, inductive dimension, subgroup, locally compact group

Classification: Primary 22A05, 22D05, 54F45; Secondary 54D45, 54H99

All topological groups considered in this note are assumed to be Hausdorff. A subset of a topological space is said to be *clopen* if it is both open and closed. A topological space X is *zero-dimensional* if it has a base consisting of clopen sets (i.e. if $\text{ind } X = 0$), and X is *strongly zero-dimensional* if every (locally) finite open cover of X consisting of functionally open sets has a finite disjoint clopen refinement (i.e. if $\dim X = 0$). Strongly zero-dimensional spaces are zero-dimensional [1, Theorem 6.2.6] but not vice versa [1, Example 6.2.20]. However, the implication can be reversed for totally bounded groups: Recently, the author proved that a zero-dimensional subgroup of a compact group is strongly zero-dimensional [3, Corollary 3.4]. The aim of our note is to extend this result over subgroups of locally compact groups.

Theorem. *A zero-dimensional subgroup of a locally compact group is strongly zero-dimensional.*

In the proof of this theorem, we need the notion of \mathbb{R} -factorizable group [4]: A topological group G is *\mathbb{R} -factorizable* if for every real-valued continuous function $f : G \rightarrow \mathbb{R}$ defined on G there exist a topological group H with a countable base, a continuous homomorphism $\pi : G \rightarrow H$ and a continuous mapping $\varphi : H \rightarrow \mathbb{R}$ such that $f = \varphi \circ \pi$. The following proposition formally improves [3, Theorem 3.3]:

Proposition. *A zero-dimensional topological group having an open \mathbb{R} -factorizable subgroup is strongly zero-dimensional.*

PROOF: Let H be an open \mathbb{R} -factorizable subgroup of a zero-dimensional topological group G . Being a subspace of a zero-dimensional space G , H is zero-dimensional, and since H is \mathbb{R} -factorizable, it is strongly zero-dimensional by [3, Theorem 3.3].

The author would like to thank cordially Dikran Dikranjan for his suggestion that [3, Corollary 3.4] might admit the improvement stated in the main theorem.

Observe that H is clopen in G , as is every open subgroup of any topological group [2, Chapter 2, Theorem 5.5]. Since G can be covered by disjoint clopen copies of H (namely, by some translations of H) and H is strongly zero-dimensional, G is also strongly zero-dimensional [1, Theorem 6.2.13]. \square

PROOF OF THEOREM: Let H be a zero-dimensional subgroup of a locally compact group G , and let U be an open neighbourhood of the neutral element of G having compact closure \bar{U} in G . Then G^* , the smallest subgroup of G that contains \bar{U} , is σ -compact. As a subgroup of a σ -compact group, $H^* = H \cap G^*$ is \mathbb{R} -factorizable [4, Corollary 1.13]. Since H^* contains the non-empty open set $U \cap H$ and is a subgroup of H , H^* is open in H [2, Chapter 2, Theorem 5.5]. Now, Proposition finishes the proof. \square

In conclusion, let us mention that, quite surprisingly, the following question remains open:

Question. Is there a normal zero-dimensional group which is not strongly zero-dimensional?

Even if we drop “normal” here, the answer to this question seems to be unknown.

REFERENCES

- [1] Engelking R., *General Topology*, Warszawa, PWN, 1977.
- [2] Hewitt E., Ross K.A., *Abstract Harmonic Analysis, vol. 1. Structure of Topological Groups. Integration Theory. Group Representations*, Die Grundlehren der mathematischen Wissenschaften, Bd. 115, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1963.
- [3] Shakhmatov D.B., *Imbeddings into topological groups preserving dimensions*, Topology Appl. **36** (1990), 181–204.
- [4] Tkačenko M.G., *Factorization theorems for topological groups and their applications*, Topology Appl. **38** (1991), 21–37.

DEPARTMENT OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY,
119 889 Moscow, USSR

(Received March 26, 1991)