

Inductive limit topologies on Orlicz spaces

MARIAN NOWAK

Abstract. Let L^φ be an Orlicz space defined by a convex Orlicz function φ and let E^φ be the space of finite elements in L^φ (= the ideal of all elements of order continuous norm). We show that the usual norm topology \mathcal{T}_φ on L^φ restricted to E^φ can be obtained as an inductive limit topology with respect to some family of other Orlicz spaces. As an application we obtain a characterization of continuity of linear operators defined on E^φ .

Keywords: Orlicz spaces, inductive limit topologies, convex functions

Classification: 46E30

1. Introduction and preliminaries.

In [1] and [2] Davis, Murray and Weber discussed the spaces

$$L^{p+} = \bigcup_{p < t < \infty} L^t[0,1] \quad \text{and} \quad l^{p-} = \bigcup_{1 \leq t < p} l^t \quad (1 < p \leq \infty)$$

(endowed with the appropriate inductive limit topologies) which turned out to be distinct from the spaces L^p and l^p , respectively.

Moreover, in [8] it is proved that if $S \subset [0, \infty)$ with $\inf S \notin S$ or $\sup S \notin S$ and μ is an infinite atomless measure (resp. $\sup S \notin S$ and μ is the counting measure on \mathbb{N}), there is no Orlicz function φ such that:

$$E^\varphi = \text{Lin} \bigcup_{p \in S} L^p \quad \text{or} \quad L^\varphi = \text{Lin} \bigcup_{p \in S} L^p.$$

On the other hand, Krasnoselskii and Rutickii [3, p. 60] showed that if μ is the finite Lebesgue measure, then

$$L^1 = \bigcup_{\varphi} L^\varphi,$$

where φ are taken over the family of all N -functions. This equality was a starting point for many results concerning a representation of an Orlicz space L^φ or a space E^φ as the union of some families of Orlicz spaces which they contain properly (see [4], [7], [9], [12]).

In [7] for a convex Orlicz function φ we found the set Ψ^φ of N -functions such that:

$$E^\varphi = \bigcup_{\psi \in \Psi^\varphi} E^\psi = \bigcup_{\psi \in \Psi^\varphi} L^\psi.$$

In this paper we show that the appropriate inductive limit topologies on E^φ defined with respect to these representations coincide with the norm topology \mathcal{T}_φ on L^φ restricted to E^φ .

We now recall some notation and terminology concerning Orlicz spaces (see [3], [5], [11] for more details).

By an Orlicz function we mean a function $\varphi : [0, \infty) \rightarrow [0, \infty]$ which is non-decreasing, left continuous, continuous at zero with $\varphi(0) = 0$, and not identically equal to zero.

We shall say that an Orlicz function φ jumps to ∞ , whenever there is a number $u_0 > 0$ such that $\varphi(u) = \infty$ for $u > u_0$. We shall say that φ vanishes near zero, whenever $\varphi(u) = 0$ for $0 \leq u \leq u_0$ for some $u_0 > 0$.

An Orlicz function φ is called convex, if $\varphi(\alpha u + \beta v) \leq \alpha\varphi(u) + \beta\varphi(v)$ for $\alpha, \beta \geq 0$, $\alpha + \beta = 1$. A convex Orlicz function is usually called a Young function. A convex Orlicz function φ , vanishing only at 0 and taking only finite values is called an N -function if $\varphi(u)/u \rightarrow 0$ as $u \rightarrow 0$ and $\varphi(u)/u \rightarrow \infty$ as $u \rightarrow \infty$. By Φ_N we will denote the collection of all N -functions.

For a convex Orlicz function φ we denote by φ^* the function complementary to φ in the sense of Young, i.e.

$$\varphi^*(v) = \sup\{uv - \varphi(u) : u \geq 0\} \text{ for } v \geq 0.$$

For a set Ψ of convex Orlicz functions we will write

$$\Psi^* = \{\psi^* : \psi \in \Psi\}.$$

Throughout this paper we will write: $\varphi_p(u) = u^p$ for $u \geq 0$, where $p \geq 1$ and

$$\varphi_0(u) = \begin{cases} 0 & \text{for } 0 \leq u \leq 1, \\ 1 & \text{for } u > 1 \end{cases}, \text{ and } \varphi_\infty(u) = \begin{cases} 0 & \text{for } 0 \leq u \leq 1, \\ \infty & \text{for } u > 1 \end{cases}.$$

We shall say that two Orlicz functions ψ and φ are equivalent for all u (resp. for small u , resp. for large u), in symbols $\psi \stackrel{a}{\sim} \varphi$ (resp. $\psi \stackrel{s}{\sim} \varphi$, resp. $\psi \stackrel{l}{\sim} \varphi$) if there exist constants $a, b, c, d > 0$ such that $a\psi(bu) \leq \varphi(u) \leq c\psi(du)$ for all $u \geq 0$ (resp. for $0 \leq u \leq u_0$, resp. for $u \geq u_0$), where $u_0 > 0$.

We say that an Orlicz function φ increases essentially more rapidly than any other ψ for all u (resp. for small u , resp. for large u), in symbols $\psi \stackrel{a}{\ll} \varphi$ (resp. $\psi \stackrel{s}{\ll} \varphi$, resp. $\psi \stackrel{l}{\ll} \varphi$) if for any $c > 0$, $\psi(cu)/\varphi(u) \rightarrow 0$ as $u \rightarrow 0$ and $u \rightarrow \infty$ (resp. as $u \rightarrow 0$, resp. $u \rightarrow \infty$) (see [3, p. 114]).

It is known that $\psi \stackrel{a}{\ll} \varphi$ (resp. $\psi \stackrel{s}{\ll} \varphi$, resp. $\psi \stackrel{l}{\ll} \varphi$) implies $\varphi^* \stackrel{a}{\ll} \psi^*$ (resp. $\varphi^* \stackrel{s}{\ll} \psi^*$, resp. $\varphi^* \stackrel{l}{\ll} \psi^*$) (see [3, Lemma 13.1]).

Let (Ω, Σ, μ) be a positive measure space, and let L^0 denote the set of equivalence classes of all real valued μ -measurable functions defined and finite a.e. on Ω . An Orlicz function φ determines a functional $m_\varphi : L^0 \rightarrow [0, \infty]$ by the formula:

$$m_\varphi(x) = \int_\Omega \varphi(|x(t)|) d\mu.$$

The Orlicz space determined by φ is the ideal of L^0 defined by

$$L^\varphi = \{x \in L^0 : m_\varphi(\lambda x) < \infty \text{ for some } \lambda > 0\}.$$

The functional m_φ restricted to L^φ is an orthogonally additive modular (see [6]).

L^φ can be equipped with the complete metrizable topology \mathcal{T}_φ of the Riesz F -norm

$$|x|_\varphi = \inf\{\lambda > 0 : m_\varphi(x/\lambda) \leq \lambda\}.$$

Moreover, if φ is convex, then the topology \mathcal{T}_φ is generated by the norm

$$\|x\|_\varphi = \inf\{\lambda > 0 : m_\varphi(x/\lambda) \leq 1\}.$$

Let

$$E^\varphi = \{x \in L^0 : m_\varphi(\lambda x) < \infty \text{ for all } \lambda > 0\}.$$

Then E^φ is a closed ideal of L^φ , and it is well known that E^φ coincides with the ideal of all elements of L^φ with order continuous F -norm $|\cdot|_\varphi$. It is known that $L^\varphi = E^\varphi$ if φ satisfies the Δ_2 -condition, i.e.

$$\limsup \frac{\varphi(2u)}{\varphi(u)} < \infty \text{ as } u \rightarrow 0 \text{ and } u \rightarrow \infty.$$

If μ is the counting measure on the set \mathbb{N} of all natural numbers, we will write l^φ and h^φ instead of L^φ and E^φ , respectively. By c_0 we will denote the space of all sequences that are convergent to 0.

Given a linear topological space (X, ξ) , by $(X, \xi)^*$ we will denote its topological dual.

2. Some equalities among Orlicz spaces.

In this section we present some equalities among Orlicz spaces, obtained in [7], that are of the key importance in the paper.

Let Φ_1 be the set of all convex Orlicz functions φ taking only finite values and such that $\varphi(u)/u \rightarrow 0$ as $u \rightarrow 0$.

Denote by

$$\begin{aligned} \Phi_{11} &= \{\varphi \in \Phi_1 : \varphi(u) > 0 \text{ for } u > 0 \text{ and } \varphi(u)/u \rightarrow \infty \text{ as } u \rightarrow \infty\}, \\ \Phi_{12} &= \{\varphi \in \Phi_1 : \varphi(u) > 0 \text{ for } u > 0 \text{ and } \varphi(u)/u \rightarrow a \text{ as } u \rightarrow \infty, a > 0\}, \\ \Phi_{13} &= \{\varphi \in \Phi_1 : \varphi(u) = 0 \text{ near zero and } \varphi(u)/u \rightarrow \infty \text{ as } u \rightarrow \infty\}, \\ \Phi_{14} &= \{\varphi \in \Phi_1 : \varphi(u) = 0 \text{ near zero and } \varphi(u)/u \rightarrow a \text{ as } u \rightarrow \infty, a > 0\}. \end{aligned}$$

Then $\Phi_1 = \bigcup_{i=1}^4 \Phi_{1i}$, where the sets are pairwise disjoint. It is seen that $\Phi_{11} = \Phi_N$.

Theorem 2.1 [7, Theorems 1.1–1.4, Theorem 1.7]. *Let $\varphi \in \Phi_{1i}$ ($i = 1, 2, 3, 4$). Then the following equalities hold:*

$$E^\varphi = \bigcup_{\psi \in \Psi_{1i}^\varphi} E^\psi = \bigcup_{\psi \in \Psi_{1i}^\varphi} L^\psi,$$

where:

$$\begin{aligned} \Psi_{11}^\varphi &= \{\psi \in \Phi_N : \varphi \ll^a \psi\}, \\ \Psi_{12}^\varphi &= \{\psi \in \Phi_N : \varphi \ll^s \psi\}, \\ \Psi_{13}^\varphi &= \{\psi \in \Phi_N : \varphi \ll^l \psi\}, \\ \Psi_{14}^\varphi &= \Phi_N. \end{aligned}$$

Moreover, if μ is an atomless measure or the counting measure on \mathbb{N} , then for each $\psi \in \Psi_{1i}^\varphi$, the strict inclusion $L^\psi \subsetneq E^\varphi$ holds.

Next, let Φ_2 be the set of all convex Orlicz functions φ vanishing only at 0 and such that $\varphi(u)/u \rightarrow \infty$ as $u \rightarrow \infty$.

Denote by

$$\begin{aligned} \Phi_{21} &= \{\varphi \in \Phi_2 : \varphi(u) < 0 \text{ for } u > 0 \text{ and } \varphi(u)/u \rightarrow 0 \text{ as } u \rightarrow 0\}, \\ \Phi_{22} &= \{\varphi \in \Phi_2 : \varphi \text{ jumps to } \infty \text{ and } \varphi(u)/u \rightarrow 0 \text{ as } u \rightarrow 0\}, \\ \Phi_{23} &= \{\varphi \in \Phi_2 : \varphi(u) < 0 \text{ for } u > 0 \text{ and } \varphi(u)/u \rightarrow a \text{ as } u \rightarrow 0, a > 0\}, \\ \Phi_{24} &= \{\varphi \in \Phi_2 : \varphi \text{ jumps to } \infty \text{ and } \varphi(u)/u \rightarrow a \text{ as } u \rightarrow 0, a > 0\}. \end{aligned}$$

Then $\Phi_2 = \bigcup_{i=1}^4 \Phi_{2i}$ and $\Phi_{21} = \Phi_N$.

Theorem 2.2 [7, Theorems 2.1–2.4, Theorem 2.6]. *Let $\varphi \in \Phi_{2i}$ ($i = 1, 2, 3, 4$). Then the following equalities hold:*

$$L^\varphi = \bigcap_{\psi \in \Psi_{2i}^\varphi} L^\psi = \bigcap_{\psi \in \Psi_{2i}^\varphi} E^\psi,$$

where:

$$\begin{aligned} \Psi_{21}^\varphi &= \{\psi \in \Phi_N : \psi \ll^a \varphi\}, \\ \Psi_{22}^\varphi &= \{\psi \in \Phi_N : \psi \ll^s \varphi\}, \\ \Psi_{23}^\varphi &= \{\psi \in \Phi_N : \psi \ll^l \varphi\}, \\ \Psi_{24}^\varphi &= \Phi_N. \end{aligned}$$

At last, according to [7, Lemma 3.1, Theorem 3.3] we have

Theorem 2.3. *Let φ_1 and φ_2 be a pair of complementary convex Orlicz functions, i.e. $\varphi_1^* = \varphi_2$. Then $\varphi_1 \in \Phi_{1i}$ iff $\varphi_2 \in \Phi_{2i}$ ($i = 1, 2, 3, 4$), and moreover, the sets $\Psi_{1i}^{\varphi_1}$ and $\Psi_{2i}^{\varphi_2}$ are mutually related in such a way that:*

$$(\Psi_{1i}^{\varphi_1})^* = \Psi_{2i}^{\varphi_2} \quad \text{and} \quad (\Psi_{2i}^{\varphi_2})^* = \Psi_{1i}^{\varphi_1}.$$

3. Inductive limit topologies on E^φ .

Let $\varphi \in \Phi_{1i}$ ($i = 1, 2, 3, 4$). Then in view of Theorem 2.1, one can consider on E^φ the inductive limit topologies $\mathcal{T}_{I_1}^\varphi$ and $\mathcal{T}_{I_2}^\varphi$ with respect to the families $\{(E^\psi, \mathcal{T}_\psi |_{E^\psi}) : \psi \in \Psi_{1i}^\varphi\}$ and $\{(L^\psi, \mathcal{T}_\psi) : \psi \in \Psi_{1i}^\varphi\}$, respectively (see [10, Chapter V, § 2]). Thus $\mathcal{T}_{I_1}^\varphi$ (resp. $\mathcal{T}_{I_2}^\varphi$) is the finest of all locally convex topologies ξ on E^φ that satisfy, for each $\psi \in \Psi_{1i}^\varphi$, the condition $\xi |_{E^\psi} \subset \mathcal{T}_\psi |_{E^\psi}$ (resp. $\xi |_{L^\psi} \subset \mathcal{T}_\psi$). It is seen that

$$(3.1) \quad \mathcal{T}_\varphi |_{E^\varphi} \subset \mathcal{T}_{I_2}^\varphi \subset \mathcal{T}_{I_1}^\varphi.$$

Our aim is to show that the topology $\mathcal{T}_\varphi |_{E^\varphi}$ coincides with $\mathcal{T}_{I_1}^\varphi$ and $\mathcal{T}_{I_2}^\varphi$. For this purpose, the following theorem will be of importance.

Theorem 3.1. *Let $\varphi \in \Phi_1$ and let μ be a σ -finite measure. Then for a linear functional f on E^φ the following statements are equivalent:*

- (a) f is $\mathcal{T}_{I_1}^\varphi$ -continuous.
- (b) There exists a unique $y \in L^{\varphi^*}$ such that

$$f(x) = f_y(x) = \int_{\Omega} x(t)y(t) d\mu \quad \text{for all } x \in E^\varphi.$$

PROOF: (a) \Rightarrow (b). Let $\varphi \in \Phi_{1i}$ ($i = 1, 2, 3, 4$). Then for each $\psi \in \Psi_{1i}^\varphi$, the functional $f |_{E^\psi}$ is continuous for $\mathcal{T}_\psi |_{E^\psi}$, so according to [5, Chapter II, § 3, Theorem 2] there exists a unique function $y_\psi \in L^{\psi^*}$ such that

$$(+) \quad f(x) = \int_{\Omega} x(t)y_\psi(t) d\mu \quad \text{for all } x \in E^\psi.$$

Assume that there exist $\psi_1, \psi_2 \in \Psi_{1i}^\varphi$ such that $y_{\psi_1} \neq y_{\psi_2}$, and $f(x) = \int_{\Omega} x(t)y_{\psi_k}(t) d\mu$ for $x \in E^{\psi_k}$, where $k = 1, 2$. Let us assume, for example, that $\mu(\{t \in \Omega : y_{\psi_1}(t) > y_{\psi_2}(t)\}) > 0$, and let $A \subset \{t \in \Omega : y_{\psi_1}(t) > y_{\psi_2}(t)\}$ be a measurable set with $0 < \mu(A) < \infty$. Denoting by χ_A the characteristic function of A , we have $\chi_A \in E^{\psi_1} \cap E^{\psi_2}$, so by (+) we get

$$\int_{\Omega} \chi_A(t)(y_{\psi_1}(t) - y_{\psi_2}(t)) d\mu = \int_A (y_{\psi_1}(t) - y_{\psi_2}(t)) d\mu = 0.$$

This contradiction establishes that there exists a unique

$$y \in \bigcap_{\psi \in \Psi_{1i}^\varphi} L^{\psi^*} \text{ such that } f(x) = \int_{\Omega} x(t)y(t) d\mu \text{ for all } x \in E^\varphi.$$

On the other hand, since $\varphi^* \in \Phi_{2i}$ and $(\Psi_{1i}^\varphi)^* = \Psi_{2i}^{\varphi^*}$ (see Theorem 2.3), according to Theorem 2.2,

$$\bigcap_{\psi \in \Psi_{1i}^\varphi} L^{\psi^*} = \bigcap_{\psi \in (\Psi_{1i}^\varphi)^*} L^\psi = \bigcap_{\psi \in \Psi_{2i}^{\varphi^*}} L^\psi = L^{\varphi^*}.$$

(b) \Rightarrow (a). Let $\varphi \in \Phi_{1i}$ ($i=1,2,3,4$). Then for each $\psi \in \Psi_{1i}^\varphi$, by Theorem 2.3, $\psi^* \in \Psi_{2i}^{\varphi^*}$. Hence $L^{\varphi^*} \subset L^{\psi^*}$, and the functional $f|_{E^\psi}$ is continuous for $\mathcal{T}_\psi|_{E^\psi}$ (see [5, Chapter 2, § 3, Theorem 2]). Therefore, in view of [10, Chapter V, Proposition 5], the functional f is continuous for $\mathcal{T}_{I_1}^\varphi$.

Thus the proof is completed. □

Now we are in a position to prove our main theorem.

Theorem 3.2. *Let $\varphi \in \Phi_1$ and μ be a σ -finite measure. Then the norm topology \mathcal{T}_φ restricted to E^φ coincides with the inductive limit topologies $\mathcal{T}_{I_1}^\varphi$ and $\mathcal{T}_{I_2}^\varphi$, that is*

$$\mathcal{T}_\varphi|_{E^\varphi} = \mathcal{T}_{I_1}^\varphi = \mathcal{T}_{I_2}^\varphi.$$

PROOF: Since the space $(E^\varphi, \mathcal{T}_\varphi|_{E^\varphi})$ is barrelled and $(E^\varphi, \mathcal{T}_\varphi|_{E^\varphi})^* = \{f_y : y \in L^{\varphi^*}\}$ (see [5, Chapter II, § 3, Theorem 2]), the equality $\mathcal{T}_\varphi|_{E^\varphi} = \beta(E^\varphi, L^{\varphi^*})$ holds (see [10, Chapter IV, § 1, Corollary 1]).

On the other hand, the space $(E^\varphi, \mathcal{T}_{I_1}^\varphi)$ is barrelled, because an inductive limit of barrelled spaces is barrelled (see [10, Chapter 2, Proposition 6]). Hence, in view of Theorem 3.1, the equality $\mathcal{T}_{I_1}^\varphi = \beta(E^\varphi, L^{\varphi^*})$ holds. Thus $\mathcal{T}_\varphi|_{E^\varphi} = \mathcal{T}_{I_1}^\varphi$, and by (3.1) our proof is completed. □

4. A characterization of continuity of linear operators on E^φ .

As an application of Theorem 3.2, in view of the general property of inductive limit topologies (see [10, Chapter V, 2, Proposition 5]), we obtain a characterization of linear operators of E^φ into a locally convex space X . The details follow.

Theorem 4.1. *Let $\varphi \in \Phi_{1i}$ ($i = 1, 2, 3, 4$) and let (X, ξ) be a locally convex space. For a linear operator $A : E^\varphi \rightarrow X$, the following statements are equivalent:*

- (a) A is $(\mathcal{T}_\varphi|_{E^\varphi}, \xi)$ -continuous.
- (b) $A|_{E^\psi}$ is $(\mathcal{T}_\psi|_{E^\psi}, \xi)$ -continuous for every $\psi \in \Psi_{1i}^\varphi$.
- (c) $A|_{E^\psi}$ is (\mathcal{T}_ψ, ξ) -continuous for every $\psi \in \Psi_{1i}^\varphi$.

We close this section with an application of Theorem 2.1 and Theorem 4.1 to the spaces: $L^p, L^1 + L^p$ ($p > 1$) and c_0 .

Examples.

A. Let $p > 1$. Then $\varphi_p \in \Phi_{11}$ and in view of Theorem 2.1 and Theorem 4.1 we get the following

Corollary 4.2. *Let $p > 1$. Then the following equalities hold:*

$$L^p = \bigcup_{\psi} E^{\psi} = \bigcup_{\psi} L^{\psi},$$

where the unions are taken over all N -functions ψ such that $\psi(u)/u^p \rightarrow \infty$ as $u \rightarrow 0$ and $u \rightarrow \infty$.

Moreover, if the measure μ is σ -finite, then for a locally convex space (X, ξ) and a linear operator $A : L^p \rightarrow X$, the following statements are equivalent:

- (a) A is (\mathcal{T}_{L^p}, ξ) -continuous.
- (b) $A|_{E^{\psi}}$ is $(\mathcal{T}_{\psi}|_{E^{\psi}}, \xi)$ -continuous for every N -function ψ such that $\psi(u)/u^p \rightarrow \infty$ as $u \rightarrow 0$ and $u \rightarrow \infty$.
- (c) $A|_{L^{\psi}}$ is $(\mathcal{T}_{\psi}, \xi)$ -continuous for every N -function ψ such that $\psi(u)/u^p \rightarrow \infty$ as $u \rightarrow 0$ and $u \rightarrow \infty$.

B. For $p > 1$ let us put

$$\varphi(u) = \begin{cases} u^p & \text{for } 0 \leq u \leq 1, \\ pu + 1 - p & \text{for } u > 1, \end{cases}$$

and let $\varphi'(u) = \min(\varphi_1(u), \varphi_p(u))$. Then φ is a convex Orlicz function and $\varphi \stackrel{a}{\sim} \varphi'$, so $E^{\varphi} = L^{\varphi} = L^{\varphi'} = L^1 + L^p$ and $\mathcal{T}_{\varphi} = \mathcal{T}_{\varphi'}$, where the topology $\mathcal{T}_{\varphi'}$ is generated by the norm:

$$\|x\|_{L^1+L^p} = \inf\{\|x_1\|_{L^1} + \|x_2\|_{L^p} : x = x_1 + x_2, \quad x_1 \in L^1, \quad x_2 \in L^p\}.$$

Since $\varphi \in \Phi_{12}$, according to Theorem 2.1 and Theorem 4.1 we have

Corollary 4.3. *Let $p > 1$. Then the following equalities hold:*

$$L^1 + L^p = \bigcup_{\psi} E^{\psi} = \bigcup_{\psi} L^{\psi},$$

where the unions are taken over the set of all N -functions ψ such that $\psi(u)/u^p \rightarrow \infty$ as $u \rightarrow 0$.

Moreover, if the measure μ is σ -finite, then for a locally convex space (X, ξ) and a linear operator $A : L^1 + L^p \rightarrow X$, the following statements are equivalent:

- (a) A is $(\mathcal{T}_{L^1+L^p}, \xi)$ -continuous.
- (b) $A|_{E^{\psi}}$ is $(\mathcal{T}_{\psi}|_{E^{\psi}}, \xi)$ -continuous for every N -function ψ such that $\psi(u)/u^p \rightarrow \infty$ as $u \rightarrow 0$.
- (c) $A|_{L^{\psi}}$ is $(\mathcal{T}_{\psi}, \xi)$ -continuous for every N -function ψ such that $\psi(u)/u^p \rightarrow \infty$ as $u \rightarrow 0$.

In particular, if the measure μ is finite, then

$$L^1 = \bigcup_{\psi} E^{\psi} = \bigcup_{\psi} L^{\psi},$$

where the unions are taken over the set of all N -functions ψ .

Moreover, for a linear operator $A : L^1 \rightarrow X$, the following statements are equivalent:

- (a) A is (\mathcal{T}_{L^1}, ξ) -continuous.
- (b) $A|_{E^{\psi}}$ is $(\mathcal{T}_{\psi}|_{E^{\psi}}, \xi)$ -continuous for every N -function ψ .
- (c) $A|_{L^{\psi}}$ is $(\mathcal{T}_{\psi}, \xi)$ -continuous for every N -function ψ .

C. Let

$$\varphi(u) = \begin{cases} 0 & \text{for } 0 \leq u \leq 1, \\ u - 1 & \text{for } u > 1. \end{cases}$$

Then φ is a convex Orlicz function and $\varphi \stackrel{s}{\sim} \varphi_0$. Hence $l^{\varphi} = l^{\varphi_0} = l^{\infty}$ and $h^{\varphi} = h^{\varphi_0} = c_0$, and the topology \mathcal{T}_{φ} on l^{φ} agrees with the topology \mathcal{T}_{∞} of the norm $\|x\|_{\infty} = \sup_i |x(i)|$ on l^{∞} . Since $\varphi \in \Phi_{14}$, in view of Theorem 2.1 and Theorem 4.1, we have

Corollary 4.4. *The following equalities hold:*

$$c_0 = \bigcup_{\psi} h^{\psi} = \bigcup_{\psi} l^{\psi},$$

where the unions are taken over the set of all N -functions.

Moreover, for a locally convex space (X, ξ) and a linear operator $A : c_0 \rightarrow X$, the following statements are equivalent:

- (a) A is $(\mathcal{T}_{\infty}|_{c_0}, \xi)$ -continuous.
- (b) $A|_{h^{\psi}}$ is $(\mathcal{T}_{\psi}|_{h^{\psi}}, \xi)$ -continuous for every N -function ψ .
- (c) $A|_{l^{\psi}}$ is $(\mathcal{T}_{\psi}, \xi)$ -continuous for every N -function ψ .

REFERENCES

- [1] Davis H.W., Murray F.J., Weber J.K., *Families of L_p -spaces with inductive and projective topologies*, Pacific J. Math. **34** (1970), 619–638.
- [2] ———, *Inductive and projective limits of L_p -spaces*, Portugal. Math. **38** (1972), 21–29.
- [3] Krasnoselskii M.A., Rutickii Ya.B., *Convex Functions and Orlicz Spaces*, Groningen, 1961.
- [4] Leśniewicz R., *On two equalities for Orlicz spaces*, Bull. Acad. Pol. Sci. **27** (1979), 557–560.
- [5] Luxemburg W.A., *Banach Functions Spaces*, Delft, 1955.
- [6] Musielak J., Orlicz W., *Some remarks on modular spaces*, Bull. Acad. Pol. Sci. **7** (1959), 661–668.
- [7] Nowak M., *Some equalities among Orlicz spaces II*, Bull. Acad. Pol. Sci. **34** (1986), 675–687.
- [8] ———, *Unions and intersections of families of L^p -spaces*, Math. Nachr. **136** (1988), 241–251.
- [9] ———, *Some equalities among Orlicz spaces I*, Comment. Math. **29** (1990), 255–275.
- [10] Robertson A.P., Robertson W.J., *Topological Vector Spaces*, Cambridge, 1973.

- [11] Turpin Ph., *Convexités dans les espaces vectoriels topologiques généraux*, Dissertationes Math. **131** (1976).
- [12] Welland R., *Inclusions relations among Orlicz spaces*, Proc. Amer. Math. Soc. **17** (1966), 135–138.

INSTITUTE OF MATHEMATICS, A. MICKIEWICZ UNIVERSITY, MATEJKI 48/49, 60–769 POZNAŃ,
POLAND

(Received August 8, 1991)