

## Two cardinal inequalities for functionally Hausdorff spaces

ALESSANDRO FEDELI

*Abstract.* In this paper, two cardinal inequalities for functionally Hausdorff spaces are established. A bound on the cardinality of the  $\tau\theta$ -closed hull of a subset of a functionally Hausdorff space is given. Moreover, the following theorem is proved: if  $X$  is a functionally Hausdorff space, then  $|X| \leq 2^{\chi(X)wcd(X)}$ .

*Keywords:* cardinal functions,  $\tau\theta$ -closed sets,  $w$ -compactness degree

*Classification:* 54A25, 54D20

A space  $X$  is said to be functionally Hausdorff if whenever  $x \neq y$  in  $X$  there is a continuous real valued function  $f$  defined on  $X$  such that  $f(x) = 0$  and  $f(y) = 1$ . A well-known Arhangel'skii's theorem states that if  $X$  is a Hausdorff space, then  $|X| \leq 2^{\chi(X)L(X)}$  ([1], [6]). Bella and Cammaroto [2] established some cardinal inequalities for Urysohn spaces that improve, for non regular spaces, the Arhangel'skii's formula. In this paper, a bound on the cardinality of the  $\tau\theta$ -closed hull of a subset of a functionally Hausdorff space and a bound on the cardinality of a functionally Hausdorff space are given. We refer the reader to [3] and [4] for notations and definitions not explicitly given. All topological spaces considered here are assumed to be infinite. Let  $E$  be a set; the cardinality of  $E$  is denoted by  $|E|$ ,  $\mathcal{P}_k(E)$  is the collection of all subsets of  $E$  of cardinality  $\leq k$ .  $\chi(X)$  and  $L(X)$  denote respectively the character and the Lindelöf degree of a space  $X$ .

**Definition 1** [5]. Let  $A$  be a subset of a space  $X$ .  $A$  is called  $\tau$ -open if  $A$  is a union of cozero-sets of  $X$ . The  $\tau$ -closure of  $A$ , denoted by  $cl_\tau(A)$ , is the set of all points  $x \in X$  such that any cozero-set neighbourhood of  $x$  intersects  $A$ . The  $\tau$ -interior of  $A$ , denoted by  $int_\tau(A)$ , is the set of all  $x$  such that there is a cozero-set neighbourhood of  $x$  contained in  $A$ .

**Definition 2.** Let  $X$  be a topological space and  $A$  a subset of  $X$ . The  $\tau\theta$ -closure of  $A$ , denoted by  $cl_{\tau\theta}(A)$ , is the set of all points  $x \in X$  such that  $cl_\tau(V) \cap A \neq \emptyset$  for every open neighbourhood  $V$  of  $x$ .  $A$  is said to be  $\tau\theta$ -closed if  $A = cl_{\tau\theta}(A)$ .

As pointed to me by S. Watson, the  $\tau\theta$ -closure is not in general idempotent.

**Definition 3.** Let  $X$  be a topological space and  $A$  a subset of  $X$ . The  $\tau\theta$ -closed hull of  $A$ , denoted by  $[A]_{\tau\theta}$ , is the smallest  $\tau\theta$ -closed subset of  $X$  containing  $A$ .

Clearly,  $[A]_{\tau\theta} = \bigcap \{F : A \subset F \text{ and } \text{cl}_{\tau\theta}(F) = F\}$ . For every space  $X$  and every  $A \subset X$  we have  $\overline{A} \subset \text{cl}_{\tau\theta}(A) \subset [A]_{\tau\theta} \subset \text{cl}_{\tau}(A)$ . It is obvious that if  $X$  is a Tychonoff space, then  $\overline{A} = \text{cl}_{\tau\theta}(A) = [A]_{\tau\theta} = \text{cl}_{\tau}(A)$  for any  $A \subset X$ .

The next result gives some conditions on a functionally Hausdorff space which are equivalent to  $\text{cl}_{\tau\theta} = \text{cl}_{\tau}$ .

**Proposition 4.** *For a functionally Hausdorff space  $X$  the following conditions are equivalent:*

- (i) *For each  $\tau$ -open set  $V$  of  $X$ ,  $\overline{V} = \text{cl}_{\tau}(V)$ .*
- (ii) *For each open set  $G$  of  $X$ ,  $G \subset \text{int}_{\tau}(\text{cl}_{\tau}(G))$ .*
- (iii) *For each subset  $A$  of  $X$ ,  $\text{cl}_{\tau\theta}(A) = \text{cl}_{\tau}(A)$ .*
- (iv) *For each  $\tau$ -open subset  $V$  of  $X$ ,  $\text{cl}_{\tau\theta}(V) = \text{cl}_{\tau}(V)$ .*

PROOF: (i)  $\Leftrightarrow$  (ii) Lemma 28 in [9]. (ii)  $\Rightarrow$  (iii) Let  $A \subset X$  and  $x \notin \text{cl}_{\tau\theta}(A)$ , then there is an open neighbourhood  $G$  of  $x$  such that  $\text{cl}_{\tau}(G) \cap A = \emptyset$ . By hypothesis  $G \subset \text{int}_{\tau}(\text{cl}_{\tau}(G))$ , then there is a cozero set  $V$  such that  $x \in V \subset \text{cl}_{\tau}(G)$ , so  $V \cap A = \emptyset$  and  $x \notin \text{cl}_{\tau}(A)$ . Hence,  $\text{cl}_{\tau\theta}(A) = \text{cl}_{\tau}(A)$ . (iii)  $\Rightarrow$  (iv) is obvious. (iv)  $\Rightarrow$  (i) Let  $V$  be a  $\tau$ -open subset of  $X$ , by hypothesis  $\text{cl}_{\tau\theta}(V) = \text{cl}_{\tau}(V)$ . Now let  $x \notin \overline{V}$ , then there is an open set  $G$  such that  $x \in G$  and  $G \cap V = \text{cl}_{\tau}(V)$ . Since  $V$  is  $\tau$ -open, we have  $\text{cl}_{\tau}(G) \cap V = \emptyset$ , hence  $x \notin \text{cl}_{\tau\theta}(V)$ . Therefore,  $\overline{V} = \text{cl}_{\tau\theta}(V) = \text{cl}_{\tau}(V)$ . □

**Remark 5.** A functionally Hausdorff space  $X$  is called weakly absolutely closed [8] provided that every  $\tau$ -open filter base on  $X$  has an adherent point. An SW space is a functionally Hausdorff space  $X$  such that every point-separating subalgebra of  $C^*(X)$  which contains the constants is uniformly dense in  $C^*(X)$  [8]. It is worth noting that by Lemma 25 in [9] and Proposition 4, a functionally Hausdorff space  $X$  is weakly absolutely closed iff it is an SW space and  $\text{cl}_{\tau\theta}(A) = \text{cl}_{\tau}(A)$  for every  $A \subset X$ .

The following result gives an upper bound on the  $\tau\theta$ -closed hull.

**Theorem 6.** *Let  $X$  be a functionally Hausdorff space. If  $A$  is a subset of  $X$ , then  $|[A]_{\tau\theta}| \leq |A|^{\chi(X)}$ .*

PROOF: Let  $m = \chi(X)$  and  $k = |A|$ . For each  $x \in X$  let  $\mathcal{B}(x)$  be a base for  $X$  at the point  $x$  such that  $|\mathcal{B}(x)| \leq m$ . If  $x \in \text{cl}_{\tau\theta}(A)$ , choose a point in  $\text{cl}_{\tau}(U) \cap A$  for every  $U \in \mathcal{B}(x)$  and let  $B_x$  be the set so obtained. Clearly,  $x \in \text{cl}_{\tau\theta}(B_x)$  and  $|B_x| \leq m$ . Let  $\mathcal{G}_x = \{\text{cl}_{\tau}(U) \cap B_x : U \in \mathcal{B}(x)\}$ . For every  $U \in \mathcal{B}_x$  we have  $x \in \text{cl}_{\tau\theta}(\text{cl}_{\tau}(U) \cap B_x)$ , in fact, if  $V \in \mathcal{B}(x)$  let  $W \in \mathcal{B}(x)$  such that  $W \subset V \cap U$ , then

$$\emptyset \neq \text{cl}_{\tau}(W) \cap B_x \subset \text{cl}_{\tau}(V \cap U) \cap B_x \subset \text{cl}_{\tau}(V) \cap (\text{cl}_{\tau}(U) \cap B_x).$$

Since  $X$  is functionally Hausdorff, then  $\bigcap \{\text{cl}_{\tau\theta}(\text{cl}_{\tau}(U) \cap B_x) : U \in \mathcal{B}(x)\} = \{x\}$ , in fact let  $y \neq x$ , then there exist open sets  $G$  and  $H$  such that  $x \in G$ ,  $y \in H$  and  $\text{cl}_{\tau}(G) \cap \text{cl}_{\tau}(H) = \emptyset$ , now let  $U \in \mathcal{B}(x)$  such that  $U \subset G$ , then  $\text{cl}_{\tau}(H) \cap \text{cl}_{\tau}(U) = \emptyset$ , so  $y \notin \bigcap \{\text{cl}_{\tau\theta}(\text{cl}_{\tau}(U) : U \in \mathcal{B}(x))\}$ , and, a fortiori,  $y \notin \bigcap \{\text{cl}_{\tau\theta}(\text{cl}_{\tau}(U) \cap B_x) :$

$U \in \mathcal{B}(x)$ . So the map  $\psi : \text{cl}_{\tau\theta}(A) \rightarrow \mathcal{P}_m(\mathcal{P}_m(A))$  defined by  $\psi(x) = \mathcal{G}_x$  for every  $x \in \text{cl}_{\tau\theta}(A)$ , is one to one. Since  $|\mathcal{P}_m(\mathcal{P}_m(A))| \leq (k^m)^m = k^m$ , then  $|\text{cl}_{\tau\theta}(A)| \leq k^m = |A|^{\chi(X)}$ . Let  $A_0 = A$  and, by transfinite induction, define for every  $\alpha < m^+$  sets  $A_\alpha$  such that  $A_\alpha = \text{cl}_{\tau\theta}(\bigcup\{A_\beta : \beta < \alpha\})$ . Clearly  $\bigcup\{A_\alpha : \alpha < m^+\} \subset [A]_{\tau\theta}$ . Now let  $x \in \text{cl}_{\tau\theta}(\bigcup\{A_\alpha : \alpha < m^+\})$ , for each  $V \in \mathcal{B}(x)$  choose a point in  $\text{cl}_\tau(V) \cap (\bigcup\{A_\alpha : \alpha < m^+\})$  and let  $B$  be the set so obtained, obviously  $B \in \mathcal{P}_m(\bigcup\{A_\alpha : \alpha < m^+\})$  and  $x \in \text{cl}_{\tau\theta}(B)$ . Since  $m^+$  is regular, there is an ordinal  $\alpha < m^+$  such that  $B \subset A_\alpha$ , so

$$x \in \text{cl}_{\tau\theta}(B) \subset \text{cl}_{\tau\theta}(A_\alpha) \subset A_{\alpha+1} \subset \bigcup\{A_\alpha : \alpha < m^+\},$$

therefore  $\bigcup\{A_\alpha : \alpha < m^+\}$  is  $\tau\theta$ -closed. Hence  $[A]_{\tau\theta} = \bigcup\{A_\alpha : \alpha < m^+\}$ . It remains to show that  $|A_\alpha| \leq k^m$  for each  $\alpha < m^+$  (this is equivalent to  $|\bigcup\{A_\alpha : \alpha < m^+\}| \leq k^m$ ). Suppose there is an ordinal  $\alpha < m^+$  such that  $|A_\alpha| > k^m$  and let  $\gamma = \min\{\alpha : |A_\alpha| > k^m\}$ . Since  $|A_\alpha| \leq k^m$  for every  $\beta < \gamma$ , we have  $|\bigcup\{A_\beta : \beta < \gamma\}| \leq k^m$ . Now  $A_\gamma = \text{cl}_{\tau\theta}(\bigcup\{A_\beta : \beta < \gamma\})$ , hence

$$|A_\gamma| = |\text{cl}_{\tau\theta}(\bigcup\{A_\beta : \beta < \gamma\})| \leq |\bigcup\{A_\beta : \beta < \gamma\}|^{\chi(X)} \leq (k^m)^m = k^m,$$

a contradiction. □

**Definition 7.** Let  $X$  be a topological space. The  $w$ -compactness degree of  $X$ , denoted by  $wcd(X)$ , is defined as the smallest infinite cardinal number  $k$  with the property that for every open cover  $\mathcal{U}$  of  $X$  there is a subcollection  $\mathcal{V} \in \mathcal{P}_k(\mathcal{U})$  for which  $X = \bigcup\{\text{cl}_\tau(V) : V \in \mathcal{V}\}$ .

For every space  $X$  we have  $wcd(X) \leq L(X)$  and this inequality can be proper.

**Example 8.** Let  $X$  be any infinite  $T_3$ -space such that every continuous real valued function defined on  $X$  is constant. Clearly  $wcd(X) = \aleph_0 < L(X)$ .

**Example 9.** For each  $\alpha < \omega_1$  let  $I(\alpha) = \{\alpha\} \times$  an open interval in the real line. Set  $X = \omega_1 \cup \bigcup\{I(\alpha) : \alpha < \omega_1\}$  and for  $x, y \in X$  define  $x < y$  if (i)  $x, y \in \omega_1$  and  $x < y$  in  $\omega_1$ , or (ii)  $x \in \omega_1, y \in I(\beta)$  and  $x \leq \beta$  in  $\omega_1$ , or (iii)  $x \in I(\gamma), y \in \omega_1$  and  $\gamma < y$  in  $\omega_1$ , or (iv)  $x \in I(\alpha), y \in I(\beta)$  and  $\alpha < \beta$  in  $\omega_1$ , or (v)  $x, y \in I(\alpha)$  and  $x < y$  in  $I(\alpha)$ . Let  $\sigma$  be the order topology on  $X$ . Let  $Y = X \cup \{\omega_1\}$ , define  $x < \omega_1$  for every  $x \in X$  and let  $\varrho$  be the order topology on  $Y$ . If  $\tau$  is the topology on  $Y$  generated by  $\varrho \cup \{Y - L : L \text{ is the set of limit ordinals in } Y - \{\omega_1\}\}$ , then  $(Y, \tau)$  is a functionally Hausdorff  $H$ -closed space which fails to be Lindelöf [7], so  $wcd(Y) = \aleph_0 < L(Y)$ .

**Theorem 10.** If  $X$  is a functionally Hausdorff space, then  $|X| \leq 2^{\chi(X)wcd(X)}$ .

PROOF: Let  $m = \chi(X)wcd(X)$  and for every  $x \in X$  let  $\mathcal{B}(x)$  be a base for  $X$  at the point  $x$  such that  $|\mathcal{B}(x)| \leq m$ . Construct a family  $\{C_\alpha : \alpha < m^+\}$  of subsets of  $X$  such that

- (1) for any  $\alpha < m^+$   $C_\alpha$  is  $\tau\theta$ -closed;
- (2) for any  $\alpha < m^+$   $|C_\alpha| \leq 2^m$ ;
- (3) if  $\alpha < \beta < m^+$ , then  $C_\alpha \subset C_\beta$ ;
- (4) for any  $\alpha < m^+$ , if  $\mathcal{U} \subset \bigcup\{\mathcal{B}(x) : x \in \bigcup\{C_\beta : \beta < \alpha\}\}$ ,  $|\mathcal{U}| \leq m$  and  $X - \bigcup\{\text{cl}_\tau(U) : U \in \mathcal{U}\} \neq \emptyset$ , then  $C_\alpha - \bigcup\{\text{cl}_\tau(U) : U \in \mathcal{U}\} \neq \emptyset$ .

The construction is done by transfinite induction. Let  $p \in X$  and  $C_0 = \{p\}$ . Let  $0 < \alpha < m^+$  and assume that  $C_\beta$  has been constructed for every  $\beta < \alpha$ . Let  $\mathcal{B}_\alpha = \bigcup\{\mathcal{B}(x) : x \in \bigcup\{C_\beta : \beta < \alpha\}\}$ , clearly  $|\mathcal{B}_\alpha| \leq 2^m$ . For any  $\mathcal{U} \subset \mathcal{B}_\alpha$  such that  $|\mathcal{U}| \leq m$  and  $X - \bigcup\{\text{cl}_\tau(U) : U \in \mathcal{U}\} \neq \emptyset$ , choose a point in  $X - \bigcup\{\text{cl}_\tau(U) : U \in \mathcal{U}\}$  and let  $A$  be the set so obtained, obviously  $|A| \leq 2^m$ . Let  $C_\alpha = [A \cup (\bigcup\{C_\beta : \beta < \alpha\})]_{\tau\theta}$ ,  $C_\alpha$  satisfies (1), (3), (4) and, by Theorem 6, also (2). The set  $C = \bigcup\{C_\alpha : \alpha < m^+\}$  is  $\tau\theta$ -closed, in fact let  $x \in \text{cl}_{\tau\theta}(C)$ , for every  $V \in \mathcal{B}(x)$  choose a point in  $\text{cl}_\tau(V) \cap C$  and let  $K$  be the set so obtained, clearly  $|K| \leq m$ , therefore there exists an  $\alpha < m^+$  such that  $K \subset C_\alpha$ , then  $x \in \text{cl}_{\tau\theta}(K) \subset \text{cl}_{\tau\theta}(C_\alpha) = C_\alpha \subset C$ . Obviously  $|C| \leq 2^m$ , so to complete the proof it suffices to show that  $C = X$ . Let us suppose that  $y \in X - C$ , since  $X$  is functionally Hausdorff, then for any  $x \in C$  there is a  $U_x \in \mathcal{B}(x)$  such that  $y \notin \text{cl}_\tau(U_x)$ ; for every  $x \in X - C$  let  $U_x \in \mathcal{B}(x)$  such that  $\text{cl}_\tau(U_x) \cap C = \emptyset$  ( $C$  is  $\tau\theta$ -closed).  $\{U_x\}_{x \in X}$  is an open cover of  $X$ , since  $wcd(X) \leq m$  there is a  $B \subset X$  such that  $|B| \leq m$  and  $X = \bigcup\{\text{cl}_\tau(U_x) : x \in B\}$ , clearly  $C \subset \bigcup\{\text{cl}_\tau(U_x) : x \in B \cap C\}$ . Since  $|B \cap C| \leq m$ , there is an  $\alpha < m^+$  such that  $B \cap C \subset C_\alpha$ . Let  $\mathcal{U} = \{U_x : x \in B \cap C\}$ ,  $\mathcal{U} \subset \bigcup\{\mathcal{B}(x) : x \in \bigcup\{C_\beta : \beta < \alpha + 1\}\}$ ,  $|\mathcal{U}| \leq m$ ,  $y \in X - \bigcup\{\text{cl}_\tau(U_x) : U_x \in \mathcal{U}\}$  and  $C_{\alpha+1} - \bigcup\{\text{cl}_\tau(U_x) : U_x \in \mathcal{U}\} = \emptyset$ , a contradiction. Hence  $C = X$  and the proof is complete.  $\square$

**Remark 11.** Let  $X$  be a functionally Hausdorff space and let  $wX$  be the completely regular space which has the same points and continuous real valued functions as those of  $X$ . Clearly  $L(wX) \leq wcd(X)$  for every functionally Hausdorff space  $X$ . On the other hand, there exist functionally Hausdorff spaces  $X$  such that  $\chi(X) < \chi(wX)$  (see e.g. [9, Example 36]). I do not know if  $\chi(wX)L(wX) \leq \chi(X)wcd(X)$  for every functionally Hausdorff space  $X$ ; if this is the case, then Theorem 10 is a consequence of the Arkhangel'skii's inequality quoted at the beginning.

#### REFERENCES

- [1] Arkhangel'skii A.V., *The power of bicompacta with the first axiom of countability*, Soviet Math. Dokl. **10** (1969), 951–955.
- [2] Bella A., Cammaroto F., *On the cardinality of Urysohn spaces*, Canad. Math. Bull. **31** (2) (1988), 153–158.
- [3] Engelking R., *General Topology. Revised and completed edition*, Sigma Series in Pure Mathematics 6, Heldermann Verlag, Berlin, 1989.
- [4] Hodel R., *Cardinal Functions I*, in Handbook of Set-Theoretic Topology (K. Kunen and J.E. Vaughan, eds.), Elsevier Science Publishers, B.V., North Holland, 1984, pp. 1–61.
- [5] Ishii T., *On the Tychonoff functor and  $w$ -compactness*, Topology Appl. **11** (1980), 173–187.
- [6] Pol R., *Short proofs of two theorems on cardinality of topological spaces*, Bull. Acad. Polon. Sci. Ser., Math. Astr. Phys. **22** (1974), 1245–1249.

- [7] Stephenson R.M., Jr., *Spaces for which the Stone-Weierstrass theorem holds*, Trans. Amer. Math. Soc. **133** (1968), 537–546.
- [8] ———, *Product spaces for which the Stone-Weierstrass theorem holds*, Proc. Amer. Math. Soc. **21** (1969), 284–288.
- [9] ———, *Pseudocompact and Stone-Weierstrass product spaces*, Pacific J. Math. **99** (1) (1982), 159–174.

DIPARTIMENTO DI MATEMATICA PURA ED APPLICATA, UNIVERSITÀ, 67100 L'AQUILA, ITALY

(Received July 27, 1993)