

On uniformly smoothing stochastic operators

WOJCIECH BARTOSZEK

Abstract. We show that a stochastic operator acting on the Banach lattice $L^1(m)$ of all m -integrable functions on (X, \mathcal{A}) is quasi-compact if and only if it is uniformly smoothing (see the definition below).

Keywords: stochastic operators, quasi-compact

Classification: Primary 47A35; Secondary 47B55, 47D07

Let (X, \mathcal{A}, m) be a σ -finite measure space. By \mathcal{D} we denote the set of all densities from $L^1(m)$, i.e. m integrable positive functions f such that $\int_X f dm = 1$.

A linear operator $P: L^1(m) \rightarrow L^1(m)$ is said to be stochastic if $P(\mathcal{D}) \subseteq \mathcal{D}$.

Stochastic operators have broad applications. The reader may find appropriate references in [LM]. Among other properties, usually the asymptotic behaviour of the iterates P^n is studied. In the middle of the eighties Komornik and Lasota introduced to the theory of stochastic operators the concept of smoothness. Namely, P is said to be smoothing if

- there exist a set $F \in \mathcal{A}$ of finite measure and
- (S) constants $0 < \eta < 1, 0 < \delta$ such that for any $f \in \mathcal{D}$
- and $E \in \mathcal{A}$ with $m(E) \leq \delta$ we have

$$\overline{\lim}_{n \rightarrow \infty} \int_{E \cup F^c} P^n f dm \leq \eta,$$

where F^c stands here and in the sequel for the complementation $X \setminus F$.

Smoothing stochastic operators have nice asymptotic properties. It is proved in [KL] that any smoothing stochastic operator P is asymptotically periodic i.e. there exist pairwise orthogonal densities g_1, \dots, g_r , positive functionals $\Lambda_1, \dots, \Lambda_r$ and a permutation α of the set $\{1, \dots, r\}$ such that $\lim_{n \rightarrow \infty} \| P^n f - \sum_{i=1}^r \Lambda_i(f) g_{\alpha^n(i)} \| = 0$ and $Pg_i = g_{\alpha(i)}$ $i = 1, 2, \dots, r$. In particular, for some constant d the sequence P^{nd} converges in the strong operator topology to $\sum_{i=1}^r \Lambda_i \otimes g_i$. The most general result in this direction was finally obtained by Komornik. Namely, it was

I thank the Foundation for Research Development for financial support

proved in [K] that any power bounded positive, and linear operator on $L^1(m)$ is asymptotically periodic.

In this note we discuss the uniform version of (S). Following [B3] (see Problem 3, page 57) we adapt here:

Definition. Let $0 < \eta < 1$. A stochastic operator P is said to be uniformly η -smoothing if there are $F \in \mathcal{A}$ with $m(F) < \infty$, and a constant $0 < \delta$ such that for some natural n_0

$$(US-\eta) \quad \sup_{f \in \mathcal{D}} \int_{E \cup F^c} P^{n_0} f \, dm \leq \eta$$

for all $E \in \mathcal{A}$ satisfying $m(E) \leq \delta$.

We will show that operators satisfying (US- η), are quasi-compact. Let us recall that an operator P is quasi-compact if $\|P^n - K\| < 1$ for some compact operator K and natural n . It is known (see for instance [B2]) that quasi-compact stochastic operators P are exactly those which satisfy $\|P^{nd} - \sum_{i=1}^r \Lambda_i \otimes g_i\| \xrightarrow[n \rightarrow \infty]{} 0$, for suitable d, r, Λ_i , and g_i . We will exploit here the characterization of quasi-compact operators obtained in [B1]. In particular we shall apply some of the results from the mentioned paper to Markov operators acting on the Banach lattice $C(\Delta)$ of all continuous functions on Δ , where Δ stands for the set of all linear and multiplicative functionals on $L^\infty(m)$ equipped with the *-weak topology, so Hausdorff and compact. We recall that a linear operator $T: C(\Delta) \rightarrow C(\Delta)$ is Markov if $T\mathbf{1} = \mathbf{1}$ and $Tf \geq 0$ for $f \geq 0$. The dual space to $C(\Delta)$ is identified with Radon, finite (signed) measures on Δ . The *-weak compact (nonempty) set of all probability measures μ on Δ such that $T^*\mu = \mu$ is denoted by $P_T(\Delta)$. Clearly the adjoint to P operator $T = P^*$ is markovian.

A linear operator R acting on a Banach space \mathcal{X} is said to be strongly ergodic if for all $x \in X$ the Cesaro means $n^{-1}(I + R + \dots + R^{n-1})x$ are convergent in the norm of \mathcal{X} . Sine's mean ergodic theorem (see [S]) provides necessary and sufficient conditions for strong ergodicity. Namely, it holds if and only if R -invariant vectors separate R^* -invariant ones. It is easy to verify that R^* -invariant vectors always separate R -invariant ones. In [B1] it is proved that a Markov operator T on $C(\Delta)$ is quasi-compact if T^* is strongly ergodic and the topological support $S(\mu)$ of any μ from $P_T(\Delta)$ is non-meager. Finally we notice that the quasi-compactness of P is equivalent to the quasi-compactness of its adjoint P^* .

Theorem. *Let P be a stochastic operator on $L^1(m)$. Then P is quasi-compact if and only if P is η -uniformly smoothing for some (for all) $0 < \eta < 1$.*

PROOF: Assume that P is η -uniformly smoothing with F, n_0, η, δ as in (US- η), and let $X = \bigcup_{j=1}^\infty X_j$ where X_j are pairwise disjoint with positive finite measure.

We assume that $X_1 = F$. Now let us define a probability measure

$$m_0 = \sum_{j=1}^{\infty} t_j m|_{X_j} \quad \text{where} \quad \sum_{j=1}^{\infty} t_j m(X_j) = 1, \quad \text{and} \quad t_j > 0.$$

Clearly m_0 and m are equivalent, so $L^\infty(m_0) = L^\infty(m)$. The measure m_0 may be transported on Δ by the Gelfand transform $\hat{\cdot}$. Then, for any $f \in L^\infty$ we have

$$\int_X f dm_0 = \int_\Delta \hat{f} d\hat{m}_0$$

where $\hat{f} \in C(\Delta)$ is the image of f by $\hat{\cdot}$. By \sim let us denote the inverse operation to $\hat{\cdot}$.

First we show that measures from $P_T(\Delta)$ are absolutely continuous with respect to \hat{m}_0 . Since $T^*L^1(\hat{m}_0) \subseteq L^1(\hat{m}_0)$, it is sufficient to show that any $\hat{\nu} \in P_T(\Delta)$ has a nonzero absolutely continuous with respect to \hat{m}_0 component. If not, let us suppose that for some $\hat{\nu} \in P_T(\Delta)$ one has $\hat{\nu} \perp \hat{m}_0$. Then there exists a clopen set $\hat{U} \subseteq \Delta$ so that

$$(\star) \quad \hat{m}_0(\hat{U}) < t_1 \delta \quad \text{with} \quad \hat{\nu}(\hat{U}) = 1.$$

Let $\hat{f} \in C(\Delta)$ be such that $\int \hat{f} d\hat{m}_0 = 1$ and $T^{*n_0}(\widehat{f m_0})(\hat{U}) > \frac{1}{2} + \frac{\eta}{2}$. We get $\int_U P^{n_0} \frac{d(\widehat{f m_0})}{dm} \sim dm > \frac{1}{2} + \frac{\eta}{2} > \eta$. This implies $m(U \cap F) > \delta$, so $m_0(U \cap F) > t_1 \delta$, and finally contradicting (\star) we get $\hat{m}_0(\hat{U}) \geq \hat{m}_0(\hat{U} \cap \hat{F}) > t_1 \delta$. Therefore $P_T(\Delta) \subseteq L^1(\hat{m}_0)$, which easily implies that the topological support of $\nu \in P_T(\Delta)$ is non-meager.

Applying Sine's mean ergodic from [S] we notice that the operator T is strongly ergodic. In particular, $A_n^* \nu = n^{-1}(I^* + T^* + \dots + T^{*(n-1)})\nu$ is $*$ -weak convergent. Since Δ has the Grothendieck property ($*$ -weak convergent sequences from $C(\Delta)^*$ are weakly convergent) thus $A_n^* \nu$ is weakly convergent. But weakly convergent Cesaro means are norm convergent. Therefore, T^* is strongly ergodic. Using results of [B1] we easily obtain quasi-compactness of $T = P^*$. By Theorem 2 from [B2], there is a natural d such that P^{*nd} is convergent in the operator norm to a finite dimensional projection. This is equivalent to the norm convergence of P^{nd} , and P is quasi-compact.

To prove the opposite let us assume that a stochastic operator P is quasi-compact. For some d we have $\lim_{n \rightarrow \infty} P^{nd} = \sum_{i=1}^r \Lambda_i \otimes g_i$, where $g_i \in \mathcal{D}$ are pairwise orthogonal (i.e. $g_i \cdot g_j = 0$ m a.e. for $i \neq j$) and $\Lambda_i(f) = \int f h_i dm$ where $\|h_i\|_\infty \leq 1$. For a given $0 < \eta < 1$ we choose a set $F \in \mathcal{A}$ of finite measure

and positive δ that if $m(E) < \delta$ then $\int_{E \cup F^c} \sum_{j=1}^r g_j dm < \frac{\eta}{2}$. If n is such that $\|P^{nd} - \sum_{j=1}^r \Lambda_j \otimes g_j\| < \frac{\eta}{2}$, then we have

$$\begin{aligned} \int_{E \cup F^c} P^{nd} f dm &= \int_{E \cup F^c} \left(P^n f - \sum_{j=1}^r \lambda_j(f) g_j \right) dm + \sum_{j=1}^r \lambda_j(f) \int_{E \cup F^c} g_j dm \\ &\leq \frac{\eta}{2} + \int_{E \cup F^c} \sum_{j=1}^r g_j dm \leq \eta \end{aligned}$$

where f is an arbitrary density. □

REFERENCES

- [B1] Bartoszek W., *On quasi-compactness and invariant measures of Markov operators on $C(X)$* , Bull. Acad. Polon. Sci. **34** (1986), 69–72.
- [B2] ———, *Asymptotic periodicity of the iterates of positive contractions on Banach lattices*, Studia Math. **XCI** (1988), 179–188.
- [B3] ———, *On the asymptotic behaviour of iterates of positive linear operators*, Die Suid-Afrikaanse Wiskundevereniging Mededelings **25:1** (1993), 48–78.
- [K] Komorník J., *Asymptotic decomposition of smoothing positive operators*, Acta Universitatis Carolinae **30:2** (1989), 77–81.
- [KL] Komorník J., Lasota A., *Asymptotic decomposition of Markov operators*, Bull. Acad. Polon. Sci. **35** no. 5–6 (1987), 321–327.
- [LM] Lasota A., Mackey M.C., *Probabilistic Properties of Deterministic Systems*, Cambridge University Press, Cambridge, 1985.
- [S] Sine R., *A mean ergodic theorem*, Proc. Amer. Math. Soc. **24** (1970), 438–439.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH AFRICA, P.O. BOX 392,
PRETORIA 0001, SOUTH AFRICA

E-mail: bartowk@risc5.unisa.ac.za

(Received August 29, 1994)