

## On the product of a compact space with an hereditarily absolutely countably compact space

MADDALENA BONANZINGA

*Abstract.* We show that the product of a compact, sequential  $T_2$  space with an hereditarily absolutely countably compact  $T_3$  space is hereditarily absolutely countably compact, and further that the product of a compact  $T_2$  space of countable tightness with an hereditarily absolutely countably compact  $\omega$ -bounded  $T_3$  space is hereditarily absolutely countably compact.

*Keywords:* compact, countably compact, absolutely countably compact, hereditarily absolutely countably compact,  $\omega$ -bounded, countable tightness, sequential space

*Classification:* 54D20, 54B10, 54D55

### Introduction and preliminary

Recently Matveev in [Mat1], [Mat2] introduced a new property called absolute countable compactness (acc) which is stronger than countable compactness. He also introduced the related property hereditarily absolutely countably compact (hacc). Matveev ([Mat1]) proved that the acc and hacc properties are not necessarily preserved by products with compact space (see [Mat1, Example 2.2 and Remark 5.3]). Matveev proved however that if  $Y$  is compact and first countable, and  $X$  is an acc or hacc  $T_2$  space, then  $X \times Y$  is acc or hacc (see [Mat1, Theorem 2.3 or 5.4]), and he raised the following two questions

**Question 1** [Mat1]. *Is  $X \times Y$  acc provided  $Y$  is a compact space with countable tightness and  $X$  is an acc space?*

**Question 2** [Mat1]. *Is  $X \times Y$  hacc provided  $Y$  is a compact space with countable tightness and  $X$  is an hacc space?*

Vaughan ([Vau2]) proved that the product of a compact sequential  $T_2$  space with an acc  $T_3$  space is acc. With the previous result and the following well known Balogh's theorem, Vaughan gave an affirmative answer to Question 1 in models of the proper forcing axiom [PFA] (Question 1 remains open in ZFC).

---

This research was supported by a grant from the C.N.R. (G.N.S.A.G.A.) and M.U.R.S.T. through "Fondi 40 %" Italy

**Theorem 1** [Bal]. [PFA] *Every compact Hausdorff space of countable tightness is sequential.*

In this paper we augment Vaughan's proof ([Vau2]) to show that the product of a compact sequential  $T_2$  space with an hacc  $T_3$  space is hacc. Then, with this result and Balogh's theorem, we can give an affirmative answer to Question 2 provided we assume the proper forcing axiom (Question 2 remains open in ZFC).

We also consider further conditions under which the product of a compact space with an hacc space is hacc. Vaughan ([Vau2]) proved that the product of a compact  $T_2$  space of countable tightness with an acc  $\omega$ -bounded  $T_3$  space is acc and that the product of a compact  $T_2$  space of countable density-tightness (defined below) with an acc  $T_3$  space of countable density-tightness is acc. In this paper we consider "analogs" of the previous results for hacc spaces, that is we prove that the product of a compact  $T_2$  space of countable tightness with an hacc  $\omega$ -bounded  $T_3$  space is hacc and that the product of a compact  $T_2$  space of countable tightness with an acc  $T_3$  space of countable tightness is hacc. Further, we prove that the product of a compact  $T_2$  space of countable density-tightness with an hacc  $T_3$  space of countable density-tightness need not be hacc.

Concerning Question 2, we note that if in some model of set theory there is a counterexample  $X \times Y$  to Question 2, then  $Y$  must be a compact non-sequential space with countable tightness, and  $X$  must be an hacc space which is not  $\omega$ -bounded and does not have countable tightness or, equivalently, there exists a closed subset of  $X$  which does not have countable density-tightness.

Recall the following definitions:

**Definition 1** [Eng]. *A space  $X$  is called countably compact provided every countable open cover of  $X$  has a finite subcover.*

Note that a characterization of countable compactness (see [Eng, 3.12.22(d)]) states that a  $T_2$  space  $X$  is countable compact iff for every open cover  $\mathcal{U}$  of  $X$  there exists a finite set  $F \subset X$  such that  $St(F, \mathcal{U}) = \bigcup\{U \in \mathcal{U} : U \cap F \neq \emptyset\} = X$ .

**Definition 2** [Mat1]. *A space  $X$  is said to be absolutely countably compact (acc) provided for every open cover  $\mathcal{U}$  of  $X$  and every dense  $D \subset X$ , there exists a finite set  $F \subset D$  such that  $St(F, \mathcal{U}) = \bigcup\{U \in \mathcal{U} : U \cap F \neq \emptyset\} = X$ .*

Matveev ([Mat1]) noted that every compact space is acc and every acc  $T_2$  space is countably compact; he proved that every countably compact space with countable density-tightness (defined below) is acc. Further Vaughan ([Vau2]) proved that every countably compact, orthocompact space (defined below) is acc.

Matveev ([Mat1]) demonstrated that acc is not hereditary with respect to closed subsets, even with respect to regular closed subsets. Then he introduced the following definition.

**Definition 3** [Mat1]. *A space  $X$  is said to be hereditarily absolutely countably compact (hacc) if all closed subspaces of  $X$  are acc.*

We also recall some other definitions. A space  $X$  has countable tightness provided that whenever  $A \subset X$  and  $x \in \overline{A}$  there exists a countable  $C \subset A$  such that  $x \in \overline{C}$ . Using different terminology, Matveev introduced ([Mat1]) the notion of countable density-tightness.

**Definition 4** [Vau2]. *The density-tightness of a space  $X$ , denoted  $d_t(X)$ , is the smallest infinite cardinal  $\kappa$  such that for every dense subset  $D \subset X$  and every  $x \in X$  there exists a subset  $E$  of  $D$  such that  $|E| \leq \kappa$  and  $x \in \overline{E}$ .*

Further  $X$  is called *orthocompact* provided for every open cover  $\mathcal{U}$  there exists an open refinement  $\mathcal{V}$  such that for every  $\mathcal{V}' \subset \mathcal{V}$ , we have  $\bigcap\{V \in \mathcal{V}' : x \in V\}$  is open for each  $x \in X$ . A set  $A \subset X$  is called *sequentially closed* if and only if  $A$  contains all limits of all sequences from  $A$ ;  $X$  is called a *sequential space* provided every sequentially closed set is closed (every sequential space has countable tightness, see [Eng, 1.7.13(c)]).  $X$  is called  $\omega$ -*bounded* if every countable subset is contained in a compact set.

Further we will use the following standard notation: for a set  $D$ ,  $[D]^{<\omega}$  denotes the set of all finite subsets of  $D$  and  $[D]^\omega$  the set of all finite or countable subsets of  $D$ . If  $D$  is a subset of a topological space  $X$ , the  $\aleph_0$ -*closure* of  $D$  (see [Arh]) is the set  $[D]_{\aleph_0} = \bigcup\{\overline{M} : M \in [D]^\omega\}$ .

**1. The product of a compact sequential  $T_2$  space with an hacc  $T_3$  space is hacc.**

**Theorem 1.1.** *If  $Y$  is a compact sequential  $T_2$  space, and  $X$  is an hacc  $T_3$  space, then  $X \times Y$  is hacc.*

PROOF: We proceed similarly to the corresponding proof of Vaughan ([Vau2, Theorem 1.2]). By contradiction, suppose there exists a closed non acc subset  $F$  of  $X \times Y$ , i.e., there exist a closed  $F \subset X \times Y$ , an open cover  $\mathcal{U}$  of  $F$  and a dense subset  $D$  of  $F$  such that for all  $B \in [D]^{<\omega}$  we have that  $St(B, \mathcal{U}) \not\supset F$ . Proceeding as in [Vau2, Theorem 1.2], we conclude that the closed sets  $F_B = \pi_Y(F \setminus St(B, \mathcal{U}))$  form a filter base on  $Y$ , where  $\pi_Y$  is the projection on  $Y$ . Hence by compactness there exists

$$y \in \bigcap\{F_B : B \in [D]^\omega\}.$$

Since  $X \times \{y\}$  is homeomorphic to the hacc space  $X$ , and  $F$  is closed in  $X \times Y$ , then  $(X \times \{y\}) \cap F$  is acc.

As in [Vau2, Theorem 1.2], there exists an open set  $V \subset X$  such that  $(V \times \{y\}) \cap F \neq \emptyset$  and

$$(1) \quad (\overline{V} \times \{y\}) \cap [D]_{\aleph_0} = \emptyset.$$

Let  $Z = \pi_Y((\overline{V} \times Y) \cap [D]_{\aleph_0})$ . Again as in [Vau2, Theorem 1.2],  $Z$  is sequentially closed in  $Y$  and, since  $Y$  is sequential,  $Z$  is closed in  $Y$ . Note that  $\pi_Y((V \times Y) \cap D) \subset \pi_Y((\overline{V} \times Y) \cap [D]_{\aleph_0}) = Z \subset \pi_Y((\overline{V} \times Y) \cap F)$ . Since  $D$  is dense in  $F$  and

$(V \times Y) \cap F$  is a nonempty open subset of  $F$ , we have that  $(V \times Y) \cap D$  is dense in  $(V \times Y) \cap F$ . As  $\pi_Y$  is a continuous mapping, we have that  $\overline{\pi_Y((V \times Y) \cap D)}$  is dense in  $\pi_Y((V \times Y) \cap F)$ , then  $\pi_Y((V \times Y) \cap F) \subseteq \pi_Y(\overline{(V \times Y) \cap D}) = \overline{Z} = Z$ . Since  $y \in \pi_Y((V \times \{y\}) \cap F)$ , we have that  $y \in Z = \pi_Y(\overline{(V \times Y) \cap D})$ . Then  $\overline{(V \times \{y\}) \cap D} \neq \emptyset$ , but this contradicts (1), and completes the proof.  $\square$

**Corollary 1.1.** [PFA] *If  $Y$  is a compact  $T_2$  space with countable tightness, and  $X$  is an hacc  $T_3$  space, then  $X \times Y$  is hacc.*

**2. The product of a compact  $T_2$  space of countable tightness with an hacc  $\omega$ -bounded  $T_3$  space is hacc.**

**Theorem 2.1.**  *$X \times Y$  is hacc provided  $Y$  is a compact  $T_2$  space of countable tightness and  $X$  is an hacc,  $\omega$ -bounded  $T_3$  space.*

PROOF: Also in this case, we proceed similarly to the corresponding proof of Vaughan ([Vau2, Theorem 1.4]). The beginning of the proof repeats the first part of the proof of Theorem 1.1, with the modification (as in [Vau2, Theorem 1.4]) that the sets  $F_B$  need not be closed. We get that there exists

$$y \in \{\overline{F_B} : B \in [D]^\omega\}$$

and an open set  $V \subset X$  such that  $(V \times \{y\}) \cap F \neq \emptyset$  and

$$(\overline{V \times \{y\}}) \cap [D]_{\aleph_0} = \emptyset.$$

Now we show that

$$y \in \overline{\pi_Y((V \times Y) \cap D)}.$$

Since  $D$  is dense in  $F$ , we have that  $\overline{\pi_Y((V \times Y) \cap D)}$  is dense in  $\pi_Y((V \times Y) \cap F)$  and then  $\pi_Y((V \times Y) \cap F) \subset \overline{\pi_Y((V \times Y) \cap D)}$ . Hence, as  $(V \times \{y\}) \cap F \neq \emptyset$  and  $y \in \pi_Y((V \times \{y\}) \cap F)$ , we have that  $y \in \overline{\pi_Y((V \times Y) \cap D)}$ .

Proceeding as in [Vau2, Theorem 1.4], we obtain the desired conclusion.  $\square$

Vaughan ([Vau2]) obtained that the product of a compact  $T_2$  space of countable tightness with a countably compact  $GO$ -space (generalized ordered spaces, i.e., spaces which are subspaces of linearly ordered topological spaces; see [FL]) is acc. By Theorem 2.1, we have the following result concerning hacc (and then acc).

**Corollary 2.1.**  *$X \times Y$  is hacc provided  $Y$  is a compact  $T_2$  space of countable tightness and  $X$  is a countably compact  $GO$ -space.*

PROOF: Since every countably compact  $GO$ -space is  $\omega$ -bounded ([GFW, Theorem 3]), and every  $GO$ -space is orthocompact ([FL, 5.23]) we have that  $X$  is an  $\omega$ -bounded, orthocompact space; further, as every countably compact, orthocompact space is acc ([Vau2]) and orthocompactness and countable compactness are hereditary with respect to closed subsets, we have that  $X$  is hacc. Then, by Theorem 2.1,  $X \times Y$  is hacc.  $\square$

**3. The product of a compact  $T_2$  space of countable tightness with an acc  $T_3$  space of countable tightness is hacc.**

We have the following result:

**Proposition 3.1.**  *$X \times Y$  is hacc provided  $Y$  is a compact  $T_2$  space with countable tightness and  $X$  is an acc  $T_3$  space with countable tightness.*

PROOF: It is well-known that  $X \times Y$  is countably compact. By Malyhin's Theorem (see [Mal]),  $X \times Y$  has countable tightness. Both properties, countable compactness and countable tightness, are hereditary with respect to closed subsets. Thus by Matveev's Theorem (see [Mat1, Theorem 1.8]),  $X \times Y$  is hacc. □

While trying to answer to the following question: *do there exist a space  $X$  and a compact space  $Y$  such that  $X$  is hacc,  $X \times Y$  is acc, but  $X \times Y$  is not hacc?*, we obtained the following example which answer this question and also demonstrates why some assumptions in Theorems 1.1 and 2.1 cannot be weakened.

**Example 3.1.** *The product of an hacc Tychonoff space of countable density-tightness with a compact  $T_2$  space of countable density-tightness need not be hacc.*

Consider the Franklin-Rajagopalan spaces (see [Eng, 3.12.17(d)])  $X = T \cup Z$  and  $Y = T \cup Z'$ , where  $T$  is homomorphic to  $\omega$  with discrete topology,  $Z$  is homomorphic to the ordinal  $\mathfrak{t}$  (see [vD], [Vau1] or [Eng, 3.12.17(d)] where it is denoted  $\delta$ ) with order topology and  $Z'$  is homomorphic to  $\mathfrak{t}+1$  with order topology. We have that  $d_t(X) = \omega$  and  $d_t(Y) = \omega$  because both  $X$  and  $Y$  contain countable dense subsets of isolated points (so, of course,  $d_t(X \times Y) = \omega$  and then  $X \times Y$  is acc ([Mat1, Lemma 1.7])). Now we show that  $X$  is hacc. Let  $F$  be a closed subspace of  $X$ . Then  $F = F_T \cup F_Z$  where  $F_T = \overline{F \cap T}$  and  $F_Z = \overline{F \setminus F_T}$ .  $F_T$  has countable density-tightness because  $T$  is a countable set of isolated points; further, since  $Z$  is a linearly ordered topological space, then  $F_Z$  is a GO-space. Then,  $F_T$  and  $F_Z$  are acc ([Mat1, Lemma 1.7] and [Vau2, Corollary 1.7], respectively). Since  $T$  is open in  $X$  and  $F$  is closed in  $X$ , we have that  $F_T$  and  $F_Z$  are regular closed in  $F$ . So,  $F$  is written as union of regular closed acc spaces; then, from Proposition 4.3 in [Mat1], it follows that  $F$  is acc. So,  $X$  is hacc. Further  $Y$  is compact, but  $X \times Y$  is not hacc because it contains a closed copy of  $\mathfrak{t} \times (\mathfrak{t} + 1)$  that is not acc (see [Bon, Theorem 1.2]). It is worth mentioning that under the assumption  $\mathfrak{t} = \omega_1$ , the space  $X$  is even first-countable.

REFERENCES

[Arh] Arhangel'skii A.V., *On bicomacta hereditarily satisfying the Souslin condition. Tightness and free sequences*, Soviet Math. Dokl. **12** (1971), 1253–1257.  
 [Bal] Balogh Z., *On compact space of countable tightness*, Proc. Amer. Math. Soc. **105** (1989), 756–764.  
 [Bon] Bonanzinga M., *Preservation and reflection of properties acc and hacc*, Comment. Math. Univ. Carolinae **37,1** (1996), 147–153.

- [vD] van Douwen E.K., *The integers and topology*, Handbook Set-theoretic Topology, K. Kunen and J.E. Vaughan Eds., Elsevier Sci. Publ., 1984, pp. 111–167.
- [Eng] Engelking R., *General Topology*, Warszawa, 1977.
- [FL] Fletcher P., Lindgren W.F., *Quasi-uniform spaces*, Marcel Dekker, New York, 1982.
- [GFW] Gulden S.L., Fleischman W.F., Weston J.H., *Linearly ordered topological spaces*, Proc. Amer. Math. Soc. **24** (1970), 197–203.
- [Mal] Malyhin V.I., *On the tightness and the Suslin number of  $\exp X$  and of a product of spaces*, Soviet Math. Dokl. **13** (1972), 496–499.
- [Mat1] Matveev M.V., *Absolutely countably compact spaces*, Topology and Appl. **58** (1994), 81–92.
- [Mat2] Matveev M.V., *A countably compact topological group which is not absolutely countably compact*, Questions and Answers **11** (1993), 173–176.
- [Vau1] Vaughan J.E., *Small uncountable cardinals and topology*, Open Problems in Topology, J. van Mill, G.M. Reed Eds., North-Holland, Amsterdam, 1990, pp. 195–218.
- [Vau2] Vaughan J.E., *On the product of a compact space with an absolutely compact space*, Papers on General Topology and Applications, Annals of the New York Academy of Sciences **788** (1996), 203–208.

DIPARTIMENTO DI MATEMATICA, UNIVERSITA' DI MESSINA, CONTRADA PAPARDO, SALITA SPERONE, 98168 MESSINA, ITALY

(Received April 24, 1996, revised March 3, 1997)