

Fixed points for multifunctions on generalized metric spaces with applications to a multivalued Cauchy problem

ADRIAN PETRUȘEL

Abstract. The purpose of this paper is to prove an existence result for a multivalued Cauchy problem using a fixed point theorem for a multivalued contraction on a generalized complete metric space.

Keywords: generalized metric space, multivalued contraction, fixed points

Classification: 34A60, 47H10

1. Introduction

In 1958 W.A.J. Luxemburg, using a fixed point theorem for a single-valued contraction on a generalized metric space, proved the existence and the uniqueness of solution of the following Cauchy problem:

$$(1) \quad x'(t) = f(t, x(t)), \quad x(t_0) = x_0,$$

where t and x are real variables and f is a real function defined on the rectangle $|t - t_0| \leq a$, $|x - x_0| \leq b$, $a, b > 0$.

The purpose of this paper is to prove an existence result for a multivalued Cauchy problem using a fixed point theorem, for a multivalued contraction defined on a complete generalized metric space.

2. Preliminaries

The concept of a generalized metric space was introduced by Luxemburg and Jung as follows:

Definition 2.1 ([6], [9]). The pair (X, d) will be called a generalized metric space if X is an arbitrary nonempty set and d is a function $d : X \times X \rightarrow [0, \infty]$ which fulfills all the standard conditions for a metric.

In this paper, the generalized metric d is allowed to take the value $+\infty$ as well. In a generalized, just as in a metric space, we can define open and closed balls, convergence of sequences, completeness of the space, etc.

If (X, d) is a generalized metric space, $Y \subset X$, $x \in X$ and $\varepsilon > 0$ then:

$$\delta(Y) = \sup\{d(a, b) \mid a, b \in Y\},$$

$$D(Y, x) = \inf\{d(y, x) \mid y \in Y\},$$

$$B_X(x, \varepsilon) = \{y \in X \mid d(x, y) < \varepsilon\},$$

$$V(Y, \varepsilon) = \{x \in X \mid D(Y, x) \leq \varepsilon\},$$

$$\mathcal{P}(X) = \{Y \mid Y \subseteq X\},$$

$$P(X) = \{Y \in \mathcal{P}(X) \mid Y \neq \emptyset\},$$

$$P_{cl}(X) = \{Y \in P(X) \mid Y = \overline{Y}\},$$

$$P_{cp,cv}(X) = \{Y \in P(X) \mid Y \text{ compact and convex in } X\} \text{ (here } X \text{ is a generalized normed space),}$$

$$H(A, B) = \begin{cases} \inf\{\varepsilon > 0 \mid A \subset V(B, \varepsilon), B \subset V(A, \varepsilon)\}, & \text{if the infimum exists} \\ +\infty, & \text{otherwise.} \end{cases}$$

The pair $(P_{cl}(X), H)$ is a generalized metric space and H is called the generalized Hausdorff-Pompeiu distance induced by d .

Lemma 2.2 ([11]). *If (X, d) is complete generalized metric space then $(P_{cl}(X), H)$ is a complete generalized metric space.*

Definition 2.3 ([3]). Let (X, d) be a generalized metric space and $T : X \rightarrow P_{cl}(X)$ be a multivalued operator. Then, T is called an a -contraction if there exists a real number $a \in [0, 1[$ such that $x, y \in X, d(x, y) < \infty \Rightarrow H(T(x), T(y)) \leq ad(x, y)$.

Definition 2.4. Let (X, d) be a generalized metric space and $T : X \rightarrow P(X)$ a multivalued operator. Then $x^* \in X$ is called a fixed point for T if $x^* \in T(x^*)$. The set of all fixed points will be denoted by $\text{Fix}T$.

The concept of semi-continuous mappings was introduced in 1932 by Bouligand and Kuratowski.

We consider here the notion of an upper semicontinuous multivalued operator.

Definition 2.5 ([7]). Let X, Y be two metric spaces. A multivalued operator $T : X \rightarrow P(Y)$ is called upper semicontinuous at $x_0 \in X$ if and only if for any neighborhood U of $T(x_0)$, there exists a neighborhood V of x_0 such that for each $x \in V$ we have $T(x) \subset U$. T is said to be upper semicontinuous (u.s.c.) on X if it is u.s.c. at any point $x_0 \in X$.

Definition 2.6. Let (X, d) be a generalized metric space and $T : X \rightarrow P_{cl}(X)$ be a multivalued operator. A sequence $(x_n)_{n \in \mathbf{N}} \subset X$ is called the sequence of successive approximations of T if and only if $x_0 \in X$ and $x_n \in T(x_{n-1}), \forall n \in \mathbf{N}^*$.

The following result is well known in the field of set-valued analysis (see [1]).

Proposition 2.7. *Let $\Omega \subset \mathbf{R} \times \mathbf{R}^n$ be an open set, $(t_0, x_0) \in \Omega$ and $F : \Omega \rightarrow P_{cp}(\mathbf{R}^n)$ an u.s.c. multivalued operator.*

Then there exist $I = [t_0 - a, t_0 + a] \subset \mathbf{R}$ (where $a > 0$) and $M > 0$ such that:

- (i) $I \times B_{\mathbf{R}^n}(x_0, aM) \subset \Omega,$
- (ii) $\|F(t, x)\| \leq M$ on $I \times B_{\mathbf{R}^n}(x_0, aM).$

An important concept is that of integrably bounded multivalued operator.

Definition 2.8 ([4]). Let (S, \mathcal{A}, μ) be a complete σ -finite measure space and $(X, \|\cdot\|)$ be a separable Banach space. A multivalued operator $T : S \rightarrow P_{cl}(X)$ is said to be integrably bounded if and only if there is a function $r \in L^1(S)$ such that for all $v \in T(s)$ we have $\|v\| \leq r(s)$ a.e.

For $1 \leq p \leq \infty$ we define the set:

$$S_T^p := \{f \in L^p(\Omega, X) \mid f(s) \in T(s), \text{ a.e.}\},$$

i.e. S_T^p contains all selectors of T that belong to Lebesgue-Bochner space $L^p(\Omega, X)$.

It is easy to see that S_T^1 is a closed subset of $L^1(\Omega, X)$ and it is nonempty if and only if T is integrably bounded (see [2] and [4]).

Finally, the following theorem is a slight version of a result given in [10].

Theorem 2.9. *Let (X, d) be a complete generalized metric space and $T : X \rightarrow P_{cl}(X)$ be a multivalued α -contraction. We suppose that there is a sequence $(x_n)_{n \in \mathbf{N}} \subset X$ of successive approximations of T such that there exists an index $N(x_0) \in \mathbf{N}$ with the following property: $d(x_N, x_{N+l}) < \infty$, for all $l \in \mathbf{N}^*$. Then $\text{Fix } T \neq \emptyset$.*

3. Main result

Consider the following multivalued Cauchy problem

$$(2) \quad \begin{cases} x'(t) \in F(t, x(t)) \\ x(t_0) = x^0 \end{cases}$$

where $F : \Omega \subset \mathbf{R} \times \mathbf{R}^n \rightarrow P_{cp}(\mathbf{R}^n)$, with $\Omega = [t_0 - a, t_0 + a] \times \tilde{B}_{\mathbf{R}^n}(x^0, b)$, $(a, b > 0)$.

The main result of this note is the following existence theorem:

Theorem 3.1. *Consider the multivalued Cauchy problem (2). We suppose that:*

- (i) $F : \Omega \rightarrow P_{cp}(\mathbf{R}^n)$ is u.s.c. and integrably bounded,
- (ii) $|t - t_0|H(F(t, u), F(t, v)) \leq k\|u - v\|$, for every $(t, u), (t, v) \in \Omega$,
- (iii) $|t - t_0|^\beta H(F(t, u), F(t, v)) \leq A\|u - v\|^\alpha$, for every $(t, u), (t, v) \in \Omega$,
- (iv) $A, k > 0, 0 < \alpha < 1, \beta < \alpha$ and $k(1 - \alpha) < 1 - \beta$.

Then, the multivalued Cauchy problem (2) has a solution.

PROOF: From Proposition 2.7 it follows the existence of a real constant $M > 0$ such that $\|F(t, x)\| \leq M$ on Ω .

We denote by I the interval $I = [t_0 - h, t_0 + h]$, where $h = \min\{a, \frac{b}{M}\}$. We shall prove, by an application of Theorem 2.9, the existence of a solution of problem (2) on this interval I .

For this purpose we shall consider a space X with a generalized metric d , as follows:

$$X = \{\varphi \in C(I, \mathbf{R}^n) \mid \|\varphi(t) - x_0\| \leq b, \forall t \in I, \varphi(t_0) = x^0\}$$

$$d : X \times X \rightarrow \mathbf{R}_+ \cup \{+\infty\}$$

$$d(\varphi_1, \varphi_2) := \sup \left\{ \frac{\|\varphi_1(t) - \varphi_2(t)\|}{|t - t_0|^{pk}} \mid t \in I \right\},$$

where $p > 1, pk(1 - \alpha) < 1 - \beta$.

From [9] we have that (X, d) is a complete generalized metric space.

Finally, we choose the multivalued operator $T : X \multimap X$,

$$T(x) := \left\{ v \in X \mid v(t) \in x^0 + \int_{t_0}^t F(s, x(s)) ds \text{ a.e. } I \right\},$$

(where $\int_{t_0}^t F(s, x(s)) ds$ denotes the multivalued integral of Aumann).

It is easy to see that a function φ^* is a fixed point of T if and only if φ^* is a solution of problem (2).

We shall prove now that T satisfies all the hypotheses of Theorem 2.9.

(a) $T(x) \neq \emptyset$ for each $x \in X$.

Consider the multivalued operator F_x , given by $F_x(t) = F(t, x(t))$. By the Kuratowski-Ryll-Nardzewski selection theorem, F_x has a measurable selection $w(t) \in F_x(t)$, for all $t \in I$.

Define $v(t) = x^0 + \int_{t_0}^t w(s) ds, t \in I$. We obtain $v \in T(x)$ and so $T(x) \neq \emptyset$.

(b) $T(x)$ is closed for each $x \in X$.

Suppose (x_n) is a sequence in $T(x)$ which converges to $y \in X$. But $x_n(t) \in x^0 + \int_{t_0}^t F(t, x(t))$ a.e. and $x^0 + \int_{t_0}^t F(t, x(t))$ is closed (see [7]). Hence $y(t) \in x^0 + \int_{t_0}^t F(t, x(t))$ a.e.

(c) T is a multivalued contraction.

We shall prove that there exists $L \in (0, 1)$ such that for each $x, y \in X$ with $d(x, y) < \infty$ one obtains $H(T(x), T(y)) \leq Ld(x, y)$.

To see this, let $v_1 \in T(x)$. Then $v_1 \in X$ and $v_1(t) \in x^0 + \int_{t_0}^t F(s, x(s)) ds$, a.e. on I . It follows that there is a mapping $f_x \in S_{F(\cdot, x(\cdot))}^1$ such that $v_1(t) = x^0 +$

$\int_{t_0}^t f_x(s) ds$ a.e. on I . Since $H(F(t, x(t)), F(t, y(t))) \leq k \frac{\|x(t)-y(t)\|}{|t-t_0|}$ one obtains that there exists $w \in F(t, y(t))$ such that $\|f_x(t) - w\| \leq k \frac{\|x(t)-y(t)\|}{|t-t_0|}$ on I . Thus the multivalued operator G defined by $G(t) = F_y(t) \cap K(t)$, $t \in I$ (where $F_y(t) = F(t, y(t))$ and $K(t) = \{w \in F(t, y(t)) \mid \|f_x(t) - w\| \leq k \frac{\|x(t)-y(t)\|}{|t-t_0|}\}$) has nonempty values.

F_y and K are measurable and hence G is also measurable. Let f_y be a measurable selection for G (which exists by the Kuratowski-Ryll-Nardzewski selection theorem). Then $f_y(t) \in F(t, y(t))$ a.e. on I and $\|f_x(t) - f_y(t)\| \leq k \frac{\|x(t)-y(t)\|}{|t-t_0|}$ on I .

Define $v_2(t) = x^0 + \int_{t_0}^t f_y(s) ds$, $t \in I$. It follows that $v_2 \in T(y)$ and

$$\begin{aligned} \|v_1(t) - v_2(t)\| &= \|x^0 + \int_{t_0}^t f_x(s) ds - x^0 - \int_{t_0}^t f_y(s) ds\| \\ &\leq \int_{t_0}^t \|f_x(s) - f_y(s)\| ds \leq k \int_{t_0}^t \frac{\|x(s) - y(s)\|}{|s - t_0|} ds \\ &= k \int_{t_0}^t \frac{\|x(s) - y(s)\|}{|s - t_0|^{pk}} |s - t_0|^{pk-1} ds \leq kd(x, y) \int_{t_0}^t |s - t_0|^{pk-1} ds \\ &= kd(x, y) \frac{|t - t_0|^{pk}}{pk}. \end{aligned}$$

Finally, one obtains:

$$\frac{\|v_1(t) - v_2(t)\|}{|t - t_0|^{pk}} \leq \frac{1}{p}d(x, y) \text{ a.e.}$$

Hence $d(v_1, v_2) \leq \frac{1}{p}d(x, y)$.

From this and the analogous inequality obtained by interchanging the roles of x and y , we get

$$H(T(x), T(y)) \leq \frac{1}{p}d(x, y), \text{ for each } x, y \in X \text{ with } d(x, y) < \infty.$$

(d) T admits a sequence of successive approximations $(\varphi_n)_{n \in \mathbf{N}}$ with the property that there exists an index $N \in \mathbf{N}$ such that $d(\varphi_N, \varphi_{N+l}) < \infty$, for all $l \in \mathbf{N}^*$.

To see this, let $(\varphi_n)_{n \in \mathbf{N}}$ a sequence of successive approximations for T (where $\varphi_0 \in X$ is arbitrary). Let $\varphi_1 \in T(\varphi_0)$. It follows that there exists $f_0 \in L^1(I, \mathbf{R}^n)$, $f_0(s) \in F(s, \varphi_0(s))$ a.e. such that

$$\varphi_1(t) = x^0 + \int_{t_0}^t f_0(s) ds \text{ a.e.}$$

Let $\varphi_2 \in T(\varphi_1)$. By the definition of T , one obtains again that there exists $f_1 \in L^1(I, \mathbf{R}^n)$, $f_1(s) \in F(s, \varphi(s))$ a.e. such that

$$\varphi_2(t) = x_0 + \int_{t_0}^t f_1(s) ds \text{ a.e.}$$

By the boundedness of F we have

$$\|\varphi_2(t) - \varphi_1(t)\| = \left\| \int_{t_0}^t (f_1(s) - f_0(s)) ds \right\| \leq \int_{t_0}^t \|f_1(s) - f_0(s)\| ds \leq 2M|t - t_0|.$$

Since $f_1(s) \in F(s, \varphi(s))$ a.e. and F has compact values, we obtain that there exists $w \in F(s, \varphi_2(s))$, for each $s \in I$ such that

$$\|w - f_1(s)\| \leq H(F(s, \varphi_2(s)), F(s, \varphi_1(s))).$$

Consider the multivalued operator G defined by $G(s) = F_{\varphi_2}(s) \cap H^*(s)$ (where $F_{\varphi_2}(s) := F(s, \varphi_2(s))$ and

$$H^*(s) := \{w \in X \mid \|w - f_1(s)\| \leq H(F(s, \varphi_2(s)), F(s, \varphi_1(s))) \text{ a.e.}\}.$$

Clearly G is measurable and by the Kuratowski-Ryll-Nardzewski selection theorem it admits a measurable selection $f_2(s) \in G(s)$ a.e. on I . Thus $f_2(s) \in F(s, \varphi_2(s))$ a.e. and

$$\|f_2(s) - f_1(s)\| \leq H(F(s, \varphi_2(s)), F(s, \varphi_1(s))).$$

Let $\varphi_3(t) := x_0 + \int_{t_0}^t f_2(s) ds$. We have:

$$\begin{aligned} \|\varphi_3(t) - \varphi_2(t)\| &\leq \int_{t_0}^t \|f_2(s) - f_1(s)\| ds \leq \int_{t_0}^t H(F(s, \varphi_2(s)), F(s, \varphi_1(s))) ds \\ &\leq A \int_{t_0}^t \frac{\|\varphi_2(s) - \varphi_1(s)\|}{|s - t_0|^\beta} ds \leq A(2M)^\alpha \int_{t_0}^t |s - t_0|^{\alpha - \beta} ds \\ &= A(2M)^\alpha \frac{|t - t_0|^{1 + \alpha - \beta}}{1 + \alpha - \beta} \leq A(2M)^\alpha |t - t_0|^{1 + \alpha - \beta}. \end{aligned}$$

Generally

$$\begin{aligned} \|\varphi_{n+1}(t) - \varphi_n(t)\| &\leq A^{1+\alpha+\dots+\alpha^{n-2}} (2M)^{\alpha^{n-1}} |t - t_0|^{(1-\beta)(1+\dots+\alpha^{n-2})+\alpha^{n-1}} \\ &< B|t - t_0|^{(1-\beta)(1+\alpha+\dots+\alpha^{n-2})+\alpha^{n-1}}, \end{aligned}$$

where $B = A^{\frac{1}{1-\alpha}} \max\{2M, 1\}$.

In view of $pk(1 - \alpha) < 1 - \beta$ there exists an index $N \in \mathbf{N}$ such that $(1 - \beta)(1 + \alpha + \dots + \alpha^{n-2}) + \alpha^{n-1} > pk$, for each $n \geq N$. Hence for $n \geq N$, we have:

$$\frac{\|\varphi_{n+1}(t) - \varphi_n(t)\|}{|t - t_0|^{pk}} \leq B|t - t_0|^{\gamma_n},$$

where $\gamma_n = (1 - \beta)(1 + \dots + \alpha^{n-2}) + \alpha^{n-1} - pk > 0$.

This shows that $d(\varphi_{n+1}, \varphi_n) < \infty$, for all $n \geq N$, which completes the proof.

After these verifications, an application of Theorem 2.9 in the preceding section gives the desired conclusion. □

Remark 3.2. For $\beta = 0$, we get an existence result which is an improvement of Theorem 2 from [5].

REFERENCES

- [1] Aubin J.P., Cellina A., *Differential Inclusions*, Springer Verlag, Berlin, 1984.
- [2] Aumann R.J., *Integrals of set-valued functions*, J. Math. Anal. Appl. **12** (1965), 1–12.
- [3] Covitz H., Nadler S.B., Jr., *Multivalued contraction mappings in generalized metric spaces*, Israel J. Math. **8** (1970), 5–11.
- [4] Hiai F., Umegake H., *Integrals, conditional expectations and martingales of multivalued functions*, J. Multivariate Anal. **7** (1977), 149–182.
- [5] Himmelberg C.J., Van Vleck F.S., *Lipschitzian generalized differential equations*, Rend. Sem. Mat. Univ. Parma **48** (1973), 159–169.
- [6] Jung C.K., *On generalized complete metric space*, Bull. A.M.S. **75** (1969), 113–116.
- [7] Kisielewicz M., *Differential Inclusions and Optimal Control*, Kluwer Acad. Publ., Dordrecht, 1991.
- [8] Kuratowski K., Ryll-Nardzewski C., *A general theorem on selectors*, Bull. Polish Acad. Sci. **13** (1965), 397–403.
- [9] Luxemburg W.A.J., *On the convergence of successive approximations in the theory of ordinary differential equations, II*, Indag. Math. **20** (1958), 540–546.
- [10] Petruşel A., *On a theorem by Roman Wegrzyk*, Demonstratio Math. **29** (1996), 637–641.
- [11] Wegrzyk R., *Fixed point theorems for multivalued functions and their applications to functional equations*, Diss. Math. **201** (1982), 1–28.

“BABEŞ-BOLYAI” UNIVERSITY CLUJ-NAPOCA, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, STR. KOGĂLNICEANU 1, 3400 CLUJ-NAPOCA, ROMANIA

(Received December 12, 1996)