

The $\mathcal{L}_\nu^{(\rho)}$ - transformation on McBride’s spaces of generalized functions

D.I. CRUZ-BÁEZ, J. RODRÍGUEZ

Abstract. An integral transform denoted by $\mathcal{L}_\nu^{(\rho)}$ that generalizes the well-known Laplace and Meijer transformations, is studied in this paper on certain spaces of generalized functions introduced by A.C. McBride by employing the adjoint method.

Keywords: Krätzel integral transformation, L_p -spaces, distributions

Classification: 44A15, 46F12

1. Introduction

E. Krätzel [5] introduced a generalized Laplace transformation defined by

$$(1.1) \quad \mathcal{L}_\nu^{(\rho)}(f)(x) = \int_0^\infty \lambda_\nu^{(\rho)}(xy)f(y) dy, \quad x > 0,$$

where

$$(1.2) \quad \lambda_\nu^{(\rho)}(x) = \frac{(2\pi)^{(\rho-1)/2} \rho^{1/2}}{\Gamma(\nu + 1 - (1/\rho))} \left(\frac{x}{\rho}\right)^{\rho\nu} \int_1^\infty (t^\rho - 1)^{\nu-(1/\rho)} e^{-xt} dt, \quad x > 0$$

for $\rho \in \mathbf{N}$ and $\text{Re } \nu > -1 + 1/\rho$. He studied in a series of papers ([5], [6] and [7]) the main classical properties of $\mathcal{L}_\nu^{(\rho)}$. J.J. Betancor and J. Barrios ([1] and [2]) continued the investigations of E. Krätzel, and they established that $\lambda_\nu^{(\rho)}(z)$ is a solution of a differential equation of fractional order. In [11] the $\mathcal{L}_\nu^{(\rho)}$ -transform is investigated on certain spaces of distributions following Zemanian by means of the kernel method. We will consider the $\mathcal{L}_\nu^{(\rho)}$ transform on McBride’s spaces of test functions $\mathcal{F}_{p,\mu}$ and define it on their duals $\mathcal{F}'_{p,\mu}$ by means of the method of adjoints.

Throughout this paper $\rho \in \mathbf{R}$ and $\rho > 0$.

The asymptotic behaviour of $\lambda_\nu^{(\rho)}(x)$ can be found in [2]; for $x \rightarrow 0$ we have

$$(1.3) \quad \lambda_\nu^{(\rho)}(x) = \begin{cases} c_1 + o(1), & \text{Re } \nu > 0, \\ c_2 (x/\rho)^{\rho\nu} + o(1), & \text{Re } \nu = 0, \nu \neq 0, \\ c_3 \log(x/\rho) + o(1), & \nu = 0, \\ c_4 (x/\rho)^{\rho\nu} + o(1), & \text{Re } \nu < 0, \end{cases}$$

where c_1, c_2, c_3 and c_4 are suitable constants, and for $x \rightarrow \infty$,

$$(1.4) \quad \lambda_\nu^{(\rho)}(x) = O(e^{-x}).$$

The following property will be useful in the sequel ([5])

$$(1.5) \quad D\lambda_\nu^{(\rho)}(z) = - \left(\frac{z}{\rho}\right)^{\rho-1} \lambda_{\nu-1}^{(\rho)}(z).$$

Here, D denotes ordinary differentiation.

2. Krätzel transform on spaces $\mathcal{F}_{p,\mu}$ and $\mathcal{F}'_{p,\mu}$

A.C. McBride [9] defines $\mathcal{F}_{p,\mu}$ as follows, let $\mu \in \mathbf{C}$,

$$\mathcal{F}_{p,\mu} = \left\{ \varphi \in \mathcal{C}^\infty(\mathbf{R}^+) : x^k \frac{d^k}{dx^k}(x^{-\mu}\varphi(x)) \in L^p(\mathbf{R}^+), \forall k \in \mathbf{N} \right\},$$

where $1 \leq p < \infty$ and

$$\mathcal{F}_{\infty,\mu} = \left\{ \varphi \in \mathcal{C}^\infty(\mathbf{R}^+) : x^k \frac{d^k}{dx^k}(x^{-\mu}\varphi(x)) \rightarrow 0 \text{ as } x \rightarrow 0 \text{ and } x \rightarrow \infty, \forall k \in \mathbf{N} \right\},$$

where $p = \infty$. $\mathcal{F}_{p,\mu}$ is a complete countable multinormed space (Fréchet space) equipped with the topology generated by the family of seminorms in $\mathcal{F}_{p,\mu}$ given by

$$\gamma_k^{p,\mu}(\varphi) = \left\| x^k \frac{d^k}{dx^k}(x^{-\mu}\varphi) \right\|_p \quad (k \in \mathbf{N}; 1 \leq p \leq \infty, \mu \in \mathbf{C}).$$

In [10] we can see that the space $\mathcal{F}_{p,\mu}$ is closely connected with the Banach space $L_{p,\mu}$ of Lebesgue measurable functions $f(x)$ such that $\|f\|_{p,\mu} = \|x^{-\mu}f\|_p < \infty$. $\mathcal{F}'_{p,\mu}$ is the space of the continuous linear functionals on $\mathcal{F}_{p,\mu}$ equipped with the weak topology.

Next, we establish a series of results for finally to define the $\mathcal{L}_\nu^{(\rho)}$ -transformation by using the adjoint method.

Proposition 2.1. *Let $1 \leq p \leq \infty, \mu, \nu \in \mathbf{C}, \rho > 0, 1/p + 1/p' = 1$ and*

$$(2.1) \quad \operatorname{Re} \mu > -\frac{1}{p} - \min \{0, \rho \operatorname{Re} \nu\}.$$

Then $\mathcal{L}_\nu^{(\rho)}$ is a continuous linear mapping from $L_{p,\mu}$ into $L_{p,2/p-\mu-1}$ and from $\mathcal{F}_{p,\mu}$ into $\mathcal{F}_{p,2/p-\mu-1}$.

PROOF: By (1.3) and (1.4) the integral

$$\int_0^\infty x^{\operatorname{Re} \mu - \frac{1}{p}} \left| \lambda_\nu^{(\rho)}(x) \right| dx < \infty$$

converges provided that (21) is satisfied. Then Proposition 2.1 follows from [9, pp. 158–159, Theorem 8.1 and Corollary 8.2] and the proof concludes. \square

The Mellin transform $(\mathcal{M}\varphi)(s)$ of a suitable function $\varphi(x)$, $x > 0$, is defined by

$$(2.2) \quad (\mathcal{M}\varphi)(s) = \int_0^\infty x^{s-1}\varphi(x) dx.$$

Lemma 2.1. *Let $\rho > 0$, $\nu, s \in \mathbf{C}$ and*

$$(2.3) \quad \operatorname{Re} s + \min \{0, \rho \operatorname{Re} \nu\} > 0.$$

Then

$$(2.4) \quad \mathcal{M} \left\{ \lambda_\nu^{(\rho)}(x) \right\} (s) = (2\pi)^{\frac{\rho-1}{2}} \rho^{-1/2-\rho\nu} \frac{\Gamma(s + \rho\nu)\Gamma\left(\frac{s}{\rho}\right)}{\Gamma\left(\frac{s}{\rho} + \nu + 1 - \frac{1}{\rho}\right)}.$$

PROOF: By the asymptotic behaviour of $\lambda_\nu^{(\rho)}$ we can guarantee (2.3). By (2.2) and (1.2) we have after changing the order of integration (Fubini's theorem)

$$\begin{aligned} \mathcal{M} \left\{ \lambda_\nu^{(\rho)}(x) \right\} &= \int_0^\infty x^{s-1} \lambda_\nu^{(\rho)}(x) dx \\ &= \frac{(2\pi)^{\frac{\rho-1}{2}} \rho^{1/2-\rho\nu}}{\Gamma\left(\nu + 1 - \frac{1}{\rho}\right)} \int_0^\infty x^{s+\rho\nu-1} \int_1^\infty (t^\rho - 1)^{\nu-(1/\rho)} e^{-xt} dt dx \\ &= \frac{(2\pi)^{\frac{\rho-1}{2}} \rho^{1/2-\rho\nu}}{\Gamma\left(\nu + 1 - \frac{1}{\rho}\right)} \int_1^\infty (t^\rho - 1)^{\nu-(1/\rho)} \int_0^\infty x^{s+\rho\nu-1} e^{-xt} dx dt. \end{aligned}$$

The relation $\Gamma(z) = \int_0^\infty \tau^{z-1} e^{-\tau} d\tau$ ($\operatorname{Re} z > 0$) holds, hence

$$\begin{aligned} \mathcal{M} \left\{ \lambda_\nu^{(\rho)}(x) \right\} &= \frac{(2\pi)^{\frac{\rho-1}{2}} \rho^{1/2-\rho\nu} \Gamma(s + \rho\nu)}{\Gamma\left(\nu + 1 - \frac{1}{\rho}\right)} \int_1^\infty (t^\rho - 1)^{\nu-(1/\rho)} t^{-s-\rho\nu} dt \\ &= (2\pi)^{\frac{\rho-1}{2}} \rho^{-1/2-\rho\nu} \frac{\Gamma(s + \rho\nu)\Gamma\left(\frac{s}{\rho}\right)}{\Gamma\left(\frac{s}{\rho} + \nu + 1 - \frac{1}{\rho}\right)} \end{aligned}$$

and (2.4) is proved. \square

The Mellin transform \mathcal{M} for $\varphi \in \mathcal{F}_{p,\mu}$ is defined by

$$(2.5) \quad (\mathcal{M}\varphi)(s) = \int_0^\infty t^{s-1}\varphi(t) dt, \quad \operatorname{Re} s = 1/p - \operatorname{Re} \mu.$$

By [10, p. 531, Theorem 5.1], we have for $1 \leq p \leq 2$ and $\mu \in \mathbf{C}$ that \mathcal{M} is a continuous linear mapping from $\mathcal{F}_{p,\mu}$ into $L_{p'}(\mathbf{R}^+)$.

Proposition 2.2. *Let $1 \leq p \leq 2$, $\mu, \nu \in \mathbf{C}$, $\rho > 0$ and*

$$(2.6) \quad \operatorname{Re} \mu > -\frac{1}{p'} - \min \{0, \rho \operatorname{Re} \nu\}, \quad \operatorname{Re} s = \frac{1}{p} + \operatorname{Re} \mu.$$

Then for $\varphi \in \mathcal{F}_{p,\mu}$ we have

$$(2.7) \quad \mathcal{M} \left\{ \mathcal{L}_\nu^{(\rho)} \varphi \right\} (s) = (2\pi)^{\frac{p-1}{2}} \rho^{-1/2-\rho\nu} \frac{\Gamma(s + \rho\nu) \Gamma\left(\frac{s}{\rho}\right)}{\Gamma\left(\frac{s}{\rho} + \nu + 1 - \frac{1}{\rho}\right)} \mathcal{M}\varphi(1-s),$$

where $s = \frac{1}{p} - \operatorname{Re} \mu + it$.

PROOF: By Fubini's theorem and (2.4), for a sufficiently good function $\varphi \in \mathcal{C}_0^\infty(\mathbf{R}^+)$ we have

$$\begin{aligned} \mathcal{M} \left\{ \mathcal{L}_\nu^{(\rho)} \varphi \right\} (s) &= \int_0^\infty y^{s-1} \int_0^\infty \lambda_\nu^{(\rho)}(xy) \varphi(x) dx dy \\ &= \int_0^\infty \varphi(x) \int_0^\infty y^{s-1} \lambda_\nu^{(\rho)}(xy) dx dy. \end{aligned}$$

Making the change $xy = t$, we get

$$\begin{aligned} \mathcal{M} \left\{ \mathcal{L}_\nu^{(\rho)} \varphi \right\} (s) &= \int_0^\infty x^{-s} \varphi(x) dx \int_0^\infty t^{s-1} \lambda_\nu^{(\rho)}(t) dt \\ &= (2\pi)^{\frac{p-1}{2}} \rho^{-1/2-\rho\nu} \frac{\Gamma(s + \rho\nu) \Gamma\left(\frac{s}{\rho}\right)}{\Gamma\left(\frac{s}{\rho} + \nu + 1 - \frac{1}{\rho}\right)} \mathcal{M}\varphi(1-s) \end{aligned}$$

and (2.7) is proved for $\varphi \in \mathcal{C}_0^\infty(\mathbf{R}^+)$. By [9, p. 18, Corollary 2.7], $\mathcal{C}_0^\infty(\mathbf{R}^+)$ is dense in $\mathcal{F}_{p,\mu}$ and hence the relation (2.7) holds for $\varphi \in \mathcal{F}_{p,\mu}$. \square

Theorem 2.1. *Let $1 \leq p \leq \infty$, $\mu, \nu \in \mathbf{C}$, $\rho > 0$ and*

$$(2.8) \quad \operatorname{Re} \mu > -\frac{1}{p'} - \min \{0, \rho \operatorname{Re} \nu\}.$$

Then we have

$$(2.9) \quad \int_0^\infty \left(\mathcal{L}_\nu^{(\rho)} f \right) (x) \varphi(x) dx = \int_0^\infty f(x) \left(\mathcal{L}_\nu^{(\rho)} \varphi \right) (x) dx$$

for $\varphi \in \mathcal{F}_{p,\mu}$, $f \in \mathcal{F}_{p',\mu-1+2/p'}$ and $\varphi \in L_{p,\mu}$, $f \in L_{p',\mu-1+2/p'}$.

PROOF: By Proposition 2.1 $\mathcal{L}_\nu^{(\rho)} f$ and $\mathcal{L}_\nu^{(\rho)} \varphi$ exist for $f \in \mathcal{F}_{p',\mu-1+2/p'}$ and $\varphi \in \mathcal{F}_{p,\mu}$, respectively, provided that (2.8) is valid. In the beginning we will prove that the equality (2.9) is true for functions of $\mathcal{C}_0^\infty(\mathbf{R}^+)$.

If $f, \varphi \in \mathcal{C}_0^\infty(\mathbf{R}^+)$ we obtain

$$\begin{aligned} \int_0^\infty \left(\mathcal{L}_\nu^{(\rho)} f\right)(x)\varphi(x) dx &= \int_0^\infty \varphi(x) dx \int_0^\infty \lambda_\nu^{(\rho)}(yx)f(y) dy \\ &= \int_0^\infty f(y) dy \int_0^\infty \lambda_\nu^{(\rho)}(yx)\varphi(x) dx \\ &= \int_0^\infty f(y) \left(\mathcal{L}_\nu^{(\rho)} \varphi\right)(y) dy, \end{aligned}$$

since Fubini's theorem allows the exchange in the integration order.

Then, to prove (2.9) for $\varphi \in \mathcal{F}_{p,\mu}$, $f \in \mathcal{F}'_{p',\mu-1+2/p'}$ and $\varphi \in L_{p,\mu}$, $f \in L_{p',\mu-1+2/p'}$, it is sufficient to show that both sides of (2.9) are bounded linear functionals on $L_{p,\mu} \times L_{p',\mu-1+2/p'}$. Applying the Hölder inequality and the definition of the norm of $L_{p,\mu}$ we obtain

$$\begin{aligned} \int_0^\infty \left| \left(\mathcal{L}_\nu^{(\rho)} f\right)(x)\varphi(x) \right| dx &= \int_0^\infty |x^{-\mu}\varphi(x)| \left| x^\mu \left(\mathcal{L}_\nu^{(\rho)} f\right)(x) \right| dx \\ &\leq \left(\int_0^\infty |x^{-\mu}\varphi(x)|^p dx \right)^{1/p} \left(\int_0^\infty |x^\mu \left(\mathcal{L}_\nu^{(\rho)} f\right)(x)|^{p'} dx \right)^{1/p'} \\ &= \|\varphi\|_{p,\mu} \left\| \mathcal{L}_\nu^{(\rho)} f \right\|_{p',-\mu}. \end{aligned}$$

Moreover, by Proposition 2.1 with p replaced by p' and μ by $\mu - 1 + 2/p'$

$$\left\| \mathcal{L}_\nu^{(\rho)} f \right\|_{p',-\mu} \leq k \|f\|_{p',\mu-1+2/p'} \quad (k > 0)$$

and hence

$$\left| \int_0^\infty \left(\mathcal{L}_\nu^{(\rho)} f\right)(x)\varphi(x) dx \right| \leq k \|\varphi\|_{p,\mu} \|f\|_{p',\mu-1+2/p'}.$$

This shows that the left hand side of (2.9) is a bounded linear functional on $L_{p,\mu} \times L_{p',\mu-1+2/p'}$. The same result for the right hand side of (2.9) is proved similarly. This completes the proof of Theorem 2.1. \square

Theorem 2.1 allows us to define the generalized $\mathcal{L}_\nu^{(\rho)}$ f -transform on $\mathcal{F}'_{p,\mu}$ when $1 \leq p \leq \infty$, $\mu, \nu \in \mathbf{C}$ and $\rho > 0$, as follows. For every $f \in \mathcal{F}'_{p,\mu}$ the generalized $\mathcal{L}_\nu^{(\rho)}$ f -transform is defined through

$$(2.10) \quad \langle \mathcal{L}_\nu^{(\rho)} f, \varphi \rangle = \langle f, \mathcal{L}_\nu^{(\rho)} \varphi \rangle$$

with $\varphi \in \mathcal{F}_{p,2/p-\mu-1}$.

Then by Proposition 2.1 and (2.10) we arrive at the following result.

Proposition 2.3. *Let $1 \leq p \leq \infty$, $\mu \in \mathbf{C}$, $\nu \in \mathbf{C}$, and $\operatorname{Re} \mu < 1/p + \min\{0, \rho \operatorname{Re} \nu\}$. Then the operator $\mathcal{L}_\nu^{(\rho)}$ is a continuous linear mapping of $\mathcal{F}'_{p,\mu}$ into $\mathcal{F}'_{p,2/p-\mu-1}$.*

Next, we investigate compositions of the operator $\mathcal{L}_\nu^{(\rho)}$ with a differential operator on the spaces $\mathcal{F}_{p,\mu}$ and $\mathcal{F}'_{p,\mu}$.

Proposition 2.4. *Let $1 \leq p \leq \infty$, $\mu, \nu \in \mathbf{C}$, $\rho > 0$, $m \in \mathbf{N}$, $1/p + 1/p' = 1$ and*

$$(2.11) \quad \operatorname{Re} \mu > -1/p' - \min\{0, \rho \operatorname{Re} \nu\} + \rho m.$$

Then for $\varphi \in \mathcal{F}_{p,\mu}$

$$(2.12) \quad \left(\left(\frac{x}{\rho} \right)^{1-\rho} D \right)^m \mathcal{L}_\nu^{(\rho)} \{y^{-\rho m} \varphi(y)\} (x) = (-1)^m \mathcal{L}_{\nu-m}^{(\rho)} \{\varphi(y)\} (x).$$

PROOF: According to Proposition 2.1 and [9, p. 21, Theorem 2.11 and p. 26, Corollary 2.15] the left and right hand sides of (2.12) are continuous linear mapping from $\mathcal{F}_{p,\mu}$ into $\mathcal{F}_{p,2/p-\mu-1}$ provided that the condition (2.11) holds. Applying (1.2) and (1.5) we have

$$\begin{aligned} & \left(\left(\frac{x}{\rho} \right)^{1-\rho} D \right)^m \mathcal{L}_\nu^{(\rho)} \{y^{-\rho m} \varphi(y)\} (x) \\ &= \left(\left(\frac{x}{\rho} \right)^{1-\rho} D \right)^m \int_0^\infty \lambda_\nu^{(\rho)}(xy) \cdot y^{-\rho m} \varphi(y) dy \\ &= \int_0^\infty \left(\left(\frac{x}{\rho} \right)^{1-\rho} D \right)^m \{ \lambda_\nu^{(\rho)}(xy) \} y^{-\rho m} \varphi(y) dy. \end{aligned}$$

After the substitution $xt = z$, we obtain

$$\begin{aligned} & \left(\left(\frac{x}{\rho} \right)^{1-\rho} D \right)^m \mathcal{L}_\nu^{(\rho)} \{y^{-\rho m} \varphi(y)\} (x) \\ &= \int_0^\infty \left(\left(\frac{z}{\rho} \right)^{1-\rho} D \right)^m \{ \lambda_\nu^{(\rho)}(z) \} \varphi(z/x) \frac{dz}{x} \\ &= \int_0^\infty (-1)^m \{ \lambda_{\nu-m}^{(\rho)}(xy) \} \varphi(y) dy \\ &= (-1)^m \mathcal{L}_{\nu-m}^{(\rho)} \{\varphi(y)\} (x) \end{aligned}$$

and Proposition 2.4 is proved. □

Proposition 2.5. *Let $1 \leq p \leq \infty$, $\mu, \nu \in \mathbf{C}$, $\rho > 0$ and $m \in \mathbf{N}$. For every $f \in$ and $\mathcal{F}'_{p,\mu}$ we have*

$$(2.13) \quad x^{-\rho m} \mathcal{L}_\nu^{(\rho)} \left(D \left(\frac{x}{\rho} \right)^{1-\rho} \right)^m f(x) = \mathcal{L}_{\nu-m}^{(\rho)} f(x)$$

provided

$$(2.14) \quad \operatorname{Re} \mu < 1/p + \min \{0, \rho \operatorname{Re} \nu\} - \rho m.$$

PROOF: By the condition (2.14), [9, p. 32, Theorem 2.22] and Proposition 2.3, the left and right hand sides of are continuous linear mapping from $\mathcal{F}'_{p,\mu}$ into $\mathcal{F}'_{p,2/p-\mu-1}$.

By (2.10) and [9, p. 32, Theorem 2.22] we have

$$\begin{aligned} & \langle x^{-\rho m} \mathcal{L}_\nu^{(\rho)} \left(D \left(\frac{x}{\rho} \right)^{1-\rho} \right)^m f(x), \varphi(x) \rangle \\ &= \langle f, (-1)^m \left(\left(\frac{x}{\rho} \right)^{1-\rho} D \right)^m \mathcal{L}_\nu^{(\rho)} x^{-\rho m} \varphi(x) \rangle \end{aligned}$$

(and by Proposition 2.4, (2.10) and [9, p. 32, Theorem 2.22] we get)

$$= \langle f, \mathcal{L}_{\nu-m}^{(\rho)} \varphi(x) \rangle = \langle \mathcal{L}_{\nu-m}^{(\rho)} f, \varphi(x) \rangle,$$

which concludes the proof. \square

REFERENCES

- [1] Barrios J.A., Betancor J.J., *On a generalization of Laplace transform due to E. Krätzel*, J. Inst. Math. & Comp. Sci. (Math. Ser.) **3** (4) (1990), 273–291.
- [2] Barrios J.A., Betancor J.J., *A Real Inversion Formula for the Krätzel's Generalized Laplace Transform*, Extracta Mathematicae **6** (2) (1991), 55–57.
- [3] Erdelyi A., Magnus W., Oberhettinger F., Tricomi F., *Tables of Integral Transforms, Vol. II*, McGraw-Hill, New York, 1954.
- [4] Kilbas A.A., Bonilla B., Rivero M., Rodríguez J., Trujillo J., *Bessel type function and Bessel type integral transform on spaces $\mathcal{F}_{p,\mu}$ and $\mathcal{F}'_{p,\mu}$* , to appear.
- [5] Krätzel E., *Eine verallgemeinerung der Laplace- und Meijer-transformation*, Wiss. Z. Univ. Jena Math. Naturw. Reihe **5** (1965), 369–381.
- [6] Krätzel E., *Die faltung der L-transformation*, Wiss. Z. Univ. Jena Math. Naturw. Reihe **5** (1965), 383–390.
- [7] Krätzel E., *Integral transformations of Bessel-type*, Proceeding of International Conference on Generalized Functions and Operational Calculus, Varna, 1975, pp. 148–155.
- [8] Krätzel E., Menzer H., *Verallgemeinerte Hankel-Funktionen*, Pub. Math. Debrecen **18**, fasc. 1-4 (1973), 139–148.

- [9] McBride A.C., *Fractional Calculus and Integral Transforms of Generalized Functions*, Res. Notes Math., 31, Pitman Press, San Francisco, London, Melbourne, 1979.
- [10] McBride A.C., *Fractional powers of a class ordinary differential operators*, Proc. London Math. Soc., Ser. 3 **45** (1982), no. 3, 519–546.
- [11] Rao G.L.N., Debnath L., *A generalized Meijer transformation*, Int. J. Math. & Math. Sci. **8:2** (1985), 359–365.
- [12] Zemanian A.H., *A distributional K transformation*, Siam J. Appl. Math. **14** (1966), no. 6, 1350–1365.
- [13] Zemanian A.H., *Generalized Integral Transformation*, Interscience Publisher, New York, 1968.

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE LA LAGUNA, 38271 LA LAGUNA (TENERIFE), SPAIN

E-mail: dicruz@ull.es

joroguez@ull.es

(Received October 13, 1997, revised March 26, 1998)