

A retractible non-locally connected dendroid

ALEJANDRO ILLANES

Abstract. A retractible non-locally connected dendroid is constructed.

Keywords: continua, dendroid, retractible

Classification: 54F50, 54F15, 54C15

A *continuum* is a compact connected metric space. A continuum X is *retractible* if for every subcontinuum A of X there exists a retraction $r : X \rightarrow A$. Retractable continua were introduced by J.J. Charatonik in [1], where he posed the following problem:

Problem. Give a structural (internal) characterization of retractible continua.

In the same paper, this problem is partially solved by showing that a locally connected continuum is retractible if and only if it is hereditarily locally connected.

A different approach for attacking this problem is to add requirements to the retractions. A continuum X is said to be *d-retractible* (resp., *sd-retractible*, *m-retractible*, *c-retractible*, *o-retractible*), provided that for each subcontinuum A of X , there exists a deformation (resp. strong deformation, monotone, confluent, open) retraction from X onto A . In [2], G.R. Gordh and L. Lum proved that a continuum is *m-retractible* if and only if it is a dendrite. Recently, the author has shown the following results:

Theorem ([3]). *If X is a continuum, then the following assertions are equivalent:*

- (a) X is a dendrite,
- (b) X is *d-retractible* and,
- (c) X is *sd-retractible*.

Theorem ([4]). *If X is a pathwise connected *c-retractible* continuum, then X is hereditarily locally connected.*

Theorem ([4]). *If X is a pathwise connected continuum, then the following assertions are equivalent:*

- (a) X is *o-retractible* and,
- (b) X is homeomorphic to an interval or to a simple closed curve.

In [1] and [5], J.J. Charatonik and L. Lum, respectively, asked the following question:

Question. Does there exist an arcwise connected retractible continuum which is not locally connected?

In [5, p. 337], L. Lum mentioned that A. Lelek had a candidate for answering this question in the positive.

In this paper, we answer the question in the positive by constructing a non-locally connected retractible dendroid.

In a recent private communication with J.J. Charatonik, A. Lelek told him that his example had a similar construction as the example presented here and that his example was never written for publication.

Preliminary constructions

Given two points p and q in the Euclidean plane R^2 , denote by $\langle p, q \rangle$ the segment joining them, if $p \neq q$ and $\langle p, q \rangle = \{p\}$, if $p = q$. For a point $p = (x, y) \in R^2$, define $p' = (-x, y)$. Given a subset B in R^2 , define $B' = \{p' \in R^2 : p \in B\}$. The origin in R^2 is denoted by Θ . Let $\pi_1 : R^2 \rightarrow R$ be the projection on the first coordinate. We will define, inductively, a sequence A_0, \dots, A_n, \dots of subsets of R^2 such that, for each integer $n > 0$, A_n is a polygon joining Θ to a point $a_n = (u_n, v_n)$. Let $a_0 = \Theta$.

Let $A_0 = \{\Theta\}$, $A_1 = \langle \Theta, (1, -\frac{1}{4}) \rangle$. Suppose that A_0, \dots, A_n have been defined and $n \geq 1$. Define $b_n = a_n + a'_{n-1}$ and $A_{n+1} = A_n \cup (a_n + A'_{n-1}) \cup (b_n + A_n)$. See Figure 1.

It is easy to prove the following:

Assertion 1. For each $n > 0$, A_n is a polygon, $v_n < 0$, $A_n \subset (0, n] \times [v_n, 0] \cup \{\Theta\}$, $u_n = n$, $a_{n+1} = 2a_n + a'_{n-1}$ and $A_n = a_n - A_n$.

Given points p and q in A_n , $\langle\langle p, q \rangle\rangle$ will denote the subarc in A_n joining p and q , if $p \neq q$ and $\langle\langle p, q \rangle\rangle = \{p\}$, if $p = q$.

Assertion 2. For each $n > 0$, there exists a homeomorphism $\alpha_n : A_n \rightarrow A_{n+1}$ such that $\alpha_n(\Theta) = \Theta$, $\alpha_n(a_n) = a_{n+1}$ and, for each $p \in A_n$, $|\pi_1(p) - \pi_1(\alpha_n(p))| \leq 2$.

PROOF: We proceed by induction. Define $\alpha_1 : A_1 \rightarrow A_2$ by $\alpha_1(p) = 2p$. Define $\alpha_2 : A_2 \rightarrow A_3$ by sending homeomorphically the segment $A_1 \subset A_2$ onto the arc $A_2 \cup (a_2 + A'_1)$ in such a way that $\alpha_2(\Theta) = \Theta$ and then sending linearly the segment $a_1 + A_1$ onto $b_2 + A_2$. Suppose that $\alpha_1, \dots, \alpha_n$, have been constructed, define $\alpha_{n+1} : A_{n+1} \rightarrow A_{n+2}$ by:

$$\alpha_{n+1}(p) = \begin{cases} \alpha_n(p) & \text{if } p \in A_n, \\ b_{n+1} - (\alpha_{n-1}(-p' + b'_n))' & \text{if } p \in a_n + A'_{n-1} \text{ and,} \\ b_{n+1} + \alpha_n(p - b_n) & \text{if } p \in b_n + A_n. \end{cases}$$

It is easy to verify that α_{n+1} has the required properties. □

The key for proving the retractibility of the continuum presented in this paper is the following:

Assertion 3. For each $n > 0$ and for each $q \in A_n$, there exists a map $\sigma : A_n \rightarrow A_n$ (which depends on q) such that:

- (a) $\sigma|_{\langle\langle \Theta, q \rangle\rangle} = Id_{\langle\langle \Theta, q \rangle\rangle}$ (the identity map on $\langle\langle \Theta, q \rangle\rangle$);
- (b) $|\pi_1(p) - \pi_1(\sigma(p))| \leq 2$ for every $p \in A_n$;
- (c) if $q \in A_{n-1}$, then $\sigma^{-1}(\Theta) \cap \langle\langle q, a_n \rangle\rangle \neq \emptyset$: and if r is the first point in $\sigma^{-1}(\Theta) \cap \langle\langle q, a_n \rangle\rangle$ (in the natural ordering of $\langle\langle q, a_n \rangle\rangle$ from q to a_n), then $\sigma \langle\langle q, r \rangle\rangle \subset \langle\langle \Theta, q \rangle\rangle$ and $\sigma(a_n) = a_n$;
- (d) if $q \in A_n - A_{n-1}$, then $\sigma(A_n) \subset \langle\langle \Theta, q \rangle\rangle$.

In order to prove Assertion 3, we will use the following assertion which is easy to prove.

Assertion 4. If C and D are two arcs and $\alpha, \beta : C \rightarrow D$ are maps such that α is onto, then there exists $p \in C$ such that $\alpha(p) = \beta(p)$.

PROOF OF ASSERTION 3: We apply induction. It is easy to prove the assertion for $n = 1, n = 2$ and $n = 3$. Now, suppose that, for every $i = 1, \dots, n$ and for every $q \in A_i$, it is possible to construct σ and take $n \geq 3$. Take a point $q \in A_{n+1} = A_n \cup (a_n + A'_{n-1}) \cup (b_n + A_n)$. For defining σ we consider seven cases, in each one of which it is easy to check that the map defined has the required properties. A geometric representation of Cases 2–6 is given in Figures 2 and 3.

Case 1. $q \in A_{n-1} \subset A_n$. Apply the induction hypothesis to q and obtain the corresponding map $\sigma_0 : A_n \rightarrow A_n$. Define $\sigma : A_{n+1} \rightarrow A_{n+1}$ by:

$$\sigma(p) = \begin{cases} \sigma_0(p) & \text{if } p \in A_n \text{ and,} \\ p & \text{if } p \in (a_n + A'_{n-1}) \cup (b_n + A_n). \end{cases}$$

Case 2. $q \in A_n - A_{n-1}$. The induction hypothesis implies the existence of $\sigma_0 : A_n \rightarrow A_n$. Define $\alpha, \beta : a_n + A'_{n-1} \rightarrow A_n$ by $\alpha(p) = a_n - \alpha_{n-1}((p - a_n)')$ and $\beta(p) = \sigma_0(-(p - a_n)' + a_n)$. Applying Assertion 4, there exists $q_0 \in a_n + A'_{n-1}$ such that $\alpha(q_0) = \beta(q_0)$.

Define $\sigma : A_{n+1} \rightarrow A_{n+1}$ by:

$$\sigma(p) = \begin{cases} \sigma_0(p) & \text{if } p \in A_n, \\ \beta(p) & \text{if } \langle\langle a_n, q_0 \rangle\rangle, \\ \alpha(p) & \text{if } \langle\langle q_0, b_n \rangle\rangle \text{ and,} \\ a_{n+1} - \alpha_n(a_{n+1} - p) & \text{if } p \in b_n + A_n. \end{cases}$$

Case 3. $q \in a_n + A'_{n-2} \subset a_n + A'_{n-1}$ and $q \neq a_n$. Define $q_1 = (q - a_n)' \in A_{n-2} \subset A_{n-1}$. Apply the induction hypothesis to q_1 to obtain a map $\sigma_0 : A_{n-1} \rightarrow A_{n-1}$. Let r be the first point in $\langle\langle q_1, a_{n-2} \rangle\rangle$, in the ordering from q_1 to a_{n-2} , such that $\sigma_0(r) = \Theta$.

Define $\sigma : A_{n+1} \rightarrow A_{n+1}$ by:

$$\sigma(p) = \begin{cases} p & \text{if } p \in A_n, \\ a_n + (\sigma_0((p - a_n)'))' & \text{if } p \in \langle\langle a_n, a_n + r' \rangle\rangle, \\ a_n - \sigma_0((p - a_n)') & \text{if } p \in \langle\langle a_n + r', b_n \rangle\rangle \text{ and,} \\ b_{n-1} + \alpha_{n-1}^{-1}(p - b_n) & \text{if } p \in b_n + A_n. \end{cases}$$

Case 4. $q \in a_n + A'_{n-1} - (a_n + A'_{n-2})$. Define $q_1 = (q - a_n)' \in A_{n-1} - A_{n-2}$. Apply the induction hypothesis to q_1 to obtain a map $\sigma_0 : A_{n-1} \rightarrow A_{n-1}$.

Define $\sigma : A_{n+1} \rightarrow A_{n+1}$ by:

$$\sigma(p) = \begin{cases} p & \text{if } p \in A_n, \\ a_n + (\sigma_0((p - a_n)'))' & \text{if } p \in a_n + A'_{n-1}, \\ a_n - (\sigma_0(-(p - (b_n + a_{n-1}))))' & \text{if } p \in b_n + A_{n-1}, \\ a_n - \alpha_{n-2}((p - (b_n + a_{n-1})))' & \text{if } p \in b_n + a_{n-1} + A'_{n-2} \text{ and,} \\ b_{n-1} + p - (b_n + b_{n-1}) & \text{if } p \in b_n + b_{n-1} + A_{n-1}. \end{cases}$$

Case 5. $q \in b_n + A_{n-2} \subset b_n + A_n$. Define $q_1 = q - a_n - a'_{n-1} \in A_{n-2} \subset A_{n-1}$. Apply the induction hypothesis to q_1 to obtain a map $\sigma_0 : A_{n-1} \rightarrow A_{n-1}$. Let r be the first point in $\langle\langle q_1, a_{n-1} \rangle\rangle$, in the ordering from q_1 to a_{n-1} , such that $\sigma_0(r) = \Theta$.

Define $\sigma : A_{n+1} \rightarrow A_{n+1}$ by:

$$\sigma(p) = \begin{cases} p & \text{if } p \in A_n \cup (a_n + A'_{n-1}), \\ b_n + \sigma_0(p - b_n) & \text{if } p \in \langle\langle b_n, b_n + r \rangle\rangle, \\ b_n - (\sigma_0(p - b_n))' & \text{if } p \in \langle\langle b_n + r, b_n + a_{n-1} \rangle\rangle, \\ a_n - \alpha_{n-2}((p - (b_n + a_{n-1})))' & \text{if } p \in b_n + a_{n-1} + A'_{n-2} \text{ and,} \\ b_{n-1} + p - (b_n + b_{n-1}) & \text{if } p \in b_n + b_{n-1} + A_{n-1}. \end{cases}$$

Case 6. $q \in (b_n + A_{n-1}) - (b_n + A_{n-2})$. Define $q_1 = q - b_n \in A_{n-1} - A_n$. Apply the induction hypothesis to q_1 to obtain a map $\sigma_0 : A_{n-1} \rightarrow A_{n-1}$. Define $\alpha, \beta : b_n + a_{n-1} + A'_{n-2} \rightarrow b_n + A_{n-1}$ by: $\alpha(p) = b_n + a_{n-1} - \alpha_{n-2}((p - b_n - a_{n-1}))'$ and $\beta(p) = b_n + \sigma_0(a_{n-1} - (p - b_n - a_{n-1}))'$. From Assertion 4, there exists $p_0 \in b_n + a_{n-1} + A'_{n-2}$ such that $\alpha(p_0) = \beta(p_0)$.

Define $\sigma : A_{n+1} \rightarrow A_{n+1}$ by:

$$\sigma(p) = \begin{cases} p & \text{if } p \in A_n \cup (a_n + A'_{n-1}), \\ b_n + \sigma_0(p - b_n) & \text{if } p \in b_n + A_{n-1}, \\ \beta(p) & \text{if } p \in \langle\langle b_n + a_{n-1}, p_0 \rangle\rangle, \\ \alpha(p) & \text{if } p \in \langle\langle p_0, b_n + b_{n-1} \rangle\rangle \text{ and,} \\ b_n - (p - b_n - b_{n-1})' & \text{if } p \in b_n + b_{n-1} + A_{n-1}. \end{cases}$$

Case 7. $q \in (b_n + A_n) - (b_n + A_{n-1})$. Define $q_1 = q - b_n \in A_n - A_{n-1}$. Apply the induction hypothesis to q_1 to obtain a map $\sigma_0 : A_n \rightarrow A_n$.

Define $\sigma : A_{n+1} \rightarrow A_{n+1}$ by:

$$\sigma(p) = \begin{cases} p & \text{if } p \in A_n \cup (a_n + A'_{n-1}), \text{ and} \\ b_n + \sigma_0(p - b_n) & \text{if } p \in b_n + A_n. \end{cases}$$

This completes the construction of σ and the proof of Assertion 3. □

For each $n > 1$, define $C_n = (0, \frac{1}{2^{n-2}}) + \{(\frac{x}{n}, \frac{-y}{2^n v_n}) \in R^2 : (x, y) \in A_n\}$ and $B_n = C_n \cup \langle (0, \frac{1}{2^{n-2}}), \Theta \rangle \cup \langle \Theta, (1, 0) \rangle$. Given points p and q in B_n , $\langle\langle p, q \rangle\rangle$ will denote the subarc in B_n joining p and q if $p \neq q$ and $\langle\langle p, q \rangle\rangle = \{p\}$ if $p = q$.

Assertion 5. For each $n \geq 2$ and $q \in C_n$ there exists a retraction $\phi : B_n \rightarrow \langle\langle q, (1, 0) \rangle\rangle$ (which depends on q) such that $|\pi_1(p) - \pi_1(\phi(p))| \leq \frac{3}{n}$ for all $p \in B_n$.

PROOF: Let $\lambda : A_n \rightarrow C_n$ be the homeomorphism defined by $\lambda(x, y) = (0, \frac{1}{2^{n-2}}) + (\frac{x}{n}, \frac{-y}{2^n v_n})$. Let $q_0 = \lambda^{-1}(q) \in A_n$. Let $\sigma : A_n \rightarrow A_n$ be as in Assertion 3 applied to q_0 . Define $\sigma_1 : C_n \rightarrow C_n$ by $\sigma_1 = \lambda \circ \sigma \circ \lambda^{-1}$. We consider two cases. In both cases, it is easy to check that ϕ has the mentioned properties.

Case 1. $q_0 \in A_{n-1}$.

Let $r \in \langle\langle q_0, a_n \rangle\rangle$ be the first point, in the ordering from q_0 to a_n , such that $\sigma(r) = \Theta$. Choose a homeomorphism $\delta : [0, \frac{1}{n}] \rightarrow \langle\langle\langle (0, \frac{1}{2^{n-2}}), (\frac{1}{n}, 0) \rangle\rangle\rangle$ such that $\delta(0) = (0, \frac{1}{2^{n-2}})$ and $\delta(\frac{1}{n}) = (\frac{1}{n}, 0)$.

Define $\phi : B_n \rightarrow \langle\langle q, (1, 0) \rangle\rangle$ by:

$$\phi(p) = \begin{cases} p & \text{if } p \in \langle\langle\langle (0, \frac{1}{2^{n-2}}), (1, 0) \rangle\rangle\rangle, \\ \sigma_1(p) & \text{if } p \in \langle\langle\langle (0, \frac{1}{2^{n-2}}), \lambda(r) \rangle\rangle\rangle, \\ (\pi_1(\sigma_1(p)), 0) & \text{if } p \in \langle\langle\langle \lambda(r), \lambda(a_n) \rangle\rangle\rangle \cap (\pi_1 \circ \sigma_1)^{-1}([\frac{1}{n}, 1]) \text{ and,} \\ \delta(\pi_1(\sigma_1(p))) & \text{if } p \in \langle\langle\langle \lambda(r), \lambda(a_n) \rangle\rangle\rangle \cap (\pi_1 \circ \sigma_1)^{-1}([0, \frac{1}{n}]). \end{cases}$$

Case 2. $q_0 \in A_n - A_{n-1}$.

In this case define $\phi : B_n \rightarrow \langle\langle q, (1, 0) \rangle\rangle$ by:

$$\phi(p) = \begin{cases} p & \text{if } p \in \langle\langle\langle (0, \frac{1}{2^{n-2}}), (1, 0) \rangle\rangle\rangle \text{ and,} \\ \sigma_1(p) & \text{if } p \in C_n. \end{cases}$$

□

The example

Define $X = \bigcup\{B_n : n \geq 2\} = \langle \Theta, (0, 1) \rangle \cup \langle \Theta, (1, 0) \rangle \cup (\bigcup\{C_n : n \geq 2\})$. Clearly, X is a non-locally connected dendroid. The continuum X is illustrated in Figure 4.

In order to prove that X is retractible, define $J = \langle \Theta, (1, 0) \rangle$ and take a subcontinuum A of X . We may assume that $A \cap J \neq \emptyset$ and $A \not\subseteq J$. Since A is a retract of $A \cup J$, we only have to prove that there is a retraction $\rho : X \rightarrow A \cup J$. Let $N \geq 2$, be such that $A \cap C_n \neq \emptyset$ for every $n \geq N$ and $A \cap C_n = \emptyset$ for every $n \leq N$. Notice that, for each $n \geq 2$, C_n is an arc in R^2 which joins $(0, \frac{1}{2^{n-2}})$ and $(1, \frac{1}{2^{n-2}} - \frac{1}{2^n})$. For each $n \geq N$, let q_n be the last element in C_n , in the ordering from $(0, \frac{1}{2^{n-2}})$ to $(1, \frac{1}{2^{n-2}} - \frac{1}{2^n})$, such that $q_n \in A$. Then there exists a retraction $\phi_n : B_n \rightarrow \langle\langle q_n, (1, 0) \rangle\rangle$, such that $|\pi_1(p) - \pi_1(\phi_n(p))| \leq \frac{3}{n}$. Finally, let p_0 be the last point in $\langle \Theta, (0, 1) \rangle$, in the ordering from Θ to $(0, 1)$, such that $p_0 \in A$. We are ready to define ρ .

Define $\rho : X \rightarrow A \cup J$ by:

$$\rho(p) = \begin{cases} p_0 & \text{if } p \in \langle p_0, (0, 1) \rangle \cup (\bigcup\{C_n : n < N\}), \\ p & \text{if } p \in \langle (0, \frac{1}{2^{N-2}}, p_0 \rangle \text{ and,} \\ \phi_n(p) & \text{if } p \in B_n \text{ for some } n \geq N. \end{cases}$$

Then ρ is a retraction.

Therefore, X is retractible.

A dendroid is called a *fan* provided that it has exactly one ramification point. A continuum is said to be *rational* provided that each of its points has arbitrarily small neighborhoods with countable boundaries. Shrinking in the constructed dendroid X the arc $\langle \Theta, (0, 1) \rangle$ to a point, i.e., applying a monotone mapping $\mu : X \rightarrow Y$ such that $\mu(\langle \Theta, (0, 1) \rangle)$ is a singleton, while the partial mapping $\mu|(X - \langle \Theta, (0, 1) \rangle)$ is a homeomorphism, we get a rational plane fan Y that keeps the main property of X of being retractible.

Acknowledgment. The author wishes to thank the referee for his suggestions.




Figure 1




Figure 2





Figure 3




Figure 4

REFERENCES

- [1] Charatonik J.J., *On retractible continua*, General Topology and its Relations to Modern Analysis and Algebra VI, Proc. Sixth Prague Topological Symposium 1986, Z. Frolík (ed.), Heldermann Verlag, Berlin, 1988, pp. 83–90.
- [2] Gordh G.R., Jr., Lum L., *Monotone retracts and some characterizations of dendrites*, Proc. Amer. Math. Soc. **59** (1976), 156–158.
- [3] Illanes A., *Characterizing dendrites by deformation retractions*, Topology Proc. **21** (1996), 129–141.

- [4] Illanes A., *Confluent and open retractible continua*, preprint.
- [5] Lum L., *A characterization of local connectivity in dendroids*, Studies in Topology, Academic Press, New York, 1975, pp. 331–338.

INSTITUTO DE MATEMÁTICAS, UNAM, CIRCUITO EXTERIOR, CD. UNIVERSITARIA, MÉXICO
04510, D.F., MÉXICO

E-mail: illanes@gauss.matem.unam.mx

(Received September 2, 1996, revised October 20, 1997)