Fixed points of asymptotically regular mappings in spaces with uniformly normal structure

Jarosław Górnicki

Abstract. It is proved that: for every Banach space X which has uniformly normal structure there exists a $k > 1$ with the property: if A is a nonempty bounded closed convex subset of X and $T : A \to A$ is an asymptotically regular mapping such that

$$\liminf_{n\to\infty} |||T^n||| < k,$$

where $|||T|||$ is the Lipschitz constant (norm) of T, then T has a fixed point in A.

Keywords: asymptotically regular mappings, uniformly normal structure, fixed points

Classification: 47H10

1. Introduction.

The concept of uniformly normal structure is due to A.A. Gillespie and B.B. Williams [7]. A Banach space X has uniformly normal structure if

$$N(X) = \sup\{r_A(A) : A \subset X, \ \text{ convex, } \operatorname{diam} A = 1\} < 1,$$

where

$$r_A(A) = \inf\left\{\sup\{\|x - y\| : y \in A\} : x \in A\right\}.$$

It was proved in [4], [2] that $N(X) \le 1 - \delta_X(1)$; thus $\varepsilon_0(X) < 1$ implies uniformly normal structure. In the paper [11] X.T. Yu proved that if X is a uniformly smooth space (or more generally, $\lim_{t\downarrow 0} \rho_X(t)t^{-1} < \frac{1}{2}$), then X has a uniformly normal structure. Also, in [12] it was proved that uniformly normal structure does not necessarily imply that the space has good geometric properties.

The concept of asymptotic regularity is due to F. Browder and V. Petryshyn [1]. A mapping $T : X \to X$ is said to be asymptotically regular if

$$\lim_{n\to\infty} \|T^{n+1}x - T^n x\| = 0$$

for all $x \in X$.

If T is nonexpansive, then $T_\lambda := \lambda \cdot I + (1 - \lambda) \cdot T$ is asymptotically regular for all $0 < \lambda < 1$ (see [6]).

Recently P.K. Lin in [10] has constructed a uniformly asymptotically regular Lipschitzian mapping acting on a weakly compact subset of l^2 which has no fixed point.

E.A. Lifshitz (see [5]) associated with each metric space (M, d) a constant $\kappa(M) \geq 1$. Define Lifshitz characteristic $\kappa_0(X)$ to be the infimum of $\kappa(C)$ where C ranges over all nonempty closed bounded convex subsets of the Banach space X. D.J. Downing and B. Turett [5] proved the following

Theorem 1. *Let X be a Banach space.*

(1) *Then $\varepsilon_0(X) < 1$ if and only if $\kappa_0(X) > 1$.*
(2) *If $\gamma > 1$ satisfies $\gamma(1 - \delta_X(\gamma^{-1})) = 1$, then $\gamma \leq \kappa_0(X)$.*

In [8] the present author proved the following

Theorem 2. *Let X be a Banach space with the Lifshitz characteristic $\kappa_0(X) > 1$ and let C be a nonempty bounded closed convex subset of X. If $T : C \to C$ is an asymptotically regular mapping such that*

$$\liminf_{n\to\infty} |||T^n||| < \kappa_0(X),$$

then T has a fixed point in C.

2. Main result.

The main result of this paper is interesting in the Banach spaces X which satisfy the conditions: $\varepsilon_0(X) \geq 1$ and $N(X) < 1$ (cf. [3]).

We start with the following

Lemma 1 [3]**.** *Let X be a Banach space with $N(X) < 1$. Then for every bounded sequence $\{x_n\}$ there exists a point $z \in \overline{\mathrm{conv}}\{x_n\}$, such that:*

(i) $\limsup_{n\to\infty} \|z - x_n\| \leq N(X) \cdot \lim_{s\to\infty} \sup\{\|x_n - x_m\| : n, m \geq s\}$,
(ii) *for every $y \in X$, $\|z - y\| \leq \limsup_{n\to\infty} \|y - x_n\|$.*

Lemma 2 [9]**.** *Let A be a nonempty closed convex subset of a Banach space X and let $\{n_i\}$ be an increasing sequence of natural numbers. Assume that $T : A \to A$ is an asymptotically regular mapping such that for some $m \in \mathbb{N}$, T^m is continuous. If*

$$\hat{r}(x) = \limsup_{i\to\infty} \|x - T^{n_i} u\| = 0$$

for some $u \in A$ and $x \in A$, then $Tx = x$.

Theorem 3. *Let A be a nonempty bounded closed convex subset of a Banach space X which has uniformly normal structure, i.e. $N(X) < 1$. If $T : A \to A$ is an asymptotically regular mapping such that*

$$\liminf_{n\to\infty} |||T^n||| = k < [N(X)]^{-1/2},$$

then T has a fixed point in A.

PROOF: Let $T : A \to A$ and let $\{n_i\}$ be a sequence of natural numbers such that

$$\liminf_{n\to\infty} |||T^n||| = \lim_{i\to\infty} |||T^{n_i}||| = k < [N(X)]^{-1/2}.$$

Consider the sequence $\{T^{n_i}x\}$ for an $x \in A$. Let $z(x)$ be a point satisfying Lemma 1 for $\{T^{n_i}x\}$. Let $r(x) = \limsup_{i\to\infty} \|T^{n_i}x - x\|$. By the condition (i) of Lemma 1, we have

(1)
$$\begin{aligned}
\limsup_{i\to\infty} \|T^{n_i}x - z\| &\le N(X) \cdot \lim_{s\to\infty} \sup\{\|T^{n_i}x - T^{n_j}x\| : n_i, n_j \ge s\} \le \\
&\le N(X) \cdot \limsup_{i\to\infty}\big(\limsup_{j\to\infty} \|T^{n_i}x - T^{n_j}x\|\big) \le \\
&\le N(X) \cdot \limsup_{i\to\infty}\big(\limsup_{j\to\infty}(\|T^{n_i}x - T^{n_i+n_j}x\| + \|T^{n_i+n_j}x - T^{n_j}x\|)\big) \le \\
&\le N(X) \cdot \limsup_{i\to\infty}\Big(\limsup_{j\to\infty}(|||T^{n_i}||| \cdot \|x - T^{n_j}x\| + \sum_{v=0}^{n_i-1} \|T^{n_j+v+1}x - T^{n_j+v}x\|)\Big) \le \\
&\le N(X) \cdot \limsup_{i\to\infty} |||T^{n_i}||| \cdot \limsup_{j\to\infty} \|x - T^{n_j}x\| = \\
&= k \cdot N(X) \cdot \limsup_{j\to\infty} \|x - T^{n_j}x\|.
\end{aligned}$$

Moreover, for $i > 1$, we have

$$\begin{aligned}
\|T^{n_i}z - z\| &\le \limsup_{j\to\infty} \|T^{n_i}z - T^{n_j}x\| \le \\
&\le \limsup_{j\to\infty}\big(\|T^{n_i}z - T^{n_i+n_j}x\| + \|T^{n_i+n_j}x - T^{n_j}x\|\big) \le \\
&\le \limsup_{j\to\infty}\Big(|||T^{n_i}||| \cdot \|z - T^{n_j}x\| + \sum_{v=0}^{n_i-1} \|T^{n_j+v+1}x - T^{n_j+v}x\|\Big) \le \qquad (2) \\
&\le |||T^{n_i}||| \cdot \limsup_{j\to\infty} \|z - T^{n_j}x\|.
\end{aligned}$$

By (1) and (2)

$$r(z) \le k^2 \cdot N(X) \cdot r(x) = a \cdot r(x), \quad \text{with} \quad a < 1. \qquad (3)$$

Define a sequence $\{x_m\}$ in the following way: x_1 is an arbitrarily chosen point of A, $x_{m+1} = z(x_m)$. Then $\{x_m\}$ is a Cauchy sequence. In fact, we have

$$\begin{aligned}
\|x_{m+1} - x_m\| &\le \|x_{m+1} - T^{n_i}x_m\| + \|T^{n_i}x_m - x_m\| \le \\
&\le \|x_{m+1} - T^{n_i}x_m\| + r(x_m).
\end{aligned}$$

Taking the limit superior as $i \to +\infty$,

$$\|x_{m+1} - x_m\| \le \limsup_{i\to\infty} \|x_{m+1} - T^{n_i} x_m\| + r(x_m) \le$$
$$\le k \cdot N(X) \cdot r(x_m) + r(x_m) = [1 + k \cdot N(X)] \cdot r(x_m).$$

Hence, by (3)

$$\|x_{m+1} - x_m\| \le [1 + k \cdot N(X)] \cdot r(x_m) \le [1 + k \cdot N(X)] \cdot a^m \cdot r(x_1) \to 0$$

as $m \to +\infty$. Let $x_0 = \lim_{m\to\infty} x_m$. Finally

$$\|x_0 - T^{n_i} x_0\| \le \|x_0 - x_m\| + \|x_m - T^{n_i} x_m\| + \|T^{n_i} x_m - T^{n_i} x_0\| \le$$
$$\le \big(1 + |||T^{n_i}|||\big) \cdot \|x_0 - x_m\| + \|x_m - T^{n_i} x_m\|.$$

Taking the limit superior as $i \to +\infty$ on both sides we get

$$\limsup_{i\to\infty} \|x_0 - T^{n_i} x_0\| \le (1 + k) \cdot \|x_0 - x_m\| + r(x_m) \le$$
$$\le (1 + k) \cdot \|x_0 - x_m\| + a^m \cdot r(x_1) \to 0$$

as $m \to +\infty$. Therefore, by Lemma 2, $Tx_0 = x_0$. □

For James spaces $X_M = \big(l^2, |\cdot|_M\big)$, where $|\cdot|_M = \max\{\|\cdot\|_2, M\cdot\|\cdot\|_\infty\}$, $(M \ge 1)$ we have

1)
$$\varepsilon_0(X_M) = \begin{cases} 2\cdot(M^2-1)^{1/2} & \text{for } M < \sqrt{2}, \\ 2 & \text{for } M > \sqrt{2}, \end{cases}$$
and $\varepsilon_0(X_M) < 1$ if and only if $M < \frac{\sqrt{5}}{2}$;
2) for $1 \le M < \frac{\sqrt{5}}{2}$, the condition $\gamma < [N(X_M)]^{-1/2}$ is weaker than $\gamma < \gamma_0$, where γ_0 is the unique solution of $x\big(1 - \delta_{X_M}(\frac{1}{x})\big) = 1$;

and

$$N(X_M) = \frac{M}{\sqrt{2}} \quad \text{for} \quad 1 \le M \le \sqrt{2}, \ [3].$$

Combining these results we get the following

Corollary 1. *Let A be a nonempty bounded closed convex subset of a James space X_M, $1 \le M < \sqrt{2}$. If $T : A \to A$ is an asymptotically regular mapping such that*

$$\liminf_{n\to\infty} |||T^n||| < \frac{2^{1/4}}{\sqrt{M}},$$

then T has a fixed point in A.

References

[1] Browder F.E., Petryshyn V.W., *The solution by iteration of nonlinear functional equations in Banach spaces*, Bull. AMS **72** (1966), 571–576.
[2] Bynum W.L., *Normal structure coefficients for Banach spaces*, Pacific J. Math. **86** (1980), 427–436.
[3] Casini E., Maluta E., *Fixed points of uniformly Lipschitzian mappings in spaces with uniformly normal structure*, Nonlinear Anal., TMA **9** (1985), 103–108.
[4] Daneš J., *On densifying and related mappings and their applications in nonlinear functional analysis*, in: Theory of Nonlinear Operators (Proc. Summer School, October 1972, GDR), Akademie-Verlag, Berlin, 1974, 15–56.
[5] Downing D.J., Turett B., *Some properties of the characteristic convexity relating to fixed point theory*, Pacific J. Math. **104** (1983), 343–350.
[6] Edelstein M., O'Brien C.R., *Nonexpansive mappings, asymptotic regularity and successive approximations*, J. London Math. Soc. (2) **17** (1978), 547–554.
[7] Gillespie A.A., Williams B.B., *Fixed point theorem for nonexpansive mappings on Banach spaces with uniformly normal structure*, Appl. Anal. **9** (1979), 121–124.
[8] Górnicki J., *A fixed point theorem for asymptotically regular mappings*, to appear.
[9] Krüppel M., *Ein Fixpunktsatz für asymptotisch reguläre Operatoren in gleichmäßig konvexen Banach-Räumen*, Wiss. Z. Pädagog. Hochsch. "Liselotte Herrmann" Güstrow, Math.-naturwiss. Fak. **25** (1987), 241–246.
[10] Lin P.K., *A uniformly asymptotically regular mapping without fixed points*, Canad. Math. Bull. **30** (1987), 481–483.
[11] Yu X.T., *On uniformly normal structure*, Kexue Tongbao **33** (1988), 700–702.
[12] ———, *A geometrically aberrant Banach space with uniformly normal structure*, Bull. Austral. Math. Soc. **38** (1988), 99–103.

Institute of Mathematics, Pedagogical University of Rzeszów, Rejtana 16 A, 35–310 Rzeszów, Poland

(*Received* September 20, 1990, *revised* October 7, 1991)