

Markku Niemenmaa

On finite commutative IP-loops with elementary abelian inner mapping groups of order p^4

Comment.Math.Univ.Carolin. 51,4 (2010) 559–563.

Abstract: We show that finite commutative inverse property loops with elementary abelian inner mapping groups of order p^4 are centrally nilpotent of class at most two.

Keywords: loop, inner mapping group

AMS Subject Classification: 20D10, 20N05

REFERENCES

1. Bruck R.H., *Contributions to the theory of loops*, Trans. Amer. Math. Soc. **60** (1946), 245–354.
2. Csörgő P., *On connected transversals to abelian subgroups and loop theoretical consequences*, Arch. Math. (Basel) **86** (2006), no. 6, 499–516.
3. Csörgő P., *Abelian inner mappings and nilpotency class greater than two*, European J. Combin. **28** (2007), no. 3, 858–867.
4. Drápal A., Vojtěchovský P., *Explicit constructions of loops with commuting inner mappings*, European J. Combin. **29** (2008), no. 7, 1662–1681.
5. Kepka T., Niemenmaa M., *On multiplication groups of loops*, J. Algebra **135** (1990), 112–122.
6. Kepka T., Niemenmaa M., *On connected transversals to abelian subgroups in finite groups*, Bull. London Math. Soc. **24** (1992), 343–346.
7. Kepka T., Niemenmaa M., *On connected transversals to abelian subgroups*, Bull. Australian Math. Soc. **49** (1994), 121–128.
8. Kinyon M., *private communication*, 2009.
9. Niemenmaa M., *On finite loops whose inner mapping groups are abelian II*, Bull. Australian Math. Soc. **71** (2005), 487–492.
10. Niemenmaa M., Rytty M., *Connected transversals and multiplication groups of loops*, Quasigroups and Related Systems **15** (2007), 95–107.