

Generalized n -coherence

J. JIRÁSKO

Abstract. In this paper necessary and sufficient conditions for large subdirect products of n -flat modules from the category $Gen(Q)$ to be n -flat are given.

Keywords: relative finiteness conditions, relative coherence, large subdirect products of n -flat modules

Classification: 16D40

In what follows, R stands for an associative ring with a unit element and $R\text{-Mod}$ ($\text{Mod-}R$) denotes the category of all unitary left (right) R -modules.

Let \mathcal{F} be a filter on a set I and $\{M_i; i \in I\}$ be a family of left R -modules. We define an equivalence relation \sim on $\prod_{i \in I} M_i$ as follows: For $(m_i), (n_i) \in \prod_{i \in I} M_i$, $(m_i) \sim (n_i)$ if $\{i \in I; m_i = n_i\} \in \mathcal{F}$. The equivalence class of $(0, 0, \dots)$ is called the \mathcal{F} -product and it is denoted by $\prod_{i \in I}^{\mathcal{F}} M_i$. Clearly, $\prod_{i \in I}^{\mathcal{F}} M_i$ is a submodule of $\prod_{i \in I} M_i$. For a set X let $|X|$ denotes the cardinality of X and for $m = (m_i)_{i \in I} \in \prod_{i \in I} M_i$ let $\text{supp}(m) = \{i \in I; m_i \neq 0\}$. For an infinite cardinal number \aleph the \aleph -product is defined as $\prod_{i \in I}^{\aleph} M_i = \{m \in \prod_{i \in I} M_i; |\text{supp}(m)| < \aleph\}$. For an infinite cardinal number \aleph let \aleph^+ be its immediate successor. Let \mathcal{F} be a filter on an index set I and let \aleph be $\text{sup}\{|I \setminus X|; X \in \mathcal{F}\}$. According to [9] we define $\text{sup}(\mathcal{F})$ to be \aleph if the supremum is not attained and \aleph^+ if the supremum is attained. If \aleph is an infinite cardinal number and $|I| \geq \aleph$ then $\mathcal{F} = \{X \subseteq I; |I \setminus X| < \aleph\}$ is a filter on I with $\text{sup}(\mathcal{F}) = \aleph$ and $\prod_{i \in I}^{\aleph} M_i = \prod_{i \in I}^{\mathcal{F}} M_i$. If $|I| < \aleph$ then obviously $\prod_{i \in I}^{\aleph} M_i = \prod_{i \in I} M_i$. If $|I| = \aleph_s$ then we have $\sum_{i \in I}^{\oplus} M_i = \prod_{i \in I}^{\aleph_0} M_i \subseteq \prod_{i \in I}^{\aleph_1} M_i \subseteq \dots \subseteq \prod_{i \in I}^{\aleph_s} M_i \subseteq \prod_{i \in I}^{\aleph_{s+1}} M_i = \prod_{i \in I} M_i$. The \mathcal{F} -products (\aleph -products) of flat and projective modules were investigated in [9] and [10] by P. Loustaunau.

Let n be a nonnegative integer. A module $M \in \text{Mod-}R$ is called n -presented if there is a finite n -presentation of M i.e. an exact sequence

$$F_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_1 \rightarrow F_0 \rightarrow M \rightarrow 0$$

in which every F_i is free of finite rank. A ring R is said to be right n -coherent if every n -presented right module is $(n + 1)$ -presented. The following definition of n -flat and n -FP-injective module is due to J. Chen and N. Ding. Let n be

The author has been supported by GA ČR 201/97/1162.

a positive integer. A left R -module Q is called n -flat if $\text{Tor}_n^R(N, Q) = 0$ for all n -presented right R -modules N . A right R -module M is said to be n -FP-injective if $\text{Ext}_R^n(N, M) = 0$ for all n -presented right R -modules N .

In [3] J. Chen and N. Ding characterize right n -coherent rings as rings for which direct products of n -flat left R -modules are n -flat. In [5] (\aleph, Q) -coherent rings were introduced and they were characterized as rings for which \aleph -products of flat modules from the category $\text{Gen}(Q)$ are flat. These rings were also studied in [11]. The aim of this paper is to generalize results of J. Chen and N. Ding and the results in [5] to \aleph -products of n -flat modules from the category $\text{Gen}(Q)$ for a fixed flat module Q .

Throughout all the paper ${}_R Q$ denotes a fixed flat left R -module and \aleph denotes an infinite cardinal number.

The notions of (\aleph, Q) -finitely generated, (\aleph, Q) -finitely presented and (\aleph, Q) -coherent modules were introduced in [5]. In the following lemmas we summarize basic properties of these modules.

Lemma 1.1. *Let $\{Q_i; i \in I\}$ be a set of left R -modules. Then*

- (i) *if \mathcal{F} is a filter on I with $\text{sup}(\mathcal{F}) \leq \aleph$ then $\prod_{i \in I}^{\mathcal{F}} Q_i \subseteq \prod_{i \in I}^{\aleph} Q_i$;*
- (ii) *let \mathcal{F} be a filter on I with $\text{sup}(\mathcal{F}) = \aleph$ and $q \in \prod_{i \in I}^{\aleph} Q_i$. If $S = \text{supp}(q)$ then there is $X \in \mathcal{F}$ and an injective map $f: S \rightarrow I \setminus X$. Since $X \subseteq I \setminus f(S)$ the element \bar{q} defined by $\bar{q}_i = q_{f^{-1}(i)}$ for $i \in f(S)$ and $\bar{q}_i = 0$ for $i \in I \setminus f(S)$ belongs to $\prod_{i \in I}^{\mathcal{F}} Q_i$.*

PROOF: (i). If $q \in \prod_{i \in I}^{\mathcal{F}} Q_i$ then $|\text{supp}(q)| < \text{sup}(\mathcal{F}) \leq \aleph$ and consequently $q \in \prod_{i \in I}^{\aleph} Q_i$.

(ii). If $\text{sup}(\mathcal{F}) = \aleph$ and $|S| < \aleph$ then there is $X \in \mathcal{F}$ with $|S| \leq |I \setminus X|$. The rest is clear. \square

Lemma 1.2. *Let \mathcal{F} be a filter on I with $\text{sup}(\mathcal{F}) = \aleph$, $\{Q_i; i \in I\}$ be a family of left R -modules and M be a right R -module. Then the following conditions are equivalent:*

- (i) *the natural homomorphism $\varphi_{\mathcal{F}}: M \otimes_R \prod_{i \in I}^{\mathcal{F}} Q_i \rightarrow \prod_{i \in I}^{\mathcal{F}} (M \otimes_R Q_i)$ defined via $\varphi_{\mathcal{F}}(m \otimes (q_i)_{i \in I}) = (m \otimes q_i)_{i \in I}$ is an epimorphism;*
- (ii) *the natural homomorphism $\varphi_{\aleph}: M \otimes_R \prod_{i \in I}^{\aleph} Q_i \rightarrow \prod_{i \in I}^{\aleph} (M \otimes_R Q_i)$ defined via $\varphi_{\aleph}(m \otimes (q_i)_{i \in I}) = (m \otimes q_i)_{i \in I}$ is an epimorphism.*

PROOF: (i) implies (ii). Let $\varphi_{\mathcal{F}}$ be an epimorphism, $q \in \prod_{i \in I}^{\aleph} (M \otimes Q_i)$, $S = \text{supp}(q)$ and $\bar{q} \in \prod_{i \in I}^{\mathcal{F}} (M \otimes Q_i)$ be the element defined in Lemma 1.1(ii). Then there is an element $m_1 \otimes q_1 + \cdots + m_r \otimes q_r \in M \otimes \prod_{i \in I}^{\mathcal{F}} Q_i$ with $(m_1 \otimes q_{1_i} + \cdots + m_r \otimes q_{r_i})_{i \in I} = \bar{q}$. We can assume without loss of generality that $q_{i_j} = 0$ for $j \in I \setminus f(S)$ and $i = 1, \dots, r$. Let $p_j \in \prod_{i \in I}^{\aleph} Q_i$ such that $p_{j_t} = 0$ for $t \in I \setminus S$ and $p_{j_s} = q_{j f(s)}$ for $s \in S$, $j = 1, \dots, r$. Hence $q_s = \bar{q}_{f(s)} = m_1 \otimes q_{1 f(s)} + \cdots + m_r \otimes q_{r f(s)} = m_1 \otimes p_{1_s} + \cdots + m_r \otimes p_{r_s}$ for $s \in S$ and consequently φ_{\aleph} is an epimorphism.

(ii) implies (i). If φ_{\aleph} is an epimorphism and $q \in \prod_{i \in I}^{\mathcal{F}} (M \otimes Q_i)$ then there is an element $m_1 \otimes q_1 + \dots + m_r \otimes q_r \in M \otimes \prod_{i \in I}^{\aleph} Q_i$ with $(m_1 \otimes q_{1i} + \dots + m_r \otimes q_{ri})_{i \in I} = q$. If $S = \text{supp}(q)$ then $I \setminus S \in \mathcal{F}$. Without loss of generality we can take q_i such that $q_{ij} = 0$ for $j \in I \setminus S$ and $i = 1, \dots, r$. Thus $q_i \in \prod_{i \in I}^{\mathcal{F}} Q_i$ for $i = 1, \dots, r$ and consequently $\varphi_{\mathcal{F}}$ is an epimorphism. \square

The following definition is motivated by the definition of R.R. Colby and E.A. Rutter of the Q -finitely generated module in [4] and the definition of P. Loustau of the \aleph -finitely generated module in [9].

Definition 1.3. A right R -module M is said to be (\aleph, Q) -finitely generated if every subset T of $M \otimes_R Q$ with $|T| < \aleph$ is contained in $N \otimes_R Q$ for some finitely generated submodule N of a module M .

Lemma 1.4. Let M be a right R -module. Then the following conditions are equivalent:

- (i) M is (\aleph, Q) -finitely generated;
- (ii) if I is a set and $Q_i \in \text{Gen}(Q)$, $i \in I$ then the natural homomorphism $\varphi: M \otimes_R \prod_{i \in I}^{\aleph} Q_i \rightarrow \prod_{i \in I}^{\aleph} (M \otimes_R Q_i)$ defined via $\varphi(m \otimes (q_i)_{i \in I}) = (m \otimes q_i)_{i \in I}$ is an epimorphism;
- (iii) if I is a set then the natural homomorphism $\varphi: M \otimes_R \prod_{i \in I}^{\aleph} Q \rightarrow \prod_{i \in I}^{\aleph} (M \otimes_R Q)$ defined via $\varphi(m \otimes (q_i)_{i \in I}) = (m \otimes q_i)_{i \in I}$ is an epimorphism.

PROOF: (i) implies (ii). Let $u \in \prod_{i \in I}^{\aleph} (M \otimes Q_i)$, $T = \text{supp}(u)$ and $f_i: Q^{(J_i)} \rightarrow Q_i$, $i \in I$ be epimorphisms. Then $|T| < \aleph$ and $\text{id}_M \otimes f_i: M \otimes Q^{(J_i)} \rightarrow M \otimes Q_i$, $i \in I$ are epimorphisms. Hence $u_i = \sum_{j=1}^{n_i} m_{ij} \otimes f_i(q_{ij})$, where $m_{ij} \in M$, $q_{ij} \in Q^{(J_i)}$, $i \in I$ and $j = 1, \dots, n_i$. Now $q_{ij} = \sum_{k=1}^{t_{ij}} q_{ijk}$, where $q_{ijk} \in Q$, $k = 1, \dots, t_{ij}$. Let $S = \{m_{ij} \otimes q_{ijk}; i \in T, j = 1, \dots, n_i, k = 1, \dots, t_{ij}\}$. Then $|S| < \aleph$ and $S \subseteq M \otimes Q$. Thus $S \subseteq N \otimes Q$ for some finitely generated submodule $N = \sum_{p=1}^l n_p R$ of M . Hence $m_{ij} \otimes q_{ijk} = \sum_{p=1}^l n_p \otimes q_{ijkp}$ for some $q_{ijkp} \in Q$. Put $v_{ip} = \sum_{j=1}^{n_i} \sum_{k=1}^{t_{ij}} q_{ijkp}$ for $i \in T$ and $v_{ip} = 0$ for $i \in I \setminus T$, $p = 1, \dots, l$. Then $w_p = (f_i(v_{ip}))_{i \in I} \in \prod_{i \in I}^{\aleph} Q_i$, $p = 1, \dots, l$ and $u_i = \sum_{p=1}^l n_p \otimes f_i(\sum_{j=1}^{n_i} \sum_{k=1}^{t_{ij}} q_{ijkp}) = \sum_{p=1}^l n_p \otimes f_i(v_{ip})$, $i \in I$. Thus $\varphi(\sum_{p=1}^l n_p \otimes w_p) = (\sum_{p=1}^l n_p \otimes f_i(v_{ip}))_{i \in I} = u$ and consequently φ is an epimorphism.

(ii) implies (iii). Obvious.

(iii) implies (i). Let $S \subseteq M \otimes Q$ with $|S| < \aleph$ and I be a set such that $|S| \leq |I|$ (e.g. $I = M \otimes Q$ or I is a set with $|I| \geq \aleph$). Then there is an injective map $f: S \rightarrow I$. Let us consider $u \in \prod_{i \in I}^{\aleph} (M \otimes Q)$ defined by $u_i = f^{-1}(i)$ for $i \in f(S)$ and $u_i = 0$ for $i \in I \setminus f(S)$. Then by assumption there is $\sum_{j=1}^r m_j \otimes q_j \in M \otimes \prod_{i \in I}^{\aleph} Q$ such that $(\sum_{j=1}^r m_j \otimes q_{j_i})_{i \in I} = u$. Now if $s \in S$ then $s = f^{-1}(i) = \sum_{j=1}^r m_j \otimes q_{j_i}$ for some $i \in f(S)$ and therefore $S \subseteq N \otimes Q$, where $N = \sum_{j=1}^r m_j R$ is a finitely generated submodule of M . \square

Corollary 1.5. *The class of all (\aleph, Q) -finitely generated modules is closed under extensions, homomorphic images, finite direct sums, direct summands and contains the class of all finitely generated modules.*

PROOF: It follows immediately from Lemma 1.4(ii) and the definition of (\aleph, Q) -finitely generated module. \square

Definition 1.6. *A right R -module M is said to be (\aleph, Q) -finitely presented if there is a short exact sequence $0 \rightarrow K \rightarrow F \rightarrow M \rightarrow 0$ with F free of finite rank and K (\aleph, Q) -finitely generated.*

Lemma 1.7. *Let M be a finitely generated right R -module. Then the following conditions are equivalent:*

- (i) M is (\aleph, Q) -finitely presented;
- (ii) if $0 \rightarrow K \rightarrow P \rightarrow M \rightarrow 0$ is a projective presentation with P finitely generated then K is (\aleph, Q) -finitely generated;
- (iii) if I is a set and $Q_i \in \text{Gen}(Q)$, $i \in I$ then the natural homomorphism $\varphi: M \otimes_R \prod_{i \in I}^{\aleph} Q_i \rightarrow \prod_{i \in I}^{\aleph} (M \otimes_R Q_i)$ defined via $\varphi(m \otimes (q_i)_{i \in I}) = (m \otimes q_i)_{i \in I}$ is an isomorphism;
- (iv) if I is a set then the natural homomorphism $\varphi: M \otimes_R \prod_{i \in I}^{\aleph} Q \rightarrow \prod_{i \in I}^{\aleph} (M \otimes_R Q)$ defined via $\varphi(m \otimes (q_i)_{i \in I}) = (m \otimes q_i)_{i \in I}$ is an isomorphism.

PROOF: (i) implies (ii). Let $0 \rightarrow K_i \rightarrow P_i \rightarrow M \rightarrow 0$, $i = 1, 2$ be two projective presentations of M . By Schanuel's Lemma we have $P_1 \oplus K_2 \simeq P_2 \oplus K_1$. Now if P_1, P_2 are finitely generated and K_1 is (\aleph, Q) -finitely generated then K_2 is (\aleph, Q) -finitely generated by Corollary 1.5.

(ii) implies (iii). Let $0 \rightarrow K \rightarrow F \rightarrow M \rightarrow 0$ be an exact sequence, where F is free of finite rank and $Q_i \in \text{Gen}(Q), i \in I$. Consider the following commutative diagram

$$\begin{array}{ccccccc} K \otimes \prod_{i \in I}^{\aleph} Q_i & \longrightarrow & F \otimes \prod_{i \in I}^{\aleph} Q_i & \longrightarrow & M \otimes \prod_{i \in I}^{\aleph} Q_i & \longrightarrow & 0 \\ \downarrow \varphi_K & & \downarrow \varphi_F & & \downarrow \varphi_M & & \\ \prod_{i \in I}^{\aleph} (K \otimes Q_i) & \longrightarrow & \prod_{i \in I}^{\aleph} (F \otimes Q_i) & \longrightarrow & \prod_{i \in I}^{\aleph} (M \otimes Q_i) & \longrightarrow & 0 \end{array} .$$

Then φ_F is obviously an isomorphism since F is free of finite rank and φ_K is an epimorphism since K is (\aleph, Q) -finitely generated. Hence φ_M is an isomorphism.

(iii) implies (iv). Obvious.

(iv) implies (i). Let $0 \rightarrow K \rightarrow F \rightarrow M \rightarrow 0$ be an exact sequence with F free of finite rank. Consider the following commutative diagram

$$\begin{array}{ccccccc} K \otimes \prod_{i \in I}^{\aleph} Q & \longrightarrow & F \otimes \prod_{i \in I}^{\aleph} Q & \longrightarrow & M \otimes \prod_{i \in I}^{\aleph} Q & \longrightarrow & 0 \\ \downarrow \varphi_K & & \downarrow \varphi_F & & \downarrow \varphi_M & & \\ 0 \longrightarrow & \prod_{i \in I}^{\aleph} (K \otimes Q) & \longrightarrow & \prod_{i \in I}^{\aleph} (F \otimes Q) & \longrightarrow & \prod_{i \in I}^{\aleph} (M \otimes Q) & \longrightarrow & 0 \end{array} .$$

Now φ_F and φ_M are isomorphisms. Hence φ_K is an epimorphism and K is (\aleph, Q) -finitely generated by Lemma 1.4. \square

Remark 1.8. As it follows from Lemma 1.2 and the proof of Lemma 1.7 every \aleph -product $\prod_{i \in I}^{\aleph}$ in Lemma 1.4 and Lemma 1.7 can be replaced by \mathcal{F} -product $\prod_{i \in I}^{\mathcal{F}}$ for a filter \mathcal{F} on I with $\sup(\mathcal{F}) = \aleph$.

Definition 1.9. Let n be a nonnegative integer. A right R -module M is called n - (\aleph, Q) -presented if there is a finite n - (\aleph, Q) -presentation of M i.e. an exact sequence

$$0 \rightarrow K_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_1 \rightarrow F_0 \rightarrow M \rightarrow 0$$

in which every F_i is free of finite rank and K_n is (\aleph, Q) -finitely generated.

Definition 1.10. Let n be a nonnegative integer. A ring R is said to be right n - (\aleph, Q) -coherent if every n -presented right R -module is $(n+1)$ - (\aleph, Q) -presented.

Lemma 1.11. Let n be a positive integer, N be an n - (\aleph, Q) -presented right R -module and $\{Q_i; i \in I\}$ be a family of left R -modules from $\text{Gen}(Q)$. Then:

- (i) there is an epimorphism $\text{Tor}_n^R(N, \prod_{i \in I}^{\aleph} Q_i) \rightarrow \prod_{i \in I}^{\aleph} \text{Tor}_n^R(N, Q_i)$;
- (ii) there is an isomorphism $\text{Tor}_{n-1}^R(N, \prod_{i \in I}^{\aleph} Q_i) \cong \prod_{i \in I}^{\aleph} \text{Tor}_{n-1}^R(N, Q_i)$.

PROOF: Let

$$0 \rightarrow K_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_1 \rightarrow F_0 \rightarrow N \rightarrow 0$$

be the finite n - (\aleph, Q) -presentation of N and $K_i = \text{Ker}(F_{i-1} \rightarrow F_{i-2})$ for $i = 2, \dots, n$. Then the short exact sequence $0 \rightarrow K_i \rightarrow F_{i-1} \rightarrow K_{i-1} \rightarrow 0$ induces the commutative diagram

$$\begin{array}{ccccc} 0 \rightarrow \text{Tor}_1^R(K_{i-1}, \prod_{i \in I}^{\aleph} Q_i) & \longrightarrow & K_i \otimes \prod_{i \in I}^{\aleph} Q_i & \longrightarrow & F_{i-1} \otimes \prod_{i \in I}^{\aleph} Q_i \\ & & \downarrow \varphi_{K_i} & & \downarrow \varphi_{F_{i-1}} \\ 0 \rightarrow \prod_{i \in I}^{\aleph} \text{Tor}_1^R(K_{i-1}, Q_i) & \longrightarrow & \prod_{i \in I}^{\aleph} (K_i \otimes Q_i) & \longrightarrow & \prod_{i \in I}^{\aleph} (F_{i-1} \otimes Q_i) \end{array}$$

Then f_{n-1} is an epimorphism since K_n is (\aleph, Q) -finitely generated and f_{n-2} is an isomorphism since K_{n-1} is (\aleph, Q) -finitely presented, K_i being finitely presented for $i < n-1$. Now our lemma follows from the fact that $\text{Tor}_{n-1}^R(N, -) \cong \text{Tor}_1^R(K_{n-2}, -)$ and $\text{Tor}_n^R(N, -) \cong \text{Tor}_1^R(K_{n-1}, -)$. \square

Theorem 1.12. Let n be a nonnegative integer. Then the following conditions are equivalent:

- (i) $\prod_{i \in I}^{\aleph} Q$ is n -flat for every index set I ;
- (ii) $\prod_{i \in I}^{\aleph} Q_i$ is n -flat for every index set I and any family of n -flat modules $Q_i \in \text{Gen}(Q)$;

(iii) R is right n - (\aleph, Q) -coherent.

(iv)

$$\mathrm{Tor}_n^R(N, \prod_{i \in I}^{\aleph} Q_i) \cong \prod_{i \in I}^{\aleph} \mathrm{Tor}_n^R(N, Q_i)$$

for every n -presented right R -module N and any family of left R -modules $Q_i \in \mathrm{Gen}(Q)$.

PROOF: (ii) implies (i). Obvious.

(i) implies (iii). Suppose that N is an n -presented right R -module,

$$F_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_1 \rightarrow F_0 \rightarrow N \rightarrow 0$$

is a finite n -presentation of N and $K_i = \mathrm{Ker}(F_{i-1} \rightarrow F_{i-2})$ for $i = 2, \dots, n$. Then the exact sequence $0 \rightarrow K_n \rightarrow F_{n-1} \rightarrow K_{n-1} \rightarrow 0$ induces the following commutative diagram

$$\begin{array}{ccccc} 0 \rightarrow K_n \otimes \prod_{i \in I}^{\aleph} Q & \longrightarrow & F_{n-1} \otimes \prod_{i \in I}^{\aleph} Q & \longrightarrow & K_{n-1} \otimes \prod_{i \in I}^{\aleph} Q \rightarrow 0 \\ & \downarrow \varphi_{K_n} & \downarrow \varphi_{F_{n-1}} & & \downarrow \varphi_{K_{n-1}} \\ 0 \rightarrow \prod_{i \in I}^{\aleph} (K_n \otimes Q) & \longrightarrow & \prod_{i \in I}^{\aleph} (F_{n-1} \otimes Q) & \longrightarrow & \prod_{i \in I}^{\aleph} (K_{n-1} \otimes Q) \rightarrow 0 \end{array}$$

Then $\mathrm{Tor}_1^R(K_{n-1}, \prod_{i \in I}^{\aleph} Q) \cong \mathrm{Tor}_n^R(N, \prod_{i \in I}^{\aleph} Q) = 0$ by assumption and the upper row is exact. The lower row is exact since Q is flat. Now $\varphi_{F_{n-1}}, \varphi_{K_{n-1}}$ are isomorphisms and consequently φ_{K_n} is an isomorphism. Thus K_n is (\aleph, Q) -finitely presented by Lemma 1.7. Hence N is $(n+1)$ - (\aleph, Q) -presented.

(iii) implies (iv). It follows immediately from Lemma 1.11 (ii).

(iv) implies (ii). Obvious. □

REFERENCES

- [1] Bican L., Kepka T., Nĕmec P., *Rings, Modules and Preradicals*, Marcel Dekker Inc., New York and Basel, 1982.
- [2] Chase S.U., *Direct products of modules*, Trans. Amer. Math. Soc. **97** (1960), 457–473.
- [3] Jianlong Chen, Nanqing Ding, *On n -coherent rings*, Comm. Algebra **24** (10) (1996), 3211–3216.
- [4] Colby R.R., Rutter E.A., Jr., *\prod -flat and \prod -projective modules*, Arch. Math. **22** (1971), 246–251.
- [5] Jirásko J., *Large subdirect products of relative flat modules*, Conf. Abelian Groups and Modules, Padova, 1994.
- [6] Jirásková H., *Generalized flatness and coherence*, Comment. Math. Univ. Carolinae **21** (1980), 293–308.
- [7] Jones M.F., *Coherence relative to a hereditary torsion theory*, Comm. Algebra **10** (1982), 719–739.
- [8] Lenzig H., *Endlich präsentierbare Moduln*, Arch. Math. **20** (1969), 262–266.

- [9] Loustaunau P., *\mathcal{F} -products of injective, flat and projective modules*, Comm. Algebra **18** (1990), 3671–3683.
- [10] Loustaunau P., *Large subdirect products of projective modules*, Comm. Algebra **17** (1989), 197–215.
- [11] Oyonarte L., Torrecillas B., *Large subdirect products of flat modules*, Comm. Algebra **24** (4) (1996), 1389–1407.
- [12] Oyonarte L., Torrecillas B., *Large subdirect products of graded modules*, Comm. Algebra **27** (2) (1999), 681–701.

DEPARTMENT OF MATHEMATICS, FACULTY OF CIVIL ENGINEERING, CZECH TECHNICAL UNIVERSITY, THÁKUROVA 7, 166 29 PRAHA 6, CZECH REPUBLIC

(Received March 26, 1999)