

## On (transfinite) small inductive dimension of products\*

V.A. CHATYRKO, K.L. KOZLOV<sup>†</sup>

*Abstract.* In this paper we study the behavior of the (transfinite) small inductive dimension (*trind*) *ind* on finite products of topological spaces. In particular we essentially improve Toulmin’s estimation [T] of *trind* for Cartesian products.

*Keywords:* transfinite dimension

*Classification:* 54F45

In this paper we study the behavior of the (transfinite) small inductive dimension (*trind*) *ind* on finite products of topological spaces. It is known that if the finite sum theorem for *ind* holds in the factors  $X, Y$  then the inequality

$$(1) \quad \text{ind}(X \times Y) \leq \text{ind} X + \text{ind} Y$$

is true (Pasyukov [9] for completely regular spaces, see also [1] for regular  $T_1$ -spaces). Similar statements for the transfinite small inductive dimension *trind* one can find in [11] (the case of regular  $T_1$ -spaces) and in [2] (the case of normal  $T_1$ -spaces).

But if the finite sum theorem for *ind* fails even in one factor then the inequality (1) is not valid for two compact spaces. Filippov [5] has constructed compact spaces  $X, Y$  such that  $\text{ind} X = \text{Ind} X = \dim X = 1, \text{ind} Y = \text{Ind} Y = \dim Y = 2$  but  $\text{ind}(X \times Y) = 4$  (see also [8]).

In the sequel,  $\alpha = \lambda(\alpha) + n(\alpha)$  is the natural decomposition of the ordinal number  $\alpha$  into the sum of the limit ordinal number  $\lambda(\alpha)$  and the non-negative integer  $n(\alpha) \geq 0$ .

In [10] Toulmin has given the following estimation of the transfinite small inductive dimension for the product of two spaces  $X, Y$  ( $X \times Y$  is hereditarily normal). Namely,

$$(2) \quad \text{trind}(X \times Y) \leq \text{trind} X (+) \text{trind} Y + \psi(n(\text{trind} X), n(\text{trind} Y))$$

where (+) is the natural sum of Hessenberg [6],  $\psi(0, m) = \psi(m, 0) = 0$  if  $m$  is a non-negative integer and  $\psi(n, m) = n + m - 1 + \max\{\psi(n - 1, m), \psi(n, m - 1)\} + \psi(n - 1, m - 1)$  if  $n, m$  are positive integers.

---

\*Supported by Kungliga Vetenskapsakademien, project 12529.

<sup>†</sup>Supported by RFFFI, grant 99-01-00128.

In particular for finite dimensional spaces  $X, Y$  the inequality

$$(3) \quad \text{ind}(X \times Y) \leq \varphi_T(\text{ind } X, \text{ind } Y)$$

is valid, where  $\varphi_T(n, m) = n + m + \psi(n, m)$ ,  $n, m$  are non-negative integers (see Tab. 1).

Observe that formula (2) can be written as follows

$$(2') \quad \text{trind}(X \times Y) \leq \lambda(\text{trind } X)(+) \lambda(\text{trind } Y) + \varphi_T(n(\text{trind } X), n(\text{trind } Y)).$$

In [9] another estimation of the small inductive dimension  $\text{ind}$  has been proved. Namely,

$$(4) \quad \text{ind}(X \times Y) \leq \varphi_P(\text{ind } X, \text{ind } Y),$$

where  $\varphi_P(0, m) = \varphi_P(m, 0) = m$  if  $m$  is a non-negative integer and  $\varphi_P(n, m) = \varphi_P(n - 1, m) + \varphi_P(n, m - 1) + 2$  if  $n, m$  are positive integers (see Tab. 2) ( $X, Y$  are regular).

In this paper we essentially improve the inequalities (2)–(4).

By a space we mean a regular  $T_1$ -space. We let  $BdU$  denote the boundary of the set  $U$ . Our terminology follows [E].

The following lemma is evident.

**Lemma 1.** *Let  $X = X_1 \cup X_2$ , where  $X_i$  is a subset of  $X$ . If  $\text{Int } X_1 \cup \text{Int } X_2 = X$  and  $\text{trind } X_i \leq \alpha_i, i = 1, 2$ , then  $\text{trind } X \leq \max\{\alpha_1, \alpha_2\}$ .*

**Theorem 2.** *Let  $X = X_1 \cup X_2$ , where  $X_i$  is closed in  $X$ , and  $\text{trind } X_i \leq \alpha_i, i = 1, 2$ . Then*

$$\text{trind } X \leq \begin{cases} \max\{\alpha_1, \alpha_2\} & \text{if } \lambda(\alpha_1) \neq \lambda(\alpha_2) \\ \max\{\alpha_1, \alpha_2\} + 1 & \text{if } \lambda(\alpha_1) = \lambda(\alpha_2). \end{cases}$$

*In particular, in the finite-dimensional case we have*

$$\text{ind } X \leq \max\{\text{ind } X_1, \text{ind } X_2\} + 1.$$

PROOF: If  $\lambda(\alpha_1) \neq \lambda(\alpha_2)$  then the inequality is valid due to [4, Theorem 7.2.6]. Let  $\lambda(\alpha_1) = \lambda(\alpha_2)$ . If  $x \in X_1 \setminus X_2$  or  $x \in X_2 \setminus X_1$  then  $\text{trind}_x X \leq \max\{\alpha_1, \alpha_2\}$ . Let now  $x \in X_1 \cap X_2$  and  $A$  be a closed subset of  $X$  such that  $x \notin A$  and  $A \cap X_i \neq \emptyset, i = 1, 2$ . Choose a partition  $C_1$  in  $X_1$  between the point  $x$  and the set  $A \cap X_1$ . Obviously one can choose the partition  $C_1$  with  $\text{trind } C_1 < \alpha_1$ . Let  $X_1 \setminus C_1 = U_1 \cup V_1$ , where  $U_1, V_1$  are open in  $X_1$  and disjoint, and  $x \in U_1, A \cap X_1 \subset V_1$ . Choose a partition  $C_2$  in  $X_2$  between the point  $x$  and the closed set  $((C_1 \cup V_1) \cup A) \cap X_2$ . Obviously one can choose the partition  $C_2$  with  $\text{trind } C_2 < \alpha_2$ . Let  $X_2 \setminus C_2 = U_2 \cup V_2$ , where  $U_2, V_2$  are open in  $X_2$  and disjoint, and  $x \in U_2, ((C_1 \cup V_1) \cup A) \cap X_2 \subset V_2$ . Observe that the space  $Y = C_1 \cup C_2 \cup (X_1 \cap X_2)$  is equal to the union  $Y_1 \cup Y_2$ , where  $Y_i = C_i \cup (X_1 \cap X_2)$  is a subset of  $Y$ . Moreover  $\text{Int } Y_1 \cup \text{Int } Y_2 = Y, \text{trind } Y_i \leq \alpha_i$  (recall that  $Y_i \subset X_i$ ). So by Lemma 1 we have the inequality  $\text{trind } Y \leq \max\{\alpha_1, \alpha_2\}$ . The set  $C = X \setminus (((U_1 \setminus X_2) \cup U_2) \cup (V_1 \cup (V_2 \setminus X_1)))$  is a partition between the point  $x$  and the set  $A$ . Besides  $C \subset Y$ . Hence  $\text{trind } C \leq \max\{\alpha_1, \alpha_2\}$ .  $\square$

**Remark 3.** a) Theorem 2 is similar to [3, Theorem 3.9] in the case of regular  $T_1$ -spaces. The analog of [3, Corollary 3.10] (the finite sum theorem for closed subspaces) in the case of regular  $T_1$ -spaces is also valid.

b) Recall that there exists a compact space  $L$  with  $ind Y = 2$  which can be represented as the union of two closed subspaces  $L_1$  and  $L_2$  such that  $ind L_1 = ind L_2 = 1$  [4, Lokucievskij's example 2.2.14].

c) Recall also that van Douwen and Przymusiński [4, Problem 4.1.B] defined even a metrizable space  $Y$  with  $ind Y = 1$  which can be represented as the union of two closed subspaces  $Y_1$  and  $Y_2$  such that  $ind Y_1 = ind Y_2 = 0$ .

Let  $P = X \times Y$ . Note that for a rectangular open subset  $U \times V$  of  $P$  we have

$$(*) \quad Bd(U \times V) = (Bd(U) \times [V]) \cup ([U] \times Bd(V)).$$

The following lemma is evident.

**Lemma 4.** *Let  $trind X = 0$ . Then  $trind(X \times Y) = trind Y$  for any space  $Y$ .*

Observe that in particular Lemma 4 is also valid for  $ind$ .

Now let us consider the finite-dimensional case.

**Theorem 5.** *Let  $P = X \times Y$ . Then*

$$(5) \quad ind P \leq \varphi_1(ind X, ind Y)$$

where  $\varphi_1(0, m) = \varphi_1(m, 0) = m$  if  $m$  is a non-negative integer,  $\varphi_1(n, m) = 2(n + m) - 1$  if  $n, m$  are positive integers (see Tab. 3, observe that  $\varphi_1(n, m) = \max\{\varphi_1(n - 1, m), \varphi_1(n, m - 1)\} + 2$  if  $n, m \geq 1$ ).

PROOF: If at least one of the factors is zero-dimensional in the sense of  $ind$  then the inequality holds due to Lemma 4. Suppose that  $ind X, ind Y \geq 1$ . Apply an induction on the sum  $ind X + ind Y = k, k \geq 2$ .

Let  $k = 2$ . Then for any point  $p \in P$  and its any neighborhood  $W$  there is a rectangular neighborhood  $U \times V \subset W$  of this point with  $ind BdU \leq 0, ind BdV \leq 0$ .

By Lemma 4 each element from the right part of equality (\*) is not more than one-dimensional. From Theorem 2 it follows that  $ind Bd(U \times V) \leq 2$ . Hence formula (5) is valid.

Let the theorem hold for  $k < n, n \geq 3$ . Put  $k = n$ . For any point  $p \in P$  and its any neighborhood  $W$  there is a rectangular neighborhood  $U \times V \subset W$  of this point with  $ind BdU \leq ind X - 1, ind BdV \leq ind Y - 1$ . By induction assumption the small inductive dimension of each element from the right part of equality (\*) is not more than  $2(n - 1) - 1$ . From Theorem 2 it follows that  $ind Bd(U \times V) \leq 2(n - 1)$ . Hence  $ind P \leq 2(n - 1) + 1 = 2(ind X + ind Y) - 1$ .  $\square$

Using induction one can easily obtain the following

**Estimations.**

- (a)  $\psi(n, m) \leq \psi(n + 1, m), \psi(n, m) \leq \psi(n, m + 1)$ ;
- (b)  $\varphi_1(n, m) \leq \varphi_T(n, m) \leq \varphi_P(n, m)$ , if  $n, m \geq 1$  and if at least one of the numbers is  $> 1$  then both inequalities are strict.

**Remark 6.** It is easy to see that  $\psi(n, n) \geq 2n - 1 + 2\psi(n - 1, n - 1), n \geq 1$ . Moreover, if  $n > k$  then  $\psi(n, n) \geq 2(2^k - 1)n + 2^k\psi(n - k, n - k) + f(k)$ . Hence, for every natural number  $m$  the inequality  $\varphi_T(n, n) \geq mn$  holds for large  $n$ .

Estimation from Theorem 5 can be improved for the class of completely paracompact spaces.

Let us recall [12] that a topological space  $X$  is *completely paracompact* if, for any open cover  $\lambda$  of  $X$ , there exist open star-finite covers  $\mu_i$  of  $X, i \in \mathbb{N}$ , such that, for any  $x \in X$  there exist  $O \in \lambda, i \in \mathbb{N}$  and  $V \in \mu_i$  for which  $x \in V \subset O$ .

It is known ([12]) that:

- (a) any  $F_\sigma$  subset of a completely paracompact space is completely paracompact;
- (b) any regular completely paracompact space is paracompact and any strongly paracompact space is completely paracompact;
- (c)  $\dim X \leq \text{ind} X$  for any completely paracompact space.

**Lemma 7.** Let  $Z$  be a completely paracompact space and  $Z = Z_1 \cup Z_2$ , where  $Z_i$  is closed,  $\text{ind} Z_i \leq 1, i = 1, 2$ , and  $\text{ind}(Z_1 \cap Z_2) \leq 0$ . Then  $\text{ind} Z \leq 1$ .

PROOF: If  $x \in Z_1 \setminus Z_2$  or  $x \in Z_2 \setminus Z_1$  then  $\text{ind}_x Z \leq 1$ . Let now  $x \in Z_1 \cap Z_2$  and  $A$  be a closed subset of  $Z$  such that  $x \notin A$ . Then from the proof of Theorem 2 it follows that there exists a partition  $C$  between  $x$  and  $A$  such that  $C \subset Y = (Z_1 \cap Z_2) \cup C_1 \cup C_2$ , where  $\text{ind} C_i \leq 0, i = 1, 2$ . By property (c) and the finite sum theorem for  $\dim$  it follows that  $\dim Y \leq 0$ . From (b) it follows that  $\text{ind} Y \leq 0$ . Hence  $\text{ind} Z \leq 1$ . □

**Theorem 8.** Let  $P = X \times Y$  be completely paracompact. Then

$$(6) \quad \text{ind} P \leq \varphi_2(\text{ind} X, \text{ind} Y),$$

where  $\varphi_2(0, m) = \varphi_2(m, 0) = m$  if  $m$  is a non-negative integer,  $\varphi_2(n, m) = 2(n + m) - 2$  if  $n, m$  are positive integers (see Tab. 4, observe that  $\varphi_2(n, m) = \max\{\varphi_2(n - 1, m), \varphi_2(n, m - 1)\} + 2$  if  $n, m \geq 1$  and  $(n, m) \neq (1, 1)$ ).

PROOF: If at least one of the factors is zero-dimensional in the sense of  $\text{ind}$  then the inequality holds due to Lemma 4. Suppose that  $\text{ind} X, \text{ind} Y \geq 1$ . Apply an induction on the sum  $\text{ind} X + \text{ind} Y = k, k \geq 2$ .

Let  $k = 2$ . Then for any point  $p \in P$  and its any neighborhood  $W$  there is a rectangular neighborhood  $U \times V \subset W$  of this point with  $\text{ind} BdU \leq 0, \text{ind} BdV \leq 0$ .

Put  $Z = Bd(U \times V), Z_1 = Bd(U) \times [V], Z_2 = [U] \times Bd(V)$  then  $Z = Z_1 \cup Z_2, Z_1 \cap Z_2 = Bd(U) \times Bd(V)$ . By Lemma 7 and property (a) we have  $\text{ind} Z \leq 1$ . Hence

formula (6) is valid.

Let the theorem hold for  $k < n, n \geq 3$ . Put  $k = n$ . For any point  $p \in P$  and any its neighborhood  $W$  there is a rectangular neighborhood  $U \times V \subset W$  of this point with  $ind BdU \leq ind X - 1, ind BdV \leq ind Y - 1$ . By induction assumption the small inductive dimension of each element from the right part of equality (\*) is not more than  $2(n - 1) - 2$ . From Theorem 2 it follows that  $ind Bd(U \times V) \leq 2(n - 1) - 1$ . Hence  $ind P \leq 2(n - 1) = 2(ind X + ind Y) - 2$ .  $\square$

**Corollary 9.** *Let  $P = X \times Y$ , where  $X, Y$  are compact spaces, and  $ind X, ind Y \geq 1$ . Then*

$$(7) \quad ind P \leq 2(ind X + ind Y) - 2.$$

Observe that estimation (7) is exact (i.e. it cannot be improved) for  $ind X = ind Y = 1$  (it is evident) and for  $ind X = 1, ind Y = 2$  (the named earlier Filippov's result [5]).

**Question A.** Is estimation (7) exact for all situations?

**Question B.** Are there spaces  $X, Y$  such that  $ind X = ind Y = 1$  and  $ind X \times Y = 3$ ?

**Remark 10.** Let  $P = \prod_{i=1}^n X_i$ , where  $X_i$  is a compact space with  $ind X_i \geq 1, i = 1, \dots, n$ . Then  $ind P \leq n(\sum_{i=1}^n ind X_i - n + 1)$ . In the case when all spaces are one-dimensional in the sense of *ind* the formula coincides with Lifanov's result [7].

Now let us consider the transfinite case.

**Theorem 11.** *Let  $P = X \times Y$  and  $trind X \leq \alpha, trind Y \leq \beta$ . Then*

$$(8) \quad trind P \leq \begin{cases} \alpha(+) \beta + n(\alpha) + n(\beta) - 1 & \text{if } n(\alpha), n(\beta) \geq 1; \\ \alpha(+) \beta & \text{otherwise.} \end{cases}$$

(Observe that formula (8) can be written as follows

$$(8') \quad trind(X \times Y) \leq \lambda(\alpha)(+) \lambda(\beta) + \varphi_1(n(\alpha), n(\beta)). \quad )$$

**PROOF:** Use induction on  $\alpha(+) \beta = \gamma$ . If  $\gamma < \omega$  then the inequality holds due to Theorem 5.

Let the theorem be valid for  $\gamma < \nu \geq \omega$ . Put  $\gamma = \nu$ . Then for any point  $p \in P$  and its any neighborhood  $W$  there is a rectangular neighbourhood  $U \times V \subset W$  of this point with  $trind BdU < \alpha, trind BdV < \beta$ .

If  $\nu$  is limit then  $\nu = \lambda(\nu)$  and  $\lambda(\alpha) = \alpha, \lambda(\beta) = \beta$ . We can assume that  $\lambda(\alpha) \geq \omega$  and  $\lambda(\beta) \geq \omega$  (otherwise apply Lemma 4). By induction assumption



Tab 3.,  $\varphi_1(n, m)$  :

	0	1	2	3	...	n
0	0	1	2	3	...	
1	1	3	5	7	...	
2	2	5	7	9	...	
3	3	7	9	11	...	
...	...	...	...	...	...	
m						

Tab 4.,  $\varphi_2(n, m)$  :

	0	1	2	3	...	n
0	0	1	2	3	...	
1	1	2	4	6	...	
2	2	4	6	8	...	
3	3	6	8	10	...	
...	...	...	...	...	...	
m						

REFERENCES

[1] Basmanov V.N., *On inductive dimensions of products of spaces*, Moscow Univ. Math. Bull. **36** (1981), no. 1, 19–22.  
 [2] Chatyrko V.A., *Ordinal products of topological spaces*, Fund. Math. **144** (1994), 95–117.  
 [3] Chatyrko V.A., *On finite sum theorems for transfinite inductive dimensions*, Fund. Math., to appear.  
 [4] Engelking R., *Theory of Dimensions, Finite and Infinite*, Heldermann Verlag, Lemgo, 1995.  
 [5] Filippov V.V., *On the inductive dimension of the product of bicomacta*, Soviet Math. Dokl. **13** (1972), 250–254.  
 [6] Hessenberg G., *Grundbegriffe der Mengenlehre*, Göttingen, 1906.  
 [7] Lifanov I.K., *On the dimension of the product of one-dimensional bicomacta*, Soviet Math. Dokl. **9** (1968), 648–651.  
 [8] Malyhin D.V., *Some properties of topological products* (in Russian), Moscow State University, Dissertation, 1999.  
 [9] Pasynkov B.A., *On inductive dimensions*, Soviet Math. Dokl. **10** (1969), 1402–1406.  
 [10] Toulmin G.H., *Shuffling ordinals and transfinite dimension*, Proc. London Math. Soc. **4** (1954), 177–195.  
 [11] Vinogradov R.N., *On transfinite dimension*, Moscow Univ. Math. Bull. **48** (1993), no. 5, 15–17.  
 [12] Zarelua A.V., *On a theorem of Hurewicz*, Amer. Math. Soc. Transl. **55** (1966), no. 2, 141–152.

DEPARTMENT OF MATHEMATICS, LINKÖPING UNIVERSITY, S-581 83 LINKÖPING, SWEDEN

E-mail: vitja@mai.liu.se

DEPARTMENT OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY, 119899 MOSCOW, RUSSIA

E-mail: kkozlov@nw.math.msu.su

(Received May 21, 1999)