

## Biharmonic Green domains in a Riemannian manifold

S.I. OTHMAN, V. ANANDAM

*Abstract.* Let  $R$  be a Riemannian manifold without a biharmonic Green function defined on it and  $\Omega$  a domain in  $R$ . A necessary and sufficient condition is given for the existence of a biharmonic Green function on  $\Omega$ .

*Keywords:* biharmonic Green functions

*Classification:* 31C12, 31B30

### 1. Introduction

In a Riemannian manifold  $R$ , we say that a domain  $\Omega$  is a biharmonic Green domain if there exists a positive solution  $Q_y(x)$  for the equation  $\Delta^2 Q_y(x) = \delta_y(x)$  in  $\Omega$ , where  $y$  is some point in  $\Omega$  and  $\Delta$  is the Laplace-Beltrami operator in  $R$ . Some necessary and sufficient conditions for  $R$  to be a biharmonic Green space are given in Sario et al. [8, Chapter VIII]. In this note we give a necessary and sufficient condition for a domain  $\Omega$  in  $R$  to be a biharmonic Green domain when  $R$  itself is not a biharmonic Green space.

### 2. Preliminaries

Let  $R$  be an oriented Riemannian manifold of dimension  $n \geq 2$  with local parameters  $x = (x^1, \dots, x^n)$  and a  $C^\infty$  metric tensor  $g_{ij}$  such that  $g_{ij}x^i x^j$  is positive definite. If  $D$  is the determinant of  $g_{ij}$ , denote the volume element by  $dx = D^{\frac{1}{2}} dx^1 \dots dx^n$ ;  $\Delta = d\delta + \delta d$  is the Laplace-Beltrami operator acting on  $R$  in the sense of distributions; in the Euclidean case,  $\Delta$  reduces to the form  $\Delta u = -\sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2}$ . A continuous function  $h$  on an open set is harmonic, by definition, if  $\Delta h = 0$ . To every open set  $w$  in  $R$ , let  $H(w)$  denote the class of harmonic functions on  $w$ . Then these harmonic functions have the *sheaf property*, solve locally the *Dirichlet problem* and possess the *Harnack property*; that is, they satisfy the axioms 1, 2, 3 of Brelot in the axiomatic potential theory ([5, pp. 13–14]). Consequently, we can use all the notions and the results of this axiomatic theory in the context of a Riemannian manifold; some of these are the following:

- (1) Let  $w$  be a regular open set in  $R$ , that is  $w$  is relatively compact in  $R$  and each boundary point of  $w$  is regular for the Dirichlet problem. A compact set  $k$  in  $w$  is said to be *outerregular* if  $w \setminus k$  is a regular open set. Given

a compact set  $k$  and a domain  $w$ , one can construct a regular domain  $w_0$  and an outerregular compact set  $k_0$  such that  $k \subset \overset{\circ}{k_0} \subset k_0 \subset w_0 \subset w$  (see Loeb [6]).

- (2) (See [5, pp. 37, 38 and 47]). If  $s > 0$  is a superharmonic function on a domain  $\Omega \subset R$  and if  $e$  is a subset of  $\Omega$ , the *reduced function* by definition is

$$R_s^e(x) = \inf\{t(x) : t \geq 0 \text{ superharmonic on } \Omega \text{ and } t \geq s \text{ on } e\};$$

and its l.s.c. regularization is the *balayage*  $\widehat{R}_s^e(x) = \liminf_{y \rightarrow x} R_s^e(y)$ . In a domain  $\Omega$  with a positive potential, a set  $e$  is *polar* if and only if  $R_1^e(x) = 0$  at some point  $x$ , or, equivalently  $\widehat{R}_1^e \equiv 0$ .

- (3) If there is a positive potential on  $\Omega$ , we define on  $\Omega$  the Green function  $G(x, y) = G_y(x)$  with pole  $y \in \Omega$ , so that  $\Delta G_y = \delta_y$ . Then for any potential  $p$  on  $\Omega$ ,  $\Delta p = \mu$  is a Radon measure and  $p(x) = \int_{\Omega} G(x, y) d\mu(y)$ . Also it is proved in [7] that given a Radon measure  $\mu \geq 0$  on  $\Omega$ ,  $\int_{\Omega} G(x, y) d\mu(y)$  is a potential if and only if

$$\int_{\Omega} \widehat{R}_1^w(x) d\mu(x) < \infty \text{ for some nonempty open set } w \text{ in } \Omega.$$

- (4) More generally, we have the following result in [3]: Let  $\Omega$  be a domain in  $R$  with or without positive potentials. Let  $\mu \geq 0$  be a Radon measure on  $\Omega$ . Then there exists a superharmonic function  $s$  on  $\Omega$  such that  $\Delta s = \mu$ . This result is in fact a simple generalization of a classical result of Brelot [4] in  $R^n$ .

**Lemma 2.1.** *Let  $\Omega$  be a domain in  $R$  such that  $\Omega$  has the Green function  $G(x, y)$  defined on it. Then for a Radon measure  $\mu \geq 0$  on  $\Omega$ ,  $\int_{\Omega} G(x, y) d\mu(y)$  is a potential if and only if for one (and hence any) nonpolar compact set  $k$  in  $\Omega$ ,  $\int_{\Omega} R_1^k d\mu < \infty$ .*

PROOF: This is a more useful reformulation of Theorem 3.1 [7]. First note that  $R_1^k$  is  $\mu$ -measurable. For  $R_1^k = \inf_n R_1^{w_n}$  where  $w_n$  is a decreasing sequence of relatively compact open sets such that  $k = \bigcap w_n$ . Since each  $R_1^{w_n} = \widehat{R}_1^{w_n}$  is l.s.c., it is  $\mu$ -measurable and hence  $R_1^k$  is  $\mu$ -measurable.

- (1) Suppose  $\int_{\Omega} G(x, y) d\mu(y)$  is a potential on  $\Omega$  and  $k$  is a nonpolar compact set in  $\Omega$ . Then for some  $x_0 \in k$ ,  $\int_{\Omega} G(x_0, y) d\mu(y) < \infty$ . If  $G(x_0, y) \geq a$  on  $k$ ,  $G(x_0, y) \geq aR_1^k$  on  $\Omega$  and hence  $\int_{\Omega} R_1^k d\mu < \infty$ .
- (2) Conversely, suppose  $\int_{\Omega} R_1^k d\mu < \infty$  for some nonpolar compact set  $k$ . Since  $R_1^k = \inf_n R_1^{w_n}$ , we can find an open set  $w$  and an outerregular compact set  $A$  such that  $k \subset \overset{\circ}{A} \subset A \subset w$  and  $\int_{\Omega} R_1^w d\mu < \infty$ . Now  $p(x) =$

$\int_A G(x, y) d\mu(y)$  is a potential on  $\Omega$ ; hence  $p(x_0) < \infty$  for some  $x_0 \in k$ . If  $a \leq G(x_0, y) \leq b$  on  $\partial A$ , then  $aR_1^k(y) \leq G(x_0, y) \leq bR_1^A(y)$  on  $\Omega \setminus A$  and hence  $\int_{\Omega \setminus A} G(x_0, y) d\mu(y)$  is finite, which implies that  $\int_{\Omega} G(x_0, y) d\mu(y)$  is finite and hence  $\int_{\Omega} G(x, y) d\mu(y)$  is a potential on  $\Omega$ . □

The following form of Lemma 2.1, without an explicit reference to the reduced functions, is convenient for applications.

**Lemma 2.2.** *Let  $\Omega$  be a domain in  $R$  with the Green function  $G(x, y)$  defined on it;  $\mu \geq 0$  is a Radon measure on  $\Omega$ . Then the following are equivalent:*

- (1) *There exists a superharmonic function  $s > 0$  on  $\Omega$  such that  $\int_{\Omega} s d\mu < \infty$ .*
- (2)  *$p(x) = \int_{\Omega} G(x, y) d\mu(y)$  is a potential on  $\Omega$ .*
- (3) *For any locally bounded potential  $q(x)$  with compact harmonic support on  $\Omega$ ,  $\int_{\Omega} q d\mu < \infty$ .*

PROOF: (1)  $\Rightarrow$  (2): Let  $k$  be a nonpolar compact subset of  $\Omega$ . If  $s \geq \alpha > 0$  on  $k$ , then  $\alpha R_1^k \leq s$  on  $\Omega$  and hence  $\int_{\Omega} R_1^k d\mu \leq \frac{1}{\alpha} \int_{\Omega} s d\mu < \infty$ . Hence by Lemma 2.1,  $p(x) = \int_{\Omega} G(x, y) d\mu(y)$  is a potential on  $\Omega$ .

(2)  $\Rightarrow$  (3): Let  $q$  be a locally bounded potential on  $\Omega$ , with compact harmonic support  $A$ . Let  $k$  be an outerregular compact set such that  $A \subset \overset{\circ}{k}$ . Then  $R_q^k = q$  on  $\Omega \setminus k$ . For,  $\widehat{R}_q^k \leq q$  on  $\Omega$  and hence  $t = q - \widehat{R}_q^k$  on  $\Omega \setminus k$  extended by 0 on  $k$  is a positive subharmonic function less or equal to  $q$  on  $\Omega$ ; hence  $t \leq 0$ , so that  $q = \widehat{R}_q^k = R_q^k$  on  $\Omega \setminus k$ . Consequently, if  $q \leq \alpha$  on  $k$ , then  $q \leq \alpha R_1^k$  on  $k$ ; also on  $\Omega \setminus k$ ,  $q = R_q^k \leq R_{\alpha}^k = \alpha R_1^k$ . Thus  $q \leq \alpha R_1^k$  on  $\Omega$ . Now assumption (2) along with Lemma 2.1 shows that  $\int_{\Omega} R_1^k d\mu < \infty$ . Hence  $\int_{\Omega} q d\mu < \infty$ .

(3)  $\Rightarrow$  (1): Let  $k$  be a nonpolar compact set. Let  $s = \widehat{R}_1^k$  on  $\Omega$ . Then  $s > 0$  is a superharmonic function that is bounded on  $\Omega$  and has compact harmonic support. Hence by (3),  $\int_{\Omega} s d\mu < \infty$ . □

### 3. Biharmonic Green domains

Let  $\Omega$  be a domain in  $R$ . Given  $y \in \Omega$ , let  $w$  be a regular domain for the Dirichlet problem such that  $y \in w \subset \overline{w} \subset \Omega$ . Let  $v_w(x, y)$  be the biharmonic Green function on  $w$  with biharmonic singularity  $y$ , that is  $\Delta^2 v_w(x, y) = \delta_y(x)$ , and with boundary conditions  $v_w/\partial w = 0$  and  $\Delta v_w/\partial w = 0$ . Then  $v_w$  increases with  $w$ . Write  $v_{\Omega}(x, y) = \lim_{w \rightarrow \Omega} v_w(x, y)$  if the limit exists for some regular exhaustion  $\{w\}$ .  $v_{\Omega}(x, y)$  is called the *biharmonic Green function* on  $\Omega$  and its existence is independent of the regular exhaustion  $\{w\}$  and the choice of the singular point  $y$  (see Sairo et al. [8, pp. 300–307]). When  $v_{\Omega}(x, y)$  exists on  $\Omega$ , it can be written as  $v_{\Omega}(x, y) = \int_{\Omega} G(x, z)G(z, y) dz$ .

**Definition 3.1.** A domain  $\Omega$  in  $R$  is said to be a *biharmonic Green domain* if and only if the biharmonic Green function  $v_\Omega(x, y)$  exists on  $\Omega$ .

The following theorem is a collection of known results about  $v_\Omega(x, y)$ .

**Theorem 3.2.** Let  $\Omega$  be a domain in  $R$ , carrying the harmonic Green function  $G(x, y)$ . Then the following are equivalent:

- (1)  $\Omega$  is a biharmonic Green domain.
- (2) For one (and hence any)  $y \in \Omega$ , there exists a potential  $q_y(x)$  on  $\Omega$  such that  $\Delta^2 q_y = \delta_y$ .
- (3) There exists a potential  $Q(x) > 0$  on  $\Omega$  such that  $\Delta Q(x)$  is a superharmonic function.
- (4) There exist potentials  $p$  and  $q$  on  $\Omega$  such that  $\Delta q = p$ . ( $q$  is called a *bipotential*.)

PROOF: (1)  $\Rightarrow$  (2): Let  $v(x, y)$  be the biharmonic Green function on  $\Omega$ . Since  $v(x, y) = \int_\Omega G(x, z)G(z, y) dz$ , for fixed  $y$ ,  $v_y(x) = v(x, y)$  is a potential on  $\Omega$  and  $\Delta v_y(x) = G_y(x)$  (see [8, p. 300]); hence  $\Delta^2 v_y = \delta_y$ .

(2)  $\Rightarrow$  (3): For some potential  $q$  on  $\Omega$ , let  $\Delta^2 q = \delta_y$ . Since  $\Delta^2 q = \Delta G_y$ ,  $\Delta q(x) = G_y(x) +$  (a harmonic function) on  $\Omega$ . That is,  $\Delta q = s$  is a superharmonic function on  $\Omega$ ; note that  $s > 0$  since  $q$  is a potential  $> 0$ .

(3)  $\Rightarrow$  (4): See Theorem 3.2 in [1].

(4)  $\Rightarrow$  (1): This is a consequence of Theorem 4.2 in [1]. □

**Theorem 3.3.** A domain  $\Omega$  in  $R$  is a biharmonic Green domain if and only if there exists a superharmonic function  $s > 0$  on  $\Omega$  such that  $\int_\Omega s^2 dx < \infty$ .

PROOF: (1) Let  $\Omega$  be a biharmonic Green domain. Then there exist potentials  $p > 0$  and  $q > 0$  on  $\Omega$  such that  $\Delta q = p$ . This means that if  $G(x, y)$  is the Green function on  $\Omega$  with  $\Delta G_y = \delta_y$ ,  $q(x) = \int_\Omega G(x, y)p(y) dy$  since  $q$  is a potential with the associated measure  $d\mu(x) = (\Delta q)dx = p dx$  in the Riesz representation. This implies (by Lemma 2.1) that for any nonpolar compact set  $k$  in  $\Omega$ ,  $\int_\Omega R_1^k(y)p(y) dy < \infty$ . Moreover, since  $p$  is a potential on  $\Omega$ , for some  $\lambda > 0$ ,  $R_1^k \leq \lambda p$  on  $\Omega$ . Consequently, with  $s = \widehat{R}_1^k$  we have  $\int_\Omega s^2 dx < \infty$ .

(2) Conversely, let  $s > 0$  be superharmonic on  $\Omega$  such that  $\int_\Omega s^2 dx < \infty$ . Since for a nonpolar compact  $k$  in  $\Omega$ ,  $R_1^k \leq \lambda s$  for some  $\lambda > 0$ ,  $\int_\Omega R_1^k(y)\widehat{R}_1^k(y) dy < \infty$ . This implies (Lemma 2.1) that  $q(x) = \int_\Omega G(x, y)\widehat{R}_1^k(y) dy$  is a potential on  $\Omega$  so that  $\Delta q = \widehat{R}_1^k$ . Since  $\widehat{R}_1^k$  is a potential on  $\Omega$ , we conclude that  $\Omega$  is a biharmonic Green domain. □

**Corollary 1.** Any domain in  $\mathbb{R}^n$ ,  $n \geq 5$ , is a biharmonic Green domain; and  $\mathbb{R}^n$  for  $2 \leq n \leq 4$  is not a biharmonic Green space. (Sario et al. [8, pp. 300–302] and [2, Theorem 5.5]).

PROOF: (a) Let  $\Omega$  be a domain  $\mathbb{R}^n$ ,  $n \geq 5$ . Note that  $s(x) = |x|^{2-n}$  is a positive superharmonic function such that  $\int_{\Omega} s^2 dx \leq \infty$ . Hence  $\Omega$  is a biharmonic Green domain.

(b) Suppose  $\mathbb{R}^n$ ,  $2 \leq n \leq 4$ , is a biharmonic Green space. Then there exists a superharmonic function  $s > 0$  in  $\mathbb{R}^n$  such that  $\int_{\mathbb{R}^n} s^2 dx < \infty$ . If  $B$  is the closed unit ball in  $\mathbb{R}^n$ , then for some  $\lambda > 0$ ,  $R_1^B \leq \lambda s$  and hence  $\int_{\mathbb{R}^n} (R_1^B)^2 dx < \infty$ . But  $R_1^B = |x|^{2-n}$  on  $\mathbb{R}^n \setminus B$ . Hence we should have  $\int_1^\infty \int_{\partial B} r^{4-2n} r^{n-1} dr dw$  is finite, that is,  $\int_1^\infty r^{3-n} dr$  is finite, a contradiction when  $2 \leq n \leq 4$ . □

**Corollary 2.** *Suppose the Riemannian manifold  $R$  is not a biharmonic Green space. If  $\Omega$  is a biharmonic Green domain in  $R$ , then  $e = R \setminus \Omega$  is not compact.*

PROOF: Suppose  $e$  is compact. Let  $k$  be an outerregular compact set such that  $e \subset \overset{\circ}{k} \subset k$ . Since  $\Omega$  is a biharmonic Green domain there exists  $s > 0$  superharmonic on  $\Omega$  such that  $\int_{\Omega} s^2 dx < \infty$ . Suppose  $\inf_{\partial k} s(x) = \lambda$ . Then  $\lambda \widehat{R}_1^k \leq s$  on  $\Omega \setminus k = R \setminus k$  and hence  $\int_{\Omega \setminus k} (\widehat{R}_1^k)^2 dx < \infty$ ; also  $\int_k (\widehat{R}_1^k)^2 dx < \infty$ , and hence  $\int_R (\widehat{R}_1^k)^2 dx < \infty$ . This means that  $R$  is a biharmonic Green space, contradicting the hypothesis. □

**Corollary 3** ([2, Theorem 5.4]). *If  $\Omega$  is a biharmonic Green domain in  $\mathbb{R}^n$ ,  $2 \leq n \leq 4$ , then  $e = \mathbb{R}^n \setminus \Omega$  is neither locally polar nor compact.*

PROOF: (a) Since  $\mathbb{R}^n$ ,  $2 \leq n \leq 4$ , is not a biharmonic Green space, by the above corollary,  $e$  is not compact.

(b) Suppose  $e$  is locally polar. Since  $\Omega$  is a biharmonic Green domain, there exists a superharmonic function  $s > 0$  such that  $\int_{\Omega} s^2 dx < \infty$ . Now  $e = \mathbb{R}^n \setminus \Omega$  being locally polar by the assumption,  $\int_e dx = 0$  and  $s$  extends as a superharmonic function  $u > 0$  on  $\mathbb{R}^n$ . Hence  $\int_{\mathbb{R}^n} u^2 dx < \infty$  which means that  $\mathbb{R}^n$ ,  $2 \leq n \leq 4$ , is a biharmonic Green space, a contradiction. □

#### 4. Biharmonic potentials and quasiharmonic potentials

If there exists a nonconstant positive harmonic function on  $\Omega$ , then we can define the harmonic Green function  $G(x, y)$  on  $\Omega$ . However, we know that this sufficient condition for the existence of the harmonic Green function is not a necessary condition, as for example in  $\mathbb{R}^n$ ,  $n \geq 3$ . A corresponding result for the biharmonic Green function is the following:

**Proposition 4.1.** *Suppose that there exists a biharmonic function which is a positive potential on  $\Omega$ . Then  $\Omega$  is a biharmonic Green domain.*

PROOF: Let  $b$  be a biharmonic function which is a positive potential on  $\Omega$ . Since  $b$  is a potential such that  $\Delta b$  is harmonic, by Theorem 3.2(3),  $\Omega$  is a biharmonic Green domain. □

In view of the above proposition, we propose the following terminology.

**Definition 4.2.** In a domain  $\Omega$  in  $R$ , let  $u > 0$  be a potential.

- (1)  $u$  is said to be a *biharmonic potential* if and only if  $\Delta^2 u = 0$  on  $\Omega$ .
- (2)  $u$  is said to be a *quasiharmonic potential* if and only if  $\Delta u = 1$  on  $\Omega$ .

**Remark.** Let  $\Omega$  be a harmonic Green domain in  $R$ . Then there exists a quasiharmonic potential on  $\Omega$  if and only if  $p(x) = \int_{\Omega} G(x, y) dy$  is a potential on  $\Omega$ . For, suppose  $p(x)$  is a potential. Then  $\Delta p = 1$  so that  $p(x)$  is a quasiharmonic potential on  $\Omega$ . Conversely, suppose  $q$  is a quasiharmonic potential on  $\Omega$ . Since  $q$  is a potential and  $\Delta q = 1$ ,  $q(x) = \int_{\Omega} G(x, y) \Delta q(y) dy = \int_{\Omega} G(x, y) dy$ .

**Theorem 4.3.** Let  $\Omega$  be a harmonic Green domain in  $R$ . Then there exists a biharmonic (resp. quasiharmonic) potential on  $\Omega$  if and only if there are a superharmonic function  $s > 0$  and a harmonic function  $h > 0$  on  $\Omega$  such that  $\int_{\Omega} s(x)h(x) dx < \infty$  (resp.  $\int_{\Omega} s(x) dx < \infty$ ).

PROOF: Let  $G(x, y)$  be the Green function on  $\Omega$ . By Lemma 2.2,  $\int_{\Omega} s(x)h(x) dx$  (resp.  $\int_{\Omega} s(x) dx$ ) is finite if and only if  $Q(x) = \int_{\Omega} G(x, y)h(y) dy$  (resp.  $Q(x) = \int_{\Omega} G(x, y) dy$ ) is a potential on  $\Omega$  which is equivalent to saying that  $Q$  is a biharmonic (resp. quasiharmonic) potential on  $\Omega$ , since  $\Delta Q = h$  (resp.  $\Delta Q = 1$ ).

□

**Corollary 1.** Let  $\Omega$  be a domain in  $R$ . If there exists a quasiharmonic potential on  $\Omega$ , then for any potential  $p$  on  $\Omega$  with compact harmonic support,  $\int_{\Omega} p dx < \infty$ . Consequently, there exists a unique bipotential  $q$  on  $\Omega$  such that  $\Delta q = p$ .

PROOF: Since  $\Omega$  has a quasiharmonic potential, there exists a superharmonic function  $s > 0$  such that  $\int_{\Omega} s dx < \infty$ . Let  $p$  be a potential with compact harmonic support  $k$ . Let  $A$  be an outerregular compact set such that  $k \subset \overset{\circ}{A} \subset A$ . Then  $p = B_{AP}$  on  $\Omega \setminus A$  where  $B_{AP}$  denotes the Dirichlet solution with boundary values  $p$  on  $\partial A$  and 0 at infinity. Hence  $p \leq \lambda s$  on  $\Omega \setminus A$  for some  $\lambda > 0$  so that  $\int_{\Omega \setminus A} p dx < \infty$ . Since  $p$  is locally integrable on  $\Omega$ ,  $\int_A p dx < \infty$ . Hence  $\int_{\Omega} p dx < \infty$ . Consequently, for a nonpolar compact  $k$ ,  $\int_{\Omega} R_1^k(x)p(x) dx < \infty$ . Hence  $q(x) = \int_{\Omega} G(x, y)p(y) dy$  is a potential on  $\Omega$  such that  $\Delta q = p$  (Lemma 2.1). If  $q_1$  is another bipotential on  $\Omega$  such that  $\Delta q_1 = p$ , then  $q_1 = q +$  (a harmonic function  $h$ ) on  $\Omega$ . Note  $h \equiv 0$  by the uniqueness of the Riesz representation. □

**Corollary 2.** Let  $\Omega$  be a domain in  $\mathbb{R}^n$ ,  $n \geq 2$ , such that  $\mathbb{R}^n \setminus \Omega$  is compact. If  $u \geq 0$  is superharmonic on  $\Omega$  and if  $\Delta u$  is constant, then  $u$  is harmonic and hence is of the form

$$u(x) = \begin{cases} \alpha \log |x - a| + b(x) & \text{if } n = 2, a \notin \Omega \\ \alpha + b(x) & \text{if } n \geq 3, \end{cases}$$

where  $\alpha \geq 0$  and  $b(x)$  is harmonic on  $\Omega$  such that  $|b(x)| \leq \beta|x|^{2-n}$  near infinity.

PROOF: First we note that there is no quasiharmonic potential on  $\Omega$ . For, suppose  $\Omega$  has a quasiharmonic potential. Then there exists a superharmonic function

$s > 0$  on  $\Omega$  such that  $\int_{\Omega} s \, dx < \infty$ . Suppose  $\mathbb{R}^n \setminus \Omega = e \subset \{x : |x| < a\}$ . Let  $\lambda = \inf_{|x|=a} s(x)$ . Then  $s(x) \geq \lambda \left|\frac{x}{a}\right|^{2-n}$  on  $|x| > a$  by the minimum principle and hence  $\int_a^\infty \int_{\partial B} \lambda \left(\frac{r}{a}\right)^{2-n} r^{n-1} \, dr \, dw \leq \int_{\Omega} s(x) \, dx < \infty$ . This implies that  $\int_a^\infty r \, dr < \infty$ , a contradiction.

Now write  $u = p + h$  on  $\Omega$  where  $p$  is a potential and  $h$  is harmonic on  $\Omega$ . Since  $\Delta p = \Delta u$  is constant and since there is no quasiharmonic potential on  $\Omega$ ,  $p \equiv 0$ . Hence  $u$  is harmonic  $\geq 0$  outside a compact set. Then, applying an inversion in the unit ball to the classical representation of Bocher's, we get the stated expression for  $u$ .  $\square$

**Remarks.** (1) The above corollary implies that if a positive superharmonic function  $u$  on  $\mathbb{R}^n$ ,  $n \geq 3$ , is biharmonic, then  $u$  is constant. Apparently, it generalizes the result that every positive harmonic function on  $\mathbb{R}^n$  is constant.

(2)  $\Omega = \{x : |x| \geq 1\}$  in  $\mathbb{R}^n$ ,  $n \geq 5$ , is an example of a domain in which there exists a biharmonic potential but no quasiharmonic potential. For, if  $s(x) = h(x) = |x|^{2-n}$ , then  $\int_{\Omega} sh \, dx < \infty$  and hence by Theorem 4.3, there exists a biharmonic potential on  $\Omega$ . But there is no quasiharmonic potential on  $\Omega$ . For, suppose  $Q(x)$  is a potential  $> 0$  on  $\Omega$  such that  $\Delta Q = 1$ ; then by the above Corollary 2,  $Q(x)$  should be harmonic, a contradiction.

#### REFERENCES

- [1] Anandam V., *Biharmonic Green functions in a Riemannian manifold*, Arab J. Math. Sc. **4** (1998), 39–45.
- [2] Anandam V., Damlakhi M., *Biharmonic Green domains in  $\mathbb{R}^n$* , Hokkaido Math. J. **27** (1998), 669–680.
- [3] Anandam V., *Biharmonic classification of harmonic spaces*, Rev. Roumaine Math. Pures Appl. **45** (2000), 383–395.
- [4] BreLOT M., *Fonctions sousharmoniques associées à une mesure*, Stud. Cerc. Şti. Mat. Iaşi **2** (1951), 114–118.
- [5] BreLOT M., *Axiomatique des fonctions harmoniques*, Les presses de l'Université de Montréal, 1966.
- [6] Loeb P.A., *An axiomatic treatment of pairs of elliptic differential equations*, Ann. Inst. Fourier **16** (1966), 167–208.
- [7] Othman S.I., Anandam V., *Liouville-Picard theorem in harmonic spaces*, Hiroshima Math. J. **28** (1998), 501–506.
- [8] Sario L., Nakai M., Wang C., Chung L.O., *Classification theory of Riemannian manifolds*, Lecture Notes in Math. 605, Springer-Verlag, 1977.

DEPARTMENT OF MATHEMATICS, KING SAUD UNIVERSITY, P.O.BOX 2455, RIYADH 11451, SAUDI ARABIA

*E-mail:* sadoon@ksu.edu.sa  
vanandam@ksu.edu.sa

(Received November 16, 2001, revised October 17, 2002)