

A factorization of quasiorder hypergroups

IVAN CHAJDA, ŠÁRKA HOŠKOVÁ

Abstract. The contribution is devoted to the question of the interchange of the construction of a quasiorder hypergroup from a quasiordered set and the factorization.

Keywords: quasiorder hypergroup, congruence on a hypergroup, relational system

Classification: 20N20, 18A40

A concept of hypergroups was formalized by [1], [2], [9], [12], [14] as follows. Let H be a non-void set and “ \circ ” a mapping of $H \times H$ into $\mathcal{P}^*(H)$ (the set of all non-void subsets of H). The pair (H, \circ) is called a *hypergroupoid*. For $A, B \in H$ we denote $A \circ B = \bigcup \{a \circ b; a \in A, b \in B\}$.

A hypergroupoid (H, \circ) is called a *hypergroup* if “ \circ ” is associative, i.e. $(a \circ b) \circ c = a \circ (b \circ c)$ for all $a, b, c \in H$, and the so-called *reproduction axiom*, i.e. $a \circ M = M = M \circ a$ for any $a \in H$, is satisfied.

Let R be a binary relation on a non-void set A . The pair $\mathcal{A} = (A, R)$ is called a *relational system*. A relational system A is called *transitive* if R is transitive and \mathcal{A} is called a *quasiordered set* whenever R is a quasiorder on A , i.e. R is a reflexive and transitive relation.

The following fact is well-known. Let $\mathcal{A} = (A, R)$ be a relational system. Denote $U_R(a) = \{x \in A; \langle a, x \rangle \in R\}$ and, for $M \subseteq A$, $U_R(M) = \{x \in A; \langle a, x \rangle \in R \text{ for all } a \in M\}$. Let $\mathcal{A} = (A, \leq)$ be a quasiordered set. Define for $a, b \in A$

$$(1) \qquad a \circ b = U_{\leq}(a) \cup U_{\leq}(b).$$

Then (A, \circ) is a hypergroup which is called a *quasiorder hypergroup* (see e.g. [9]).

The concept of congruence on a hypergroup (H, \circ) was defined by several authors. It was shown in [9, p. 151] that the definitions are equivalent. Let θ be an equivalence on a set A and $M \subseteq A$. Denote

$$\theta(M) = \{x \in A; \langle a, x \rangle \in \theta \text{ for some } a \in M\}.$$

Definition 1 ([9]). Let (H, \circ) be a hypergroup and θ be an equivalence on H . We call θ a *congruence* on (H, \circ) if for each $a, b, c, d \in H$ we have:

$$\langle a, b \rangle \in \theta \text{ and } \langle c, d \rangle \in \theta \text{ imply } \theta(a \circ c) = \theta(b \circ d).$$

The motivation of our paper is the following: Let (H, \leq) be a quasiordered set and $\mathcal{H} = (H, \circ)$ be a hypergroup, where “ \circ ” is defined by (1) (i.e. it is the *induced quasiorder hypergroup*). From now on, a quasiorder will be denoted by the symbol “ \leq ”. Let θ be a congruence on (H, \circ) .

I. Does there exist an equivalence ψ on (H, \leq) such that $(H/\psi, \leq/\psi)$ is a quasiordered set and $\mathcal{H}/\theta = (H, \circ)/\theta$ is isomorphic to the quasiorder hypergroup induced by $(H/\psi, \leq/\psi)$?

It can be visualized by the following diagram:

$$(D1) \quad \begin{array}{ccc} (H, \leq) & \xrightarrow{\psi?} & (H/\psi, \leq/\psi) \\ \downarrow & & \downarrow \\ \mathcal{H} = (H, \circ) & \longrightarrow & \mathcal{H}/\theta = (H, \circ)/\theta \end{array}$$

II. Suppose that $\mathcal{H} = (H, \circ)$ be a quasiorder hypergroup induced by a quasiordered set (H, \leq) and let ψ be an equivalence on (H, \circ) such that $(H/\psi, \leq/\psi)$ is a quasiordered set again. Under what conditions on ψ does there exist a congruence θ on \mathcal{H} such that \mathcal{H}/θ is isomorphic to the quasiorder hypergroup induced by $(H/\psi, \leq/\psi)$?

It can be visualized by the following diagram:

$$(D2) \quad \begin{array}{ccc} (H, \leq) & \xrightarrow{\psi} & (H/\psi, \leq/\psi) \\ \downarrow & & \downarrow \\ \mathcal{H} = (H, \circ) & \xrightarrow{\theta?} & \mathcal{H}/\theta = (H, \circ)/\theta \end{array}$$

As θ and ψ are equivalences on the same set H , we can easily simplify our problems by considering $\theta = \psi$, i.e. we can ask what conditions must be satisfied by an equivalence on a quasiordered set to be a congruence on the induced quasiorder hypergroup and vice versa. First of all, we need several concepts and properties of relational systems.

Definition 2. Let $\mathcal{A} = (A, R)$ be a relational system and θ be an equivalence on A . For $a \in A$ denote by $[a]_\theta$ the θ -class containing the element a . Define R/θ on A/θ as follows:

$$\langle [a]_\theta, [b]_\theta \rangle \in R/\theta \text{ if and only if there exist } x \in [a]_\theta, y \in [b]_\theta \text{ with } \langle x, y \rangle \in R.$$

The system $\mathcal{A}/\theta = (A/\theta, R/\theta)$ is called a *quotient system of \mathcal{A} by θ* .

The following statement is almost trivial:

Lemma 1. Let $\mathcal{A} = (A, R)$ be a relational system and θ be an equivalence on A . If R is reflexive or symmetric, then also R/θ has the same property.

Unfortunately, a similar statement fails for transitive relational systems, see the following:

Example 1. Let $A = \{a, b, c, d\}$ and (A, \leq) be a quasiordered set visualized in Figure 1 below.




Figure 1

Let θ be an equivalence on A defined by the partition $\{a\}, \{b, c\}, \{d\}$. Then we have on A/θ the following:

$$[a]_\theta \leq/\theta [b]_\theta \text{ since } a \leq c \text{ and } c \in [b]_\theta,$$

$$[b]_\theta \leq/\theta [d]_\theta \text{ since } b \leq d,$$

but $[a]_\theta \leq/\theta [d]_\theta$ does not hold, i.e. “ \leq/θ ” is not transitive.

This example motivates us to introduce the following concept:

Definition 3. Let $\mathcal{A} = (A, R)$ be a relational system and θ be an equivalence on A . We say that θ is *compatible* (with \mathcal{A}) if either $\theta = A \times A$ or it satisfies the following condition:

- (c) for each $x, y, z \in A$ with $\langle x, y \rangle \in \theta$ and $\langle y, z \rangle \in R$ there exists $q \in A$ such that $\langle x, q \rangle \in R$ and $\langle q, z \rangle \in \theta$.

Theorem 1. Let $\mathcal{A} = (A, \leq)$ be a quasiordered set and θ be a compatible equivalence on A . Then $\mathcal{A}/\theta = (A/\theta, \leq/\theta)$ is a quasiordered set.

PROOF: By Lemma 1 " \leq/θ " is reflexive on A/θ . Suppose $[a]_\theta \leq/\theta [b]_\theta$ and $[b]_\theta \leq/\theta [c]_\theta$. Then there are $x \in [a]_\theta, y, y' \in [b]_\theta$ and $z \in [c]_\theta$ such that $x \leq y, y' \leq z$. By (c) of Definition 3 there exists $q \in [c]_\theta$ such that $y \leq q$. Due to transitivity of " \leq ", $x \leq q$, thus $[a]_\theta \leq/\theta [c]_\theta$, thus " \leq/θ " is also transitive. \square

Definition 4. Let $\mathcal{A} = (A, R), \mathcal{B} = (B, Q)$ be relational systems. Then a mapping $f: A \rightarrow B$ is called:

- (a) *monotonous* if $\langle a, b \rangle \in R$ implies $\langle f(a), f(b) \rangle \in Q$;
- (b) *strong homomorphism* if it is monotonous and for each $a, b \in A$ with $\langle f(a), f(b) \rangle \in Q$ there exist $c, d \in A$ such that $\langle c, d \rangle \in R$ and $f(c) = f(a), f(d) = f(b)$;
- (c) *U-morphism* if it is surjective and for each $x \in A$ we have

$$f(U_R(x)) = U_Q(f(x)).$$

Lemma 2. Let $\mathcal{A} = (A, R), \mathcal{B} = (B, Q)$ be relational systems and a mapping $f: A \rightarrow B$ be a U-morphism. Then f is a strong homomorphism.

PROOF: Suppose that $a, b \in A, \langle a, b \rangle \in R$. Then $b \in U_R(a)$ and since f is a U-morphism, $f(b) \in f(U_R(a)) = U_Q(f(a))$, which gives $\langle f(a), f(b) \rangle \in Q$. Thus f is monotonous.

Suppose now that $\langle f(a), f(b) \rangle \in Q$. Then $f(b) \in U_Q(f(a)) = f(U_R(a))$, thus there exists $c \in U_R(a)$ with $f(c) = f(b)$. But $c \in U_R(a)$ implies $\langle a, c \rangle \in R$. Hence, f is a strong homomorphism. \square

Theorem 2. Let $\mathcal{A} = (A, R), \mathcal{B} = (B, Q)$ be relational systems and $f: A \rightarrow B$ a surjective mapping. The following are equivalent:

- (a) f is a U-morphism;
- (b) f is monotonous and for each $x, y \in A$ with $\langle f(x), f(y) \rangle \in Q$ there exists $z \in A$ such that $\langle x, z \rangle \in R$ and $f(y) = f(z)$.

PROOF: (a) \Rightarrow (b). By Lemma 2 we have that f is monotonous. Suppose that $\langle f(x), f(y) \rangle \in Q$. Then $f(y) \in U_Q(f(x)) = f(U_R(x))$, thus there is $z \in U_R(x)$ (i.e. $\langle x, z \rangle \in R$) such that $f(z) = f(y)$.

(b) \Rightarrow (a). Let $f: A \rightarrow B$ be a surjective and monotonous mapping. Then clearly

$$f(U_R(x)) \subseteq U_Q(f(x)).$$

Let $z \in U_Q(f(x))$. Then $z = f(w)$ for some $w \in A$, where $\langle f(x), f(w) \rangle \in Q$. By (b) there exists $c \in A$ such that $\langle x, c \rangle \in R$ and $f(c) = f(w) = z$. Thus $c \in U_R(x)$ proving the converse inclusion, i.e. f is a U-morphism. \square

Theorem 3. *Let $\mathcal{A} = (A, R)$ be a relational system and θ be a compatible equivalence on \mathcal{A} . Then the canonical mapping $h_\theta: A \mapsto A/\theta$ given by $h_\theta(a) = [a]_\theta$ is a U -morphism.*

PROOF: Let θ be a compatible equivalence on $\mathcal{A} = (A, R)$. Suppose $\langle a, b \rangle \in R$, $a, b \in A$. Thus $[a]_\theta R/\theta [b]_\theta$ and hence h_θ is monotonous. Of course, h_θ is surjective. We only need to verify (b) of Theorem 2. Suppose $[x]_\theta R/\theta [y]_\theta$. Then there exist $a \in [x]_\theta$, $b \in [y]_\theta$ with $\langle a, b \rangle \in R$. By Definition 3, there exists $q \in A$ such that $\langle x, q \rangle \in R$ and $\langle q, b \rangle \in \theta$, i.e. $[q]_\theta = [b]_\theta$. Hence, (b) of Theorem 2 is satisfied. \square

Theorem 4. *Let $\mathcal{A} = (A, R)$, $\mathcal{B} = (B, Q)$ be relational systems and $f: A \rightarrow B$ a U -morphism. Then the induced equivalence*

$$\langle a, b \rangle \in \theta_f \text{ iff } f(a) = f(b)$$

is compatible with \mathcal{A} .

PROOF: Suppose $\langle x, y \rangle \in \theta_f$ and $\langle y, z \rangle \in R$. Then $f(x) = f(y)$ and by Lemma 2 we have $\langle f(y), f(z) \rangle \in Q$. Further, by Theorem 2 there exists $u \in A$ such that $\langle x, u \rangle \in R$ and $f(u) = f(z)$, i.e. $\langle u, z \rangle \in \theta_f$. Hence, condition (c) of Definition 3 is satisfied for $q = u$, i.e. θ_f is compatible with \mathcal{A} . \square

We can finish our treatment concerning the problems in the introduction:

Corollary 1. *Let (H, \leq) be a quasiordered set and $\mathcal{H} = (H, \circ)$ the induced quasiorder hypergroup. Let θ be a congruence on \mathcal{H} . Then θ is a compatible equivalence on (H, \leq) and \mathcal{H}/θ is isomorphic to the quasiorder hypergroup induced by the quasiordered set $(H/\theta, \leq/\theta)$.*

PROOF: By Lemma 1 the relation “ \leq/θ ” is reflexive; later on we will verify that it is also transitive.

First we will prove that the canonical mapping $h_\theta: (H, \leq) \mapsto (H/\theta, \leq/\theta)$ is a U -morphism.

For $x \leq y$ we have $[x]_\theta \leq/\theta [y]_\theta$, thus $[U_{\leq}(x)]_\theta \subseteq U_{\leq/\theta}([x]_\theta)$.

Let $[z]_\theta \in U_{\leq/\theta}([x]_\theta)$. Then $[x]_\theta \leq/\theta [z]_\theta$, which implies that there exist $x_1, z_1 \in H$ such that $\langle x, x_1 \rangle \in \theta$, $\langle z, z_1 \rangle \in \theta$, $x_1 \leq z_1$. Therefore, as θ is a congruence on \mathcal{H} ,

$$\langle a, b \rangle \in \theta \Rightarrow [a \circ a]_\theta = [b \circ b]_\theta \Leftrightarrow [U_{\leq}(a)]_\theta = [U_{\leq}(b)]_\theta.$$

Thus $[U_{\leq}(x)]_\theta = [U_{\leq}(x_1)]_\theta$, $[U_{\leq}(z)]_\theta = [U_{\leq}(z_1)]_\theta$. Further

$$U_{\leq}(z_1) \subseteq U_{\leq}(x_1) \Rightarrow [U_{\leq}(z_1)]_\theta \subseteq [U_{\leq}(x_1)]_\theta \Rightarrow [U_{\leq}(z)]_\theta \subseteq [U_{\leq}(x)]_\theta.$$

As $[z]_\theta \in [U_{\leq}(z)]_\theta$, we get $[z]_\theta \in [U_{\leq}(x)]_\theta$ and $U_{\leq/\theta}([x]_\theta) \subseteq [U_{\leq}(x)]_\theta$.

Together we have obtained that $[U_{\leq}(x)]_\theta = U_{\leq/\theta}([x]_\theta)$ and the canonical mapping $h_\theta: a \rightarrow [a]_\theta$ is a U -morphism.

In Theorem 4 let us put $f = h_\theta$, $A = H$, $B = H/\theta$, $R = \leq$ and $Q = \leq/\theta$. Then the induced equivalence θ_{h_θ} is compatible with (H, \leq) . But $\theta_{h_\theta} = \theta$. Now Theorem 1 implies that “ \leq/θ ” is a quasiorder on H/θ .

Due to the fact that h_θ is U -morphism we get

$$(2) \quad \begin{aligned} [a \circ b]_\theta &= [U_{\leq}(a) \cup U_{\leq}(b)]_\theta = [U_{\leq}(a)]_\theta \cup [U_{\leq}(b)]_\theta \\ &= U_{\leq/\theta}([a]_\theta) \cup U_{\leq/\theta}([b]_\theta). \end{aligned}$$

The operations “ \circ_θ ” (for the definition of a hyperoperation induced by the congruence θ on the quotient hypergroup H/θ see [9, p. 153]) and “ \star ” (compare (1)), where

$$\begin{aligned} [a]_\theta \circ_\theta [b]_\theta &= [a \circ b]_\theta, \\ [a]_\theta \star [b]_\theta &= U_{\leq/\theta}([a]_\theta) \cup U_{\leq/\theta}([b]_\theta), \end{aligned}$$

are the same due to (2). Thus Diagram (D1) (with $\theta = \psi$) commutes. □

Corollary 2. *Let ψ be a compatible equivalence on a quasiordered set (H, \leq) . Then ψ is a congruence on the quasiorder hypergroup \mathcal{H} induced by (H, \leq) and \mathcal{H}/ψ is isomorphic to the quasiorder hypergroup induced by $(H/\psi, \leq/\psi)$.*

PROOF: As ψ is compatible, by Theorem 1 we get that “ \leq/ψ ” is the quasiorder and by Theorem 3 we have $[U_{\leq}(x)]_\psi = U_{\leq/\psi}([x]_\psi)$. If $\langle a, c \rangle \in \psi$, $\langle b, d \rangle \in \psi$, then $[a]_\psi = [c]_\psi$, $[b]_\psi = [d]_\psi$, which implies

$$[U_{\leq}(a)]_\psi = U_{\leq/\psi}([a]_\psi) = U_{\leq/\psi}([c]_\psi) = [U_{\leq}(c)]_\psi.$$

Analogously $[U_{\leq}(b)]_\psi = [U_{\leq}(d)]_\psi$.

Then

$$\begin{aligned} [a \circ b]_\psi &= [U_{\leq}(a) \cup U_{\leq}(b)]_\psi = [U_{\leq}(a)]_\psi \cup [U_{\leq}(b)]_\psi \\ &= [U_{\leq}(c)]_\psi \cup [U_{\leq}(d)]_\psi = [U_{\leq}(c) \cup U_{\leq}(d)]_\psi = [c \circ d]_\psi, \end{aligned}$$

which means that ψ is the congruence on (H, \circ) . The commutativity of Diagram (D2) can be verified in the same way as in Corollary 1. □

Example 2. Consider the quasiordered set (H, \leq) , where $H = \{a, b, c, d\}$ is depicted in Figure 2, and the equivalence θ determined by the partition $\{a, b\}, \{c, d\}$. It is easy to verify that θ does not satisfy condition (c) of Definition 3.




Figure 2

Although $(H/\theta, \leq/\theta)$ is still a quasiordered set (see Figure 3), θ is not a congruence on \mathcal{H} induced by (H, \leq) (as $[d \circ d]_\theta \neq [d \circ c]_\theta$, see Table 1):

\circ	a	b	c	d
a	$\{a, b, d\}$	$\{a, b, d\}$	H	$\{a, b, d\}$
b	$\{a, b, d\}$	$\{b\}$	$\{b, c, d\}$	$\{b, d\}$
c	H	$\{b, c, d\}$	$\{b, c, d\}$	$\{b, c, d\}$
d	$\{a, b, d\}$	$\{b, d\}$	$\{b, c, d\}$	$\{d\}$

Table 1




Figure 3

Example 3. Let (H, \leq) be the quasiordered set in Figure 4(a):




Figure 4

Then (H, \leq) induces the quasiorder hypergroup $\mathcal{H} = (H, \leq)$ given by Table 2:

\circ	a	b	c
a	H	H	H
b	H	H	H
c	H	H	$\{c\}$

Table 2

The equivalence θ given by the partition $\{a, b\}, \{c\}$ is clearly a congruence on \mathcal{H} and a compatible equivalence on (H, \leq) . The quotient quasiordered set $(H/\theta, \leq/\theta)$ is visualized in Figure 4(b) and \mathcal{H}/θ is determined by Table 3:

\circ_θ	$\{a, b\}$	$\{c\}$
$\{a, b\}$	H/θ	H/θ
$\{c\}$	H/θ	$\{c\}$

Table 3

REFERENCES

- [1] Corsini P., *Prolegomena of Hypergroup Theory*, Aviani Editore, Tricesimo, 1993.
- [2] Corsini P., *Hypergraphs and hypergroups*, Algebra Universalis **35** (1996), 548–555.
- [3] Corsini P., *On the hypergroup associated with a binary relation*, Mult.-Valued Log. **5** (2000), 407–419.
- [4] Corsini P., Leoreanu V., *Applications of Hyperstructure Theory*, Kluwer Academic Publishers, Dordrecht, Hardbound, ISBN 1-4020-1222-5, 2003.
- [5] Corsini P., Leoreanu V., *Hypergroups and binary relations*, Algebra Universalis **43** (2000), 321–330.
- [6] Corsini P., Leoreanu V., *Hypergroups associated with graphs*, Ital. J. Pure Appl. Math. **4** (1998), 119–126.
- [7] Chajda I., *Algebraic Theory of Tolerance Relations*, The Palacky University Olomouc, 1991.
- [8] Chajda I., *Congruences in transitive systems*, Math. Notes (Miskolc), to appear.
- [9] Chvalina J., *Functional Graphs, Quasi-ordered Sets and Commutative Hypergroups*, Masarykova Universita, Brno, 1995 (in Czech).
- [10] Chvalina J., Hošková Š., *Abelization of quasi-hypergroups as reflexion*, Second Conf. Math. and Physics at Technical Universities, Military Academy Brno, Proc. of Contributions, MA Brno (2001), pp. 47–53 (in Czech).
- [11] Chvalina J., Hošková Š., *Abelization of weakly associative hyperstructures based on their direct squares*, Acta Math. Inform. Univ. Ostraviensis, 2003, 12p.
- [12] Marty F., *Sur une généralisation de la notion de groupe*, Huitième Congr. math. Scan. (1934), Stockholm, pp. 45–49.

- [13] Rosenberg I., *Hypergroups and join spaces determined by relations*, Ital. J. Pure Appl. Math. **4** (1998), 93–101.
- [14] Vougiouklis T., *Hyperstructures and their Representations*, Hadronic Press Monographs in Mathematics, Palm Harbor, Florida, 1994.

DEPARTMENT OF ALGEBRA AND GEOMETRY, FACULTY OF SCIENCE UP, TOMKOVA 40,
779 00 OLOMOUC, CZECH REPUBLIC

E-mail: chajda@aix.upol.cz

DEPARTMENT OF MATHEMATICS, MILITARY ACADEMY BRNO, KOUNICOVA 65, 612 00 BRNO,
CZECH REPUBLIC

E-mail: hoskova@scova.vabo.cz

(Received December 17, 2003, revised May 7, 2004)