

Affine regular decagons in GS–quasigroup

V. VOLENEC, Z. KOLAR–BEGOVIĆ

Abstract. In this article the “geometric” concept of the affine regular decagon in a general GS–quasigroup is introduced. The relationships between affine regular decagon and some other geometric concepts in a general GS–quasigroup are explored. The geometrical presentation of all proved statements is given in the GS–quasigroup $\mathbb{C}(\frac{1}{2}(1 + \sqrt{5}))$.

Keywords: GS–quasigroup, affine regular decagon, affine regular pentagon

Classification: 20N05

1. Introduction

In [1] the concept of GS–quasigroup is defined. A quasigroup (Q, \cdot) is said to be golden section quasigroup or shortly GS–quasigroup if it satisfies the (mutually equivalent) identities

$$(1) \quad a(ab \cdot c) \cdot c = b,$$

$$(1') \quad a \cdot (a \cdot bc)c = b$$

and the identity of idempotency

$$(2) \quad aa = a.$$

The considered GS–quasigroup (Q, \cdot) satisfies the identities of mediality, elasticity, left and right distributivity i.e. we have the identities

$$(3) \quad ab \cdot cd = ac \cdot bd,$$

$$(4) \quad a \cdot ba = ab \cdot a,$$

$$(5) \quad a \cdot bc = ab \cdot ac,$$

$$(5') \quad ab \cdot c = ac \cdot bc.$$

Further, the identities

$$(6) \quad a(ab \cdot b) = b,$$

$$(6') \quad (b \cdot ba)a = b,$$

$$(7) \quad a(ab \cdot c) = b \cdot bc,$$

$$(7') \quad (c \cdot ba)a = cb \cdot b,$$

$$(8) \quad a(a \cdot bc) = b(b \cdot ac),$$

$$(8') \quad (cb \cdot a)a = (ca \cdot b)b$$

and equivalencies

$$(9) \quad ab = c \Leftrightarrow a = c \cdot cb,$$

$$(9') \quad ab = c \Leftrightarrow b = ac \cdot c$$

also hold.

Let \mathbb{C} be the set of points of the Euclidean plane. For any two different points a, b we define $ab = c$ if the point b or a divides the pair a, c or the pair b, c respectively, in ratio of the golden section.

In [1] it is proved that (\mathbb{C}, \cdot) is a GS–quasigroup in both cases. We shall denote these two quasigroups by $\mathbb{C}(\frac{1}{2}(1 + \sqrt{5}))$ and $\mathbb{C}(\frac{1}{2}(1 - \sqrt{5}))$ because we have $c = \frac{1}{2}(1 + \sqrt{5})$ or $c = \frac{1}{2}(1 - \sqrt{5})$ if $a = 0$ and $b = 1$. These quasigroups can give a motivation for the definition of “geometric” notions and proving of “geometric” properties of a general GS–quasigroup. In the quasigroup $\mathbb{C}(\frac{1}{2}(1 + \sqrt{5}))$ we shall illustrate (by figures) the properties of a general GS–quasigroup. If we interchange the roles of both factors in all products we will get the presentation in the same figure for the quasigroup $\mathbb{C}(\frac{1}{2}(1 - \sqrt{5}))$.

These two mentioned quasigroups are mutually equivalent since the following statement is obviously valid.

Lemma 1.1. *If the operation \bullet on the set Q is defined by the equivalency $a \bullet b = c \Leftrightarrow ba = c$, i.e. by the identity $a \bullet b = ba$, then (Q, \bullet) is a GS–quasigroup if and only if (Q, \cdot) is a GS–quasigroup.*

From now on let (Q, \cdot) be any GS–quasigroup. The elements of the set Q are said to be *points*.

The following statements are proved in [1] and they will be used later.

Lemma 1.2. *Any three of four equalities $ab = d$, $ae = f$, $dc = e$, $fc = b$ imply the remaining equality.*

Lemma 1.3. *Any two of four equalities $ab = c$, $dc = b$, $ac = d$, $db = a$ imply the remaining two equalities.*

The points a, b, c, d are said to be the vertices of a *parallelogram* and we write $\text{Par}(a, b, c, d)$ iff the identity $a \cdot b(ca \cdot a) = d$ holds. In [1] numerous properties of the quaternary relation Par on the set Q are proved. Let us mention just the following statements which we shall use afterwards.

Lemma 1.4. *From $\text{Par}(a, b, c, d)$ and $\text{Par}(c, d, e, f)$ it follows $\text{Par}(a, b, f, e)$.*

Lemma 1.5. *Let a, b, c be any three points and $d = ac$, $e = ab$, $f = ec$, $g = df$. Then the statements $\text{Par}(a, b, d, f)$, $\text{Par}(b, e, f, g)$, $\text{Par}(a, e, d, g)$ are valid.*

We shall say that b is the *midpoint* of the pair of points a, c and write $M(a, b, c)$ if $\text{Par}(a, b, c, b)$. In [1] the following statements, by means of the properties of the quaternary relation Par , are proved.

Lemma 1.6. *For any points a, b there is only one point c such that $M(a, b, c)$. The statement $M(a, b, c)$ implies the statement $M(c, b, a)$. For any point a it is valid $M(a, a, a)$.*

Lemma 1.7. *The statement $M(a, b, c)$ holds if and only if $c = ba \cdot b$.*

Lemma 1.8. *For any point p the statements $M(a, b, c)$, $M(pa, pb, pc)$, $M(ap, bp, cp)$ are mutually equivalent.*

In [2] the concept of the GS–trapezoid is defined. The points a, b, c, d successively are said to be the vertices of the *golden section trapezoid* and it is denoted by $\text{GST}(a, b, c, d)$ if the identity $a \cdot ab = d \cdot dc$ holds. Because of (9) this identity is equivalent to the identity $d = (a \cdot ab)c$.

In [4] the concept of affine regular pentagon is defined. The points a, b, c, d, e successively are said to be the vertices of the *affine regular pentagon* and it is denoted by $\text{ARP}(a, b, c, d, e)$ if any two (and then all five) of the five statements $\text{GST}(a, b, c, d)$, $\text{GST}(b, c, d, e)$, $\text{GST}(c, d, e, a)$, $\text{GST}(d, e, a, b)$, $\text{GST}(e, a, b, c)$ are valid.

The concept of the DGS–trapezoid is introduced in [3]. Points a, b, c, d are said to be the vertices of the *double golden section trapezoid* or shorter DGS–trapezoid and we write $\text{DGST}(a, b, c, d)$ if the equality $ab = dc$ holds.

The points o, a, b, c are said to be the vertices of a *golden section deltoid* and we write $\text{GSD}(o, a, b, c)$ if and only if the identity $c = oa \cdot b$ is valid ([5]). In [5] the following statements are proved.

Lemma 1.9. *For any point p the statements $\text{GSD}(o, a, b, c)$, $\text{GSD}(po, pa, pb, pc)$ and $\text{GSD}(op, ap, bp, cp)$ are mutually equivalent.*

Lemma 1.10. *If the statements $\text{GSD}(o, a, b, c)$, $\text{GSD}(o, b, c, d)$ hold then $ab = dc = e$ i.e. $\text{DGST}(a, b, c, d)$ and $\text{Par}(o, a, e, d)$ hold.*

2. Affine regular decagon in GS–quasigroup

Now we are going to introduce the concept of the affine regular decagon in a general GS–quasigroup. Firstly, we will prove the theorem which will lead to the definition of the mentioned concept.

Theorem 2.1. *From the equations*

$$(10) \quad oa_i \cdot a_{i+1} = a_{i+2}$$

for $i = 0, 1, 2, 3, 4, 5, 6, 7$ the equations (10) for $i = 8, 9$ follow, where indexes are taken modulo 10 (Figure 1).

PROOF: If we denote by k the equality (10) for $i = k$ then we get

$$\begin{aligned}
 oa_1 \cdot a_0 &\stackrel{(1)}{=} o[o(oa_1 \cdot a_0) \cdot o] \cdot o \stackrel{(5)}{=} o[(o \cdot oa_1)(oa_0) \cdot o] \cdot o \\
 &\stackrel{(6')}{=} o[(o \cdot oa_1)(oa_0) \cdot (o \cdot oa_1)a_1] \cdot o \stackrel{(5)}{=} [o \cdot (o \cdot oa_1)(oa_0 \cdot a_1)]o \\
 &\stackrel{0}{=} [o \cdot (o \cdot oa_1)a_2]o \stackrel{(5')}{=} [o \cdot (oa_2)(oa_1 \cdot a_2)]o \stackrel{1}{=} o(oa_2 \cdot a_3) \cdot o \stackrel{2}{=} oa_4 \cdot o \\
 &\stackrel{(6')}{=} oa_4 \cdot (o \cdot oa_4)a_4 \stackrel{(5')}{=} o(o \cdot oa_4) \cdot a_4 \stackrel{(1)}{=} o(o \cdot oa_4) \cdot [o(oa_4 \cdot a_5) \cdot a_5] \\
 &\stackrel{(3)}{=} [o \cdot o(oa_4 \cdot a_5)] \cdot (o \cdot oa_4)a_5 \stackrel{(5')}{=} [o \cdot o(oa_4 \cdot a_5)][oa_5 \cdot (oa_4 \cdot a_5)] \\
 &\stackrel{4}{=} (o \cdot oa_6)(oa_5 \cdot a_6) \stackrel{5}{=} (o \cdot oa_6)a_7 \stackrel{(5')}{=} oa_7 \cdot (oa_6 \cdot a_7) \stackrel{6}{=} oa_7 \cdot a_8 \stackrel{7}{=} a_9,
 \end{aligned}$$

wherefrom it follows

$$oa_9 \cdot a_0 = o(oa_1 \cdot a_0) \cdot a_0 \stackrel{(1)}{=} a_1,$$

which means that from equality (10) where $i = 0, 1, 2, 4, 5, 6, 7$ the equality (10) for $i = 9$ follows, and similarly from equalities (10) for $i = 9, 0, 1, 3, 4, 5, 6$ the equality (10) for $i = 8$ follows. □




FIGURE 1

We shall say that $(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9)$ is the *affine regular decagon with the center o* and we shall write $Aff_o(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9)$ if in a cycle of the equalities (10) for $i = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9$, any eight adjacent (and then all ten) equalities are valid.

From Theorem 2.1 it follows immediately

Corollary 2.2. *For any points o, a_0, a_1 there is a unique octuple of the points $a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9$ so that $Aff_o(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9)$.*

Theorem 2.3. *If $(i_0, i_1, i_2, i_3, i_4, i_5, i_6, i_7, i_8, i_9)$ is any cyclic permutation of $(0, 1, 2, 3, 4, 5, 6, 7, 8, 9)$ or of $(9, 8, 7, 6, 5, 4, 3, 2, 1, 0)$ then $Aff_o(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9)$ implies $Aff_o(a_{i_0}, a_{i_1}, a_{i_2}, a_{i_3}, a_{i_4}, a_{i_5}, a_{i_6}, a_{i_7}, a_{i_8}, a_{i_9})$.*

PROOF: It is enough to prove the identity

$$(11) \quad oa_i \cdot a_{i-1} = a_{i-2}.$$

However, we get

$$oa_i \cdot a_{i-1} \stackrel{(10)}{=} o(oa_{i-2} \cdot a_{i-1}) \cdot a_{i-1} \stackrel{(1)}{=} a_{i-2}.$$

□

Further, let $Aff_o(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9)$ where $i \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$.

If we denote

$$(12) \quad oa_i = b_i,$$

then from (10) and (11) immediately follows

$$(13) \quad b_i a_{i\pm 1} = a_{i\pm 2}.$$

Theorem 2.4. *The statements $Par(o, b_i, b_{i\pm 1}, a_{i\pm 3})$, $Par(a_i, b_i, a_{i\pm 2}, a_{i\pm 3})$, $Par(o, a_i, b_{i\pm 1}, a_{i\pm 2})$ are valid.*

PROOF: On the bases of (12) and (13) we get equalities $oa_i = b_i$, $oa_{i\pm 1} = b_{i\pm 1}$, $b_i a_{i\pm 1} = a_{i\pm 2}$, $b_{i\pm 1} a_{i\pm 2} = a_{i\pm 3}$ wherefrom according to Lemma 1.5 the statements of theorem follow. □

Because of (13) we get equalities $b_i a_{i\pm 1} = a_{i\pm 2}$ and $b_{i\pm 3} a_{i\pm 2} = a_{i\pm 1}$ wherefrom owing to Lemma 1.3 it follows

$$(14) \quad b_i a_{i\pm 2} = b_{i\pm 3}.$$

Theorem 2.5. *The statements $M(a_i, o, a_{i+5})$, $M(b_i, o, b_{i+5})$ are valid.*

PROOF: Owing to Theorem 2.4 $Par(a_i, o, b_{i+3}, b_{i+2})$ and $Par(b_{i+3}, b_{i+2}, o, a_{i+5})$ are valid, wherefrom according to Lemma 1.4 it follows $Par(a_i, o, a_{i+5}, o)$, i.e. $M(a_i, o, a_{i+5})$. Hence, according to Lemma 1.8 it follows $M(oa_i, oo, oa_{i+5})$ i.e. $M(b_i, o, b_{i+5})$. □

The property $M(a_i, o, a_{i+5})$ justifies the fact that the point o is called the center of the considered affine-regular decagon.

If we apply Lemma 1.7 we get $M(a_i, o, oa_i \cdot o)$, which together with $M(a_i, o, a_{i+5})$, according to Lemma 1.6, gives the equality $a_{i+5} = oa_i \cdot o$. Whence, on the basis of (12), we get

$$(15) \quad b_i o = a_{i+5}.$$

If the statement $Aff_o(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9)$ is valid then we shall say that $(a_0, a_3, a_6, a_9, a_2, a_5, a_8, a_1, a_4, a_7)$ is the *affine regular star-shaped decagon with the center o* and we shall write $\overline{Aff_o}(a_0, a_3, a_6, a_9, a_2, a_5, a_8, a_1, a_4, a_7)$ (Figure 2).

It is obvious that the implication $\overline{Aff_o}(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9) \Rightarrow \overline{Aff_o}(a_0, a_3, a_6, a_9, a_2, a_5, a_8, a_1, a_4, a_7)$ is valid.

If $(i_0, i_1, i_2, i_3, i_4, i_5, i_6, i_7, i_8, i_9)$ is any cyclic permutation of $(0, 1, 2, 3, 4, 5, 6, 7, 8, 9)$ or of $(9, 8, 7, 6, 5, 4, 3, 2, 1, 0)$ then the implication $\overline{Aff_o}(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9) \Rightarrow \overline{Aff_o}(a_{i_0}, a_{i_1}, a_{i_2}, a_{i_3}, a_{i_4}, a_{i_5}, a_{i_6}, a_{i_7}, a_{i_8}, a_{i_9})$ is also valid.




FIGURE 2

Let us take (Figure 3)

$$(16) \quad a_i o = c_{i+5}.$$

Then we get successively

$$a_{i+5} \stackrel{(15)}{=} b_i o \stackrel{(12)}{=} oa_i \cdot o \stackrel{(4)}{=} o \cdot a_i o \stackrel{(16)}{=} oc_{i+5},$$

i.e. the equation

$$(17) \quad oc_i = a_i$$

is valid. Further, we get

$$o \cdot a_i c_{i\pm 2} \stackrel{(5)}{=} oa_i \cdot oc_{i\pm 2} \stackrel{(12), (17)}{=} b_i a_{i\pm 2} \stackrel{(14)}{=} b_{i\pm 3} \stackrel{(12)}{=} oa_{i\pm 3},$$

wherefrom it follows

$$(18) \quad a_i c_{i\pm 2} = a_{i\pm 3}.$$

Now, we obtain

$$a_{i+3} \cdot a_i o \stackrel{(16)}{=} a_{i+3} \cdot c_{i+5} \stackrel{(18)}{=} a_{i+6}.$$

If we interchange the roles of both factors in all products in the above equality, then we get the equality $oa_i \cdot a_{i+3} = a_{i+6}$, which means that in the quasigroup $\mathbb{C}(\frac{1}{2}(1 - \sqrt{5}))$ the statement $Aff_o(a_0, a_3, a_6, a_9, a_2, a_5, a_8, a_1, a_4, a_7)$ holds when in the quasigroup $\mathbb{C}(\frac{1}{2}(1 + \sqrt{5}))$ the statement $Aff_o(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9)$ i.e. $\overline{Aff_o}(a_0, a_3, a_6, a_9, a_2, a_5, a_8, a_1, a_4, a_7)$ holds.

Therefore, the affine regular decagon in the first quasigroup is the affine regular star-shaped decagon in the second quasigroup, and conversely the affine regular decagon in the second quasigroup is the affine regular star-shaped decagon in the first quasigroup.

According to Lemma 1.1 these two quasigroups are equivalent, so it means that it is a matter of convention which of these two decagons $(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9)$ and $(a_0, a_3, a_6, a_9, a_2, a_5, a_8, a_1, a_4, a_7)$ will be called affine regular decagon and which one affine regular–star shaped decagon, since we cannot differ them in a general GS–quasigroup. Besides that, it means that each statement about affine regular decagons which is proved in a general GS–quasigroup, it is also valid for affine regular star-shaped decagon and vice versa (with above mentioned interchange of both factors in all products).




FIGURE 3

In Figure 3 dashed lines present the formulas (20), (21), (30), (31), (36) and (37) for $i = 0$ and if the sign on top is considered.

On the base of (18) we get equalities $a_i c_{i\pm 2} = a_{i\pm 3}$, $a_{i\pm 3} c_{i\pm 1} = a_i$, wherefrom owing to Lemma 1.3 it follows

$$(19) \quad a_i c_{i\pm 1} = c_{i\pm 2}.$$

Because of (17) and (19) we get equalities $oc_i = a_i$, $oc_{i\pm 1} = a_{i\pm 1}$, $a_i c_{i\pm 1} = c_{i\pm 2}$, $a_{i\pm 1} c_{i\pm 2} = c_{i\pm 3}$, wherefrom owing to Lemma 1.5 we have the statements analogous to the statements of Theorem 2.4 i.e. we get the following:

Theorem 2.6. *The statements $Par(o, a_i, a_{i\pm 1}, c_{i\pm 3})$, $Par(c_i, a_i, c_{i\pm 2}, c_{i\pm 3})$, $Par(o, c_i, a_{i\pm 1}, c_{i\pm 2})$ are valid.*

Further we get

$$oc_i \cdot c_{i+1} \stackrel{(17)}{=} a_i c_{i+1} \stackrel{(19)}{=} c_{i+2},$$

so it follows:

Theorem 2.7. *The statement $Aff_o(c_0, c_1, c_2, c_3, c_4, c_5, c_6, c_7, c_8, c_9)$ is valid (Figure 3).*

Owing to (17), (12) and (18) we get equalities $oc_{i\pm 3} = a_{i\pm 3}$, $oa_i = b_i$, $a_{i\pm 3} c_{i\pm 1} = a_i$ wherefrom according to Lemma 1.2 it follows

$$(20) \quad b_i c_{i\pm 1} = c_{i\pm 3}.$$

According to (20) we get equalities $b_i c_{i\pm 1} = c_{i\pm 3}$, $b_{i\pm 4} c_{i\pm 3} = c_{i\pm 1}$, wherefrom because of Lemma 1.3 it follows

$$(21) \quad b_i c_{i\pm 3} = b_{i\pm 4}.$$

From (12) and (17) owing to Lemma 1.3 we also get

$$(22) \quad b_i a_i = c_i,$$

$$(23) \quad b_i c_i = o.$$

Further, we get because of Lemma 1.3 and $oa_i = b_i$

$$\begin{aligned} a_i c_{i+5} &\stackrel{(17),(16)}{=} oc_i \cdot a_i o \stackrel{(3)}{=} oa_i \cdot c_i o \stackrel{(12)}{=} b_i \cdot c_i o \stackrel{(5)}{=} b_i c_i \cdot b_i o \\ &\stackrel{(23),(15)}{=} oa_{i+5} \stackrel{(12)}{=} b_{i+5}, \end{aligned}$$

i.e.

$$(24) \quad a_i c_{i+5} = b_{i+5}.$$

Now, introducing the notation (Figure 3)

$$(25) \quad ob_i = d_i$$

we get

$$(26) \quad d_i b_i = a_i,$$

$$(27) \quad d_i a_i = o.$$

Because of (27), (18) and (17) we obtain the equalities

$$d_i a_i = o, \quad a_{i\pm 3} c_{i\pm 1} = a_i, \quad oc_{i\pm 1} = a_{i\pm 1},$$

wherefrom on the basis of Lemma 1.2 it follows

$$(28) \quad d_i a_{i\pm 1} = a_{i\pm 3}.$$

Using (28) we get $d_i a_{i\pm 1} = a_{i\pm 3}$, $d_{i\pm 4} a_{i\pm 3} = a_{i\pm 1}$ wherefrom according to Lemma 1.3 it follows

$$(29) \quad d_i a_{i\pm 3} = d_{i\pm 4}.$$

Theorem 2.8. *The statements $ARP(a_0, a_2, a_4, a_6, a_8)$ and $ARP(a_1, a_3, a_5, a_7, a_9)$ are valid.*

PROOF: The first statement, for example, follows on the base of equalities

$$d_3 a_2 = a_0, \quad d_3 a_4 = a_6, \quad d_5 a_4 = a_2, \quad d_5 a_6 = a_8$$

which are presented in (28). □

Using (27), (12) and (13) we obtain equalities

$$d_i a_i = o, \quad oa_{i\pm 1} = b_{i\pm 1}, \quad b_{i\pm 2} a_{i\pm 1} = a_i,$$

which, because of Lemma 1.2, imply the following

$$(30) \quad d_i b_{i\pm 1} = b_{i\pm 2}.$$

From (30) we get

$$d_i b_{i\pm 1} = b_{i\pm 2}, \quad d_{i\pm 3} b_{i\pm 2} = b_{i\pm 1},$$

whence owing to Lemma 1.3 we obtain

$$(31) \quad d_i b_{i\pm 2} = d_{i\pm 3}.$$

Now, we have

$$ob_i \cdot b_{i+1} \stackrel{(25)}{=} d_i b_{i+1} \stackrel{(30)}{=} b_{i+2},$$

so it immediately follows

Theorem 2.9. *The statement $Aff_o(b_0, b_1, b_2, b_3, b_4, b_5, b_6, b_7, b_8, b_9)$ is valid (Figure 3).*

Further we get

$$\begin{aligned} d_i o &\stackrel{(25)}{=} ob_i \cdot o \stackrel{(4)}{=} o \cdot b_i o \stackrel{(15)}{=} oa_{i+5} \stackrel{(12)}{=} b_{i+5}, \\ d_i c_i &\stackrel{(25)}{=} ob_i \cdot c_i \stackrel{(5')}{=} oc_i \cdot b_i c_i \stackrel{(17),(23)}{=} a_i o \stackrel{(16)}{=} c_{i+5}, \\ b_i a_{i+5} &\stackrel{(12),(17)}{=} oa_i \cdot oc_{i+5} \stackrel{(5)}{=} o \cdot a_i c_{i+5} \stackrel{(24)}{=} ob_{i+5} \stackrel{(25)}{=} d_{i+5}, \end{aligned}$$

i.e.

$$(32) \quad d_i o = b_{i+5},$$

$$(33) \quad d_i c_i = c_{i+5},$$

$$(34) \quad b_i a_{i+5} = d_{i+5}$$

and

$$d_i c_{i+5} \stackrel{(16)}{=} d_i \cdot a_i o \stackrel{(5)}{=} d_i a_i \cdot d_i o \stackrel{(27),(32)}{=} ob_{i+5} \stackrel{(25)}{=} d_{i+5},$$

i.e.

$$(35) \quad d_i c_{i+5} = d_{i+5}.$$

On the basis of (25), (12) and (21) we get

$$ob_i = d_i, \quad oa_{i\pm 4} = b_{i\pm 4}, \quad b_{i\pm 4} c_{i\pm 1} = b_i,$$

wherefrom according to Lemma 1.2 we get

$$(36) \quad d_i c_{i\pm 1} = a_{i\pm 4}.$$

Analogously, because of (36), (19) and (20) we get equalities

$$d_i c_{i\pm 1} = a_{i\pm 4}, \quad a_{i\pm 4} c_{i\pm 3} = c_{i\pm 2}, \quad b_{i\pm 4} c_{i\pm 3} = c_{i\pm 1}$$

whence it follows

$$(37) \quad d_i c_{i\pm 2} = b_{i\pm 4}.$$

Theorem 2.10. *The statements $M(c_i, b_i, d_i)$, $M(d_i, a_i, c_{i+5})$ are valid.*

PROOF: The statement follows on the base of Lemma 1.7, the first one from

$$d_i \stackrel{(25)}{=} ob_i \stackrel{(23)}{=} b_i c_i \cdot b_i,$$

and the second one from the equalities (27) and (16). □

Theorem 2.11. *The statement $Aff_o(d_0, d_1, d_2, d_3, d_4, d_5, d_6, d_7, d_8, d_9)$ is valid (Figure 3).*

PROOF:

$$od_i \cdot d_{i+1} \stackrel{(25)}{=} od_i \cdot ob_{i+1} \stackrel{(5)}{=} o \cdot d_i b_{i+1} \stackrel{(30)}{=} ob_{i+2} \stackrel{(25)}{=} d_{i+2}.$$

□

Now, we will introduce some new points which will also be the vertices of new affine regular decagons.

Owing to (10) the statements $GSD(o, a_{i-1}, a_i, a_{i+1})$ and $GSD(o, a_i, a_{i+1}, a_{i+2})$ are valid, wherefrom according to Lemma 1.10 the statement $DGST(a_{i-1}, a_i, a_{i+1}, a_{i+2})$ holds i.e. $a_{i-1}a_i = a_{i+2}a_{i+1}$. Let us denote now (Figure 4)

$$(38) \quad a_{i,i+1} = a_{i-1}a_i = a_{i+2}a_{i+1},$$

then we have

$$\begin{aligned} oa_{i,i+1} \cdot a_{i+1,i+2} &\stackrel{(38)}{=} (o \cdot a_{i-1}a_i) \cdot a_i a_{i+1} \stackrel{(5)}{=} (oa_{i-1} \cdot oa_i) \cdot a_i a_{i+1} \\ &\stackrel{(3)}{=} (oa_{i-1} \cdot a_i)(oa_i \cdot a_{i+1}) \stackrel{(12)}{=} b_{i-1}a_i \cdot b_i a_{i+1} \stackrel{(13)}{=} a_{i+1}a_{i+2} \\ &\stackrel{(38)}{=} a_{i+2,i+3} \end{aligned}$$

i.e.

$$(39) \quad oa_{i,i+1} \cdot a_{i+1,i+2} = a_{i+2,i+3},$$

so it immediately follows:




FIGURE 4

Theorem 2.12. *The statement $Aff_o(a_{01}, a_{12}, a_{23}, a_{34}, a_{45}, a_{56}, a_{67}, a_{78}, a_{89}, a_{90})$ holds (Figure 4).*

Because of (12) and (2) according to Lemma 1.9 the statements $GSD(o, b_{i-1}, b_i, b_{i+1})$ and $GSD(o, b_i, b_{i+1}, b_{i+2})$ hold, wherefrom according to Lemma 1.10 the statement $DGST(b_{i-1}, b_i, b_{i+1}, b_{i+2})$ follows i.e. $b_{i-1}b_i = b_{i+2}b_{i+1}$. Let us denote

$$(40) \quad b_{i,i+1} = b_{i-1}b_i = b_{i+2}b_{i+1}.$$

Because of (12) and (5) it is obviously valid

$$(41) \quad oa_{i,i+1} = b_{i,i+1}.$$

As we have

$$ob_{i,i+1} \cdot b_{i+1,i+2} \stackrel{(41)}{=} (o \cdot oa_{i,i+1}) \cdot oa_{i+1,i+2} \stackrel{(5)}{=} o(oa_{i,i+1} \cdot a_{i+1,i+2}) \\ \stackrel{(39)}{=} oa_{i+2,i+3} \stackrel{(41)}{=} b_{i+2,i+3}$$

we immediately get:

Theorem 2.13. *The statement $Aff_o(b_{01}, b_{12}, b_{23}, b_{34}, b_{45}, b_{56}, b_{67}, b_{78}, b_{89}, b_{90})$ is valid (Figure 4).*

Analogously, because of (25) and (2) and according to Lemma 1.9 $GSD(o, d_{i-1}, d_i, d_{i+1})$ and $GSD(o, d_i, d_{i+1}, d_{i+2})$ are valid whence applying Lemma 1.10 the statement $DGST(d_{i-1}, d_i, d_{i+1}, d_{i+2})$ is valid i.e. $d_{i-1}d_i = d_{i+2}d_{i+1}$. Let us take

$$(42) \quad d_{i,i+1} = d_{i-1}d_i = d_{i+2}d_{i+1}.$$

Because of (25) and (5) we obtain

$$(43) \quad ob_{i,i+1} = d_{i,i+1}.$$

If we apply (43) and Theorem 2.13 we can get

$$od_{i,i+1} \cdot d_{i+1,i+2} = d_{i+2,i+3},$$

so it follows

Theorem 2.14. *The statement $Aff_o(d_{01}, d_{12}, d_{23}, d_{34}, d_{45}, d_{56}, d_{67}, d_{78}, d_{89}, d_{90})$ (Figure 4) is valid.*

REFERENCES

- [1] Volenec V., *GS-quasigroups*, Časopis Pěst. Mat. **115** (1990), 307–318.
- [2] Kolar Z., Volenec V., *GS-trapezoids in GS-quasigroups*, Math. Commun. **7** (2002), 143–158.
- [3] Kolar–Begović Z., Volenec V., *DGS-trapezoids in GS-quasigroups*, Math. Commun. **8** (2003), 215–218.
- [4] Kolar–Begović Z., Volenec V., *Affine regular pentagons in GS-quasigroups*, Quasigroups Related Systems **12** (2004), 103–112.
- [5] Kolar–Begović Z., Volenec V., *GS-deltoids in GS-quasigroups*, Math. Commun. **10** (2005), 117–122.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ZAGREB, BIJENIČKA C. 30,
HR-10 000 ZAGREB, CROATIA

E-mail: volenec@math.hr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSIJEK, GAJEV TRG 6, HR-31 000 OSIJEK,
CROATIA

E-mail: zkolar@mathos.hr

(Received November 10, 2007, revised March 18, 2008)