

On weakly monotonically monolithic spaces

LIANG-XUE PENG

Abstract. In this note, we introduce the concept of weakly monotonically monolithic spaces, and show that every weakly monotonically monolithic space is a D -space. Thus most known conclusions on D -spaces can be obtained by this conclusion. As a corollary, we have that if a regular space X is sequential and has a point-countable wcs^* -network then X is a D -space.

Keywords: D -space, sequential space, wcs^* -network, weakly monotonically monolithic space

Classification: Primary 54F99; Secondary 54G99

1. Introduction

The notion of a D -space was first investigated by van Douwen and Pfeffer in [6]. A *neighborhood assignment* for a space X is a function ϕ from X to the topology of the space X such that $x \in \phi(x)$ for any $x \in X$. A space X is called a D -space if for any neighborhood assignment ϕ for X there exists a closed discrete subspace D of X such that $X = \bigcup\{\phi(d) : d \in D\}$ (cf. [6]). By results of [3], we know that all semi-stratifiable spaces and all metrizable spaces are D -spaces. We also know that the union of a finite family of metrizable subspaces is a D -space and every space with a point-countable base is a D -space by results of [1] and [2], respectively.

In [5], it was proved that $C_p(X)$ is hereditarily a D -space if X is compact. Some sufficiencies of D -spaces were discussed in [4], [8], [9], [16] and [19]. Let us recall that a space is called *monolithic* if $nw(\bar{A}) \leq \max\{|A|, \omega\}$ for any $A \subset X$. By results of [5] and [9], we know that many monolithic spaces have D -property. Thus V.V. Tkachuk introduced the concept of monotonically monolithic spaces in [19]. It was proved that every monotonically monolithic space is hereditarily a D -space (cf. [19]). Thus every Lindelöf Σ -space is hereditarily a D -space (cf. [9] and [19]).

In [4] and [15], it was proved that a space with a point-countable weak base is a D -space. In [16], Peng proved that a space with a point-countable cs^* -network is a D -space. In this paper, the idea of [19] and [16] is generalized and we introduce the concept of weakly monotonically monolithic spaces. It is proved that every weakly monotonically monolithic space is a D -space. Thus we have

Project supported by the National Natural Science Foundation of China (Grant No. 10971185).

that if a regular space X is sequential and has a point-countable wcs^* -network then X is a D -space. In fact, most known results on D -spaces can be obtained by the conclusions of the paper.

All the spaces in this note are at least T_1 -spaces. Let \mathbb{N} be the set of all natural numbers and $\omega = \mathbb{N} \cup \{0\}$. In notation and terminology we will follow [7] and [10].

2. Main results

Definition 1 (cf. [19]). Given a set A in a space X say that a family \mathcal{N} of subsets of X is an *external network* of A in X if for any $x \in A$ and U the neighborhood of x there exists $B \in \mathcal{N}$ such that $x \in B \subset U$. If $A = \{x\}$ for some $x \in X$ and \mathcal{N} is an external network of A in X , then we say that \mathcal{N} is an *external network of x* .

Definition 2 (cf. [19]). We say that a space X is *monotonically monolithic* if for any $A \subset X$ we can assign an external network $\mathcal{O}(A)$ to the set \overline{A} in such a way that the following conditions are satisfied:

- (1) $|\mathcal{O}(A)| \leq \max\{|A|, \omega\}$;
- (2) if $A \subset B \subset X$ then $\mathcal{O}(A) \subset \mathcal{O}(B)$;
- (3) if α is an ordinal and we have a family $\{A_\beta : \beta < \alpha\}$ of subsets of X such that $\beta < \beta' < \alpha$ implies $A_\beta \subset A_{\beta'}$ then $\mathcal{O}(\bigcup_{\beta < \alpha} A_\beta) = \bigcup_{\beta < \alpha} \mathcal{O}(A_\beta)$.

Definition 3. We say that a space X is *weakly monotonically monolithic* if for any $A \subset X$ we can assign an external network $\mathcal{W}(A)$ of A in such a way that the following conditions are satisfied:

- (1) $|\mathcal{W}(A)| \leq \max\{|A|, \omega\}$;
- (2) if $A \subset B \subset X$ then $\mathcal{W}(A) \subset \mathcal{W}(B)$;
- (3) if α is an ordinal and we have a family $\{A_\beta : \beta < \alpha\}$ of subsets of X such that $\beta < \beta' < \alpha$ implies $A_\beta \subset A_{\beta'}$ then $\mathcal{W}(\bigcup_{\beta < \alpha} A_\beta) = \bigcup_{\beta < \alpha} \mathcal{W}(A_\beta)$;
- (4) If $A \subset X$ is not closed in X then there is some $x \in \overline{A} \setminus A$ such that $\mathcal{W}(A)$ is an external network of x .

Every monotonically monolithic space is a weakly monotonically monolithic space. We also know that if a monotonically monolithic space X is separable then X is hereditarily separable and hereditarily Lindelöf.

Example 4. Let $X = \mathbb{R}$, $\mathbb{Q} = \{x_n : n \in \mathbb{N}\}$ be the set of all rational numbers of the real set \mathbb{R} and $\mathbb{I} = \mathbb{R} \setminus \mathbb{Q}$.

For each $n \in \mathbb{N}$, let $\mathcal{B}(x_n) = \{\{x_n\}\}$ be a neighborhood base of the point x_n .

For each $x \in \mathbb{I}$, let $\mathcal{B}(x) = \{\{x\} \cup A : A \subset \mathbb{Q} \text{ and } \mathbb{Q} \setminus A \text{ is finite}\}$ be a neighborhood base of the point x . We denote the topology of X by \mathcal{T} . Thus (X, \mathcal{T}) is a T_1 -space and \mathbb{I} is a closed discrete subspace of X . So (X, \mathcal{T}) is separable but it is not Lindelöf. Thus it is not monolithic.

Let $y_1 \in \mathbb{I}$. For each $B \subset X$, we let $\mathcal{W}(B) = \{\{x\} : x \in B\}$ if $B \subset \mathbb{I}$, otherwise $\mathcal{W}(B) = \{\{x\} \cup \{y_1\} : x \in B\} \cup \{\{x\} : x \in B\}$.

We can see that \mathcal{W} witnesses weak monotonic monolithicity of X . Thus X is a weakly monotonically monolithic T_1 -space but it is not a monotonically monolithic space.

Let us recall that a family \mathcal{F} of subsets of X is a *cs*-network* of X , if for any sequence $\{x_n\}_{n \in \mathbb{N}}$ which converges to a point x and for any open set U which contains x , there is some $F \in \mathcal{F}$ such that $x \in F \subset U$ and $|\{n : x_n \in F\}| = \omega$ (cf. [13] and [14]). A space is called *sequential* if for any non-closed subset A of X , there exists a sequence $\{x_n\}_{n \in \mathbb{N}}$, $x_n \in A$ for each $n \in \mathbb{N}$, such that $\{x_n\}_{n \in \mathbb{N}}$ converges to a point $x \in \overline{A} \setminus A$. We also know that every space with a point-countable weak base is sequential and has a point-countable *cs*-network*.

Lemma 5. *If X is a sequential space with a point-countable *cs*-network*, then X is a weakly monotonically monolithic space.*

PROOF: Let \mathcal{F} be a point-countable *cs*-network* of X . For any $A \subset X$, we let $\mathcal{W}(A) = \{F : F \cap A \neq \emptyset \text{ and } F \in \mathcal{F}\}$. By the sequential property we see that $\mathcal{W}(A)$ satisfies the conditions which appear in the Definition 3. \square

Corollary 6. *Let X be a space. If X has a point-countable weak base, then X is a weakly monotonically monolithic space.*

Example 7 ([11, Example 9.3]). Let $S = \{\frac{1}{n} : n \in \mathbb{N}\} \cup \{0\}$ and $Y = [0, 1] \times S$. Let $Y' = [0, 1] \times \{\frac{1}{n} : n \in \mathbb{N}\}$ have the usual Euclidean topology as a subspace of $[0, 1] \times S$. Define a typical neighborhood $(t, 0)$ in Y to be the form $\{(t, 0)\} \cup (\bigcup\{U(t, \frac{1}{k}) : k \geq n\})$, where $U(t, \frac{1}{k})$ is a neighborhood of $(t, \frac{1}{k})$ in $[0, 1] \times \{\frac{1}{k}\}$.

In [11], it is pointed out that Y is a completely regular separable space but it is not Lindelöf, and it is also pointed out that Y is a two-to-one quotient image of the topological sum M of compact metric spaces $\{[0, 1] \times \{\frac{1}{n}\} : n \in \mathbb{N}\} \cup \{t\} \times S : t \in [0, 1]\}$.

The space Y has a point-countable weak base (this is pointed out in [12, p. 26]). So we know that the space Y is not a monotonically monolithic space but Y is a weakly monotonically monolithic regular space by Corollary 6.

In [12], Lin did not give a proof that the space Y has a point-countable weak base. To assist the reader, we give a short proof.

Let $f : M \rightarrow Y$ be the two-to-one quotient map, where M is the topological sum of compact metric spaces $\{[0, 1] \times \{\frac{1}{n}\} : n \in \mathbb{N}\} \cup \{t\} \times S : t \in [0, 1]\}$. For each $y \in Y$, we let $y = (a_1, a_2)$. Suppose $\{y_n\}_{n \in \mathbb{N}}$ is a sequence of Y such that $\{y_n\}_{n \in \mathbb{N}}$ converges to the point y and we assume that $y_n \neq y$ for each $n \in \mathbb{N}$.

- (1) If $a_2 \neq 0$ then there is some $m \in \mathbb{N}$ such that $y_n \in [0, 1] \times \{a_2\}$ for each $n > m$, since Y is determined by the collection $\{[0, 1] \times \{\frac{1}{n}\} : n \in \mathbb{N}\} \cup \{t\} \times S : t \in [0, 1]\}$.
- (2) If $a_2 = 0$ then there is some $m \in \mathbb{N}$ such that $y_n \in \{a_1\} \times S$ for each $n > m$, since $[0, 1] \times \{0\}$ is a closed discrete subspace of Y and Y is determined by the collection $\{[0, 1] \times \{\frac{1}{n}\} : n \in \mathbb{N}\} \cup \{t\} \times S : t \in [0, 1]\}$.

Thus we have the following conclusion:

If $y \in Y$ then there is some point $x_y \in f^{-1}(y)$ such that whenever a sequence $\{y_n\}_{n \in \mathbb{N}}$ of Y converges to the point y and $y_n \neq y$ for each $n \in \mathbb{N}$ then there is a sequence $\{x_n\}_{n \in \mathbb{N}}$ of X such that the sequence $\{x_n\}_{n \in \mathbb{N}}$ converges to the point x_y and $x_n \in f^{-1}(y_n)$ for each $n \in \mathbb{N}$. (This is pointed out in [12, p. 26]).

Let \mathcal{B} be a point-countable base of M . For each $y \in Y$, let $\mathcal{B}'_y = \{B : x_y \in B \text{ and } B \in \mathcal{B}\}$ and let $\mathcal{B}^*_y = \{f(B) : B \in \mathcal{B}'_y\}$. We will show that $\mathcal{B}^* = \bigcup \{\mathcal{B}^*_y : y \in Y\}$ is a point-countable weak base of Y .

We only need to prove that U is open if for any $y \in U$ there is some $B^* \in \mathcal{B}^*_y$ such that $y \in B^* \subset U$.

Suppose U is not open. Then there is a sequence $\{y_n\}_{n \in \mathbb{N}}$ such that $\{y_n\}_{n \in \mathbb{N}}$ converges to a point $y \in U$ and $y_n \notin U$ for each $n \in \mathbb{N}$, since Y is sequential. Thus there is a sequence $\{x_n\}_{n \in \mathbb{N}}$ of M such that $x_n \in f^{-1}(y_n)$ and $\{x_n\}_{n \in \mathbb{N}}$ converges to x_y . There is some $B^* \in \mathcal{B}^*_y$ such that $y \in B^* \subset U$. We let $B^* = f(B)$ for some $B \in \mathcal{B}'_y$. Thus $x_y \in B$, and hence there is some $m \in \mathbb{N}$ such that $x_n \in B$ for each $n > m$. So $y_n = f(x_n) \in f(B) = B^* \subset U$. This contradicts $y_n \notin U$. Thus \mathcal{B}^* is a weak base of Y . Since \mathcal{B} is point-countable and $|f^{-1}(y)| = 2$ for each $y \in Y$, we know that \mathcal{B}^* is point-countable in Y . Thus Y has a point-countable weak base.

A family \mathcal{P} of subsets of X is called a *wcs**-network of X if for any sequence $\{x_n\}_{n \in \mathbb{N}}$ which converges to a point x , and any open set U which contains x , there is some $P \in \mathcal{P}$ such that $P \subset U$ and $|\{n : x_n \in P\}| = \omega$ (cf. [14]). We know that every k -network of X is a *wcs**-network of X and if X is regular and \mathcal{P} is a *wcs**-network of X then $\{\overline{P} : P \in \mathcal{P}\}$ is a *cs**-network of X .

Example 8 ([18, Example 78]). Let \mathcal{T} be the usual Euclidean topology of \mathbb{R}^2 . Let $S_1 = \{(x, y) : x, y \in \mathbb{R}, y > 0\}$, $L = \{(x, 0) : x \in \mathbb{R}\}$ and $X = S_1 \cup L$. Let $\mathcal{T}^* = \{\mathcal{T}|X\} \cup \{\{x\} \cup (S_1 \cap U) : x \in L, x \in U \text{ and } U \in \mathcal{T}\}$, where $\mathcal{T}|X = \{U \cap X : U \in \mathcal{T}\}$. The space (X, \mathcal{T}^*) is a non-regular T_2 -space.

In [12, p. 28], it is pointed out that the space X which appears in Example 8 has a locally countable k -network but it has no point-countable *cs**-network. Thus it has a point-countable *wcs**-network but it has no point-countable *cs**-network.

Lemma 9. *If a regular space X is sequential and has a point-countable *wcs**-network, then X is a weakly monotonically monolithic space.*

PROOF: Let \mathcal{F} be a point-countable *wcs**-network of X . For any $A \subset X$, let $\mathcal{W}(A) = \{\overline{F} : F \cap A \neq \emptyset \text{ and } F \in \mathcal{F}\}$. We see that $\mathcal{W}(A)$ satisfies the conditions which appear in the Definition 3, since X is regular and sequential. \square

Similarly to Definition 2.11 from [19], we have:

Definition 10. Assume that Y is a weakly monotonically monolithic space and fix an operator \mathcal{W} which witnesses that. Let ϕ be any neighborhood assignment on X . For every $P \subset X$, we denote $P^* = \{x : x \in P \text{ and } P \subset \phi(x)\}$. For any open set $U \subset X$ say that a set $A \subset X$ is *U-saturated* if $P^* \subset \phi(A) \cup U$ for any $P \in \mathcal{W}(A)$, where $\phi(A) = \bigcup \{\phi(x) : x \in A\}$. If $U = \emptyset$ then *U-saturated* sets will be called *saturated*.

Lemma 11. *Let ϕ be any neighborhood assignment for X . If X is a weakly monotonically monolithic space, $A \subset X$ is a countable closed discrete subspace of X and U is any open subset of X , then there is a closed discrete subspace $B \subset X \setminus U \cup \phi(A)$, such that $A \cup B$ is U -saturated and $|B| \leq \omega$.*

PROOF: X is a weakly monotonically monolithic space. We fix an operator \mathcal{W} which witnesses that. Let $\mathcal{F}_0^* = \{P^* : P \in \mathcal{W}(A)\}$. Thus \mathcal{F}_0^* is a countable family. Enumerate it by prime numbers p . We take the first member F^* of \mathcal{F}_0^* such that $F^* \setminus U \cup \phi(A) \neq \emptyset$. We choose a point $x_1 \in F^* \setminus U \cup \phi(A)$. Then $F \subset \phi(x_1)$. The family $\mathcal{W}(A \cup \{x_1\})$ is countable. We denote $\mathcal{F}_1^* = \{F^* : F \in \mathcal{W}(A \cup \{x_1\}) \setminus \mathcal{W}(A)\}$. We enumerate \mathcal{F}_1^* by the squares p^2 of prime numbers.

Suppose we have finished n steps. We have $\phi(A)$, $U \cup \phi(A \cup \{x_1\})$, \dots , $U \cup \phi(A \cup \{x_1, \dots, x_n\})$, and families \mathcal{F}_i^* of subsets of X , the family \mathcal{F}_i^* is enumerated by the i -th powers of prime numbers for each $0 \leq i \leq n$.

If $U \cup \phi(A \cup \{x_1, \dots, x_n\}) = X$, then we stop the induction and let $B = \{x_i : 1 \leq i \leq n\}$. So we assume that $U \cup \phi(A \cup \{x_1, \dots, x_n\}) \neq X$. If $\bigcup\{\bigcup\mathcal{F}_i^* : 0 \leq i \leq n\}$ is contained in $U \cup \phi(A \cup \{x_1, \dots, x_n\})$ then we choose a point $x_{n+1} \in X \setminus U \cup \phi(A \cup \{x_1, \dots, x_n\})$. If $\bigcup\{\bigcup\mathcal{F}_i^* : 0 \leq i \leq n\} \setminus (U \cup \phi(A \cup \{x_1, \dots, x_n\})) \neq \emptyset$ then we take the first member F^* of $\bigcup\{\mathcal{F}_i^* : 0 \leq i \leq n\}$ such that $F^* \setminus U \cup \phi(A \cup \{x_1, \dots, x_n\}) \neq \emptyset$ and choose a point $x_{n+1} \in F^* \setminus U \cup \phi(A \cup \{x_1, \dots, x_n\})$. Thus $F^* \subset F \subset \phi(x_{n+1})$. We let $\mathcal{F}_{n+1}^* = \{F^* : F \in \mathcal{W}(A \cup \{x_1, \dots, x_{n+1}\}) \setminus \mathcal{W}(A \cup \{x_1, \dots, x_n\})\}$ and enumerate it by the $(n+1)$ -st powers of prime numbers.

In this way, we have a set $B = \{x_n : n \in \mathbb{N}\}$ and $B \subset X \setminus U \cup \phi(A)$. To prove that B is a closed discrete subspace of X , we only need to prove that B is closed since $x_m \notin \phi(x_n)$ whenever $m > n$.

Suppose $\overline{B} \setminus B \neq \emptyset$. Then there is some $x \in \overline{B} \setminus B$ such that $\mathcal{W}(B)$ is an external network of x . Thus there is some $P \in \mathcal{W}(B)$ such that $x \in P \subset \phi(x)$ and hence $x \in P^*$. Since $B = \bigcup\{\{x_1, \dots, x_n\} : n \in \mathbb{N}\} \subset \bigcup\{A \cup \{x_1, \dots, x_n\} : n \in \mathbb{N}\}$, we see that $P \in \mathcal{W}(\bigcup\{A \cup \{x_1, \dots, x_n\} : n \in \mathbb{N}\})$. So $P \in \mathcal{W}(A \cup \{x_1, \dots, x_n\})$ for some $n \in \mathbb{N}$. Thus $P^* \in \mathcal{F}_i^*$ for some $0 \leq i \leq n$. So there is some $m > n$ such that $P^* \subset U \cup \phi(A \cup \{x_1, \dots, x_m\})$. Thus $x \in \phi(A \cup \{x_1, \dots, x_m\})$. So $x \notin \overline{B}$. This contradicts the fact $x \in \overline{B}$. So B is closed and hence it is a closed discrete subspace of X and $B \subset X \setminus U \cup \phi(A)$, $|B| \leq \omega$.

For any $P \in \mathcal{W}(A \cup B)$, since $A \cup B = \bigcup\{A \cup \{x_1, \dots, x_n\} : n \in \mathbb{N}\}$, we have $P \in \mathcal{W}(A \cup \{x_1, \dots, x_n\})$ for some $n \in \mathbb{N}$. Hence $P^* \in \mathcal{F}_i^*$ for some $0 \leq i \leq n$. So P^* will be covered by $U \cup \phi(A \cup \{x_1, \dots, x_m\})$ for some $m > n$. Thus $A \cup B$ is U -saturated. \square

Lemma 12. *Let ϕ be any neighborhood assignment for X . If X is a weakly monotonically monolithic space and $A \subset X$ is a closed discrete subspace of X and U is any open subset of X , then there is a closed discrete subspace $D \subset X \setminus U \cup \phi(A)$, such that $A \cup D$ is U -saturated and $|D| \leq \max\{|A|, \omega\}$.*

PROOF: We will use induction on the cardinal $\kappa = |A|$. If $|A| \leq \omega$, then it is true by Lemma 11. Now assume that κ is an uncountable cardinal and our theorem is proved whenever U an open subset of X and $A \subset X$, $|A| < \kappa$.

Take an arbitrary set $A \subset X$ such that $|A| = \kappa$ and let $A = \{x_\alpha : \alpha < \kappa\}$. Let $A_\alpha = \{x_\beta : \beta < \alpha\}$ for each $\alpha \in [\omega, \kappa)$ and let $U' = U \cup \phi(A)$. Proceeding inductively assume that $\omega < \alpha < \kappa$ and we have a family $\{D_\beta : \omega \leq \beta < \alpha\}$ of closed discrete subspaces of X with the following properties:

- (1) $A_\omega \cup D_\omega$ is U' -saturated;
- (2) If $\omega < \beta < \alpha$ then $U_\beta = U' \cup \phi(\bigcup\{D_\gamma : \omega \leq \gamma < \beta\})$, $D_\beta \subset X \setminus U_\beta$ and $|D_\beta| \leq |\beta|$;
- (3) If $B_\beta = A_\beta \cup (\bigcup\{D_\gamma : \omega \leq \gamma < \beta\})$ then $B_\beta \cup D_\beta$ is U_β -saturated whenever $\omega < \beta < \alpha$;
- (4) The set $\bigcup\{D_\gamma : \omega \leq \gamma < \beta\}$ is a closed discrete subspace of X for each $\beta \in (\omega, \alpha)$.

We first prove that $\bigcup\{D_\beta : \omega \leq \beta < \alpha\}$ is a closed discrete subspace of X . We only need to prove that $\bigcup\{D_\beta : \beta < \alpha\}$ is a closed subspace of X . If $\alpha = \gamma + 1$, then $\bigcup\{D_\beta : \omega \leq \beta < \alpha\} = \bigcup\{D_\beta : \omega \leq \beta < \gamma\} \cup D_\gamma$. The sets D_γ and $\bigcup\{D_\beta : \omega \leq \beta < \gamma\}$ are closed discrete and hence $\bigcup\{D_\beta : \omega \leq \beta < \alpha\}$ is closed discrete. Let α be a limit ordinal. Suppose $\bigcup\{D_\beta : \omega \leq \beta < \alpha\}$ is not closed in X . Then there is some $x \in \overline{\bigcup\{D_\beta : \omega \leq \beta < \alpha\}} \setminus \bigcup\{D_\beta : \omega \leq \beta < \alpha\}$ such that $\mathcal{W}(\bigcup\{D_\beta : \omega \leq \beta < \alpha\})$ is an external network of the point x . So there is some $P \in \mathcal{W}(\bigcup\{D_\beta : \omega \leq \beta < \alpha\})$ such that $x \in P \subset \phi(x)$ and hence $x \in P^*$. The set $\bigcup\{D_\beta : \omega \leq \beta < \alpha\} \subset \bigcup\{A_{\beta_1} \cup (\bigcup\{D_\gamma : \omega \leq \gamma \leq \beta_1\}) : \omega \leq \beta_1 < \alpha\}$, so $P \in \mathcal{W}(A_{\beta_1} \cup (\bigcup\{D_\gamma : \omega \leq \gamma \leq \beta_1\}))$ for some $\beta_1 \in [\omega, \alpha)$.

The set $A_{\beta_1} \cup (\bigcup\{D_\gamma : \omega \leq \gamma \leq \beta_1\})$ is U_{β_1} -saturated, so $P^* \subset U_{\beta_1} \cup \phi(D_{\beta_1})$ and hence $x \in U' \cup \phi(A_{\beta_1} \cup (\bigcup\{D_\gamma : \omega \leq \gamma \leq \beta_1\}))$. We have $x \in \phi(\bigcup\{D_\gamma : \omega \leq \gamma \leq \beta_1\})$, since $x \in \overline{\bigcup\{D_\beta : \omega \leq \beta < \alpha\}} \setminus \bigcup\{D_\beta : \omega \leq \beta < \alpha\}$ and $U' \cap (\bigcup\{D_\beta : \omega \leq \beta < \alpha\}) = \emptyset$. Let γ_x be the smallest ordinal such that $x \in \phi(D_{\gamma_x})$. Since $x \in \overline{\bigcup\{D_\beta : \omega \leq \beta < \alpha\}}$, we see that $\gamma_x \neq \omega$. The set $\bigcup\{D_\gamma : \omega \leq \gamma < \gamma_x\}$ is a closed discrete subspace of X by induction and $x \notin \phi(\bigcup\{D_\gamma : \omega \leq \gamma < \gamma_x\})$. If $V_x = (X \setminus \bigcup\{D_\gamma : \omega \leq \gamma < \gamma_x\}) \cap \phi(D_{\gamma_x}) \cap O_x$, where $x \in O_x$ and $|O_x \cap D_{\gamma_x}| \leq 1$ for an open set O_x , then $|V_x \cap (\bigcup\{D_\beta : \omega \leq \beta < \alpha\})| \leq 1$. This contradicts $x \in \overline{\bigcup\{D_\beta : \omega \leq \beta < \alpha\}} \setminus \bigcup\{D_\beta : \omega \leq \beta < \alpha\}$. Thus $\bigcup\{D_\beta : \omega \leq \beta < \alpha\}$ is a closed discrete subspace of X .

Let $U_\alpha = U' \cup \phi(A_\alpha \cup (\bigcup\{D_\beta : \omega \leq \beta < \alpha\}))$, so $|A_\alpha \cup (\bigcup\{D_\beta : \omega \leq \beta < \alpha\})| \leq |\alpha| < \kappa$ and $A_\alpha \cup (\bigcup\{D_\beta : \omega \leq \beta < \alpha\}) \subset U_\alpha$. Thus by induction, there exists a closed discrete subspace $D_\alpha \subset X \setminus U_\alpha$ such that $(A_\alpha \cup (\bigcup\{D_\beta : \omega \leq \beta < \alpha\})) \cup D_\alpha$ is U_α -saturated.

If $D = \bigcup\{D_\alpha : \omega \leq \alpha < \kappa\}$ then $D \subset X \setminus U \cup \phi(A)$ and $|D| \leq \kappa$. In what follows, we show that D is a closed discrete subspace of X .

For any $\beta \in (\omega, \kappa)$, the set $\bigcup\{D_\alpha : \omega \leq \alpha < \beta\}$ is closed and $\phi(D_\beta) \cap D_\gamma = \emptyset$ for any $\gamma \in (\beta, \kappa)$. Thus to prove that D is a closed discrete subspace of X we only need to prove that D is closed in X . Suppose D is not closed; then there is some $x \in \overline{D} \setminus D$ such that $\mathcal{W}(D)$ is an external network of x . Since $\bigcup\{D_\gamma : \omega \leq \gamma < \beta\}$ is closed discrete for each $\beta < \kappa$, we can see that $x \notin \phi(D)$. There is some $P \in \mathcal{W}(D)$ such that $x \in P \subset \phi(x)$ and hence $x \in P^*$. The set

$D \subset \bigcup\{A_\alpha \cup (\bigcup\{D_\beta : \omega \leq \beta \leq \alpha\}) : \omega \leq \alpha < \kappa\}$, so there exists some $\gamma < \kappa$ such that $P \in \mathcal{W}(A_\gamma \cup (\bigcup\{D_\beta : \omega \leq \beta \leq \gamma\}))$. Let $U_\gamma = U' \cup \phi(A_\gamma \cup (\bigcup\{D_\beta : \omega \leq \beta < \gamma\}))$, we know that $A_\gamma \cup (\bigcup\{D_\beta : \omega \leq \beta < \gamma\}) \cup D_\gamma$ is U_γ -saturated. So $P^* \subset U' \cup \phi(A_\gamma \cup (\bigcup\{D_\beta : \omega \leq \beta \leq \gamma\}))$. Thus $x \in \phi(D)$. This contradicts the fact $x \notin \phi(D)$. So D is a closed discrete subspace of X and $D \subset X \setminus U'$.

Consider the set $A \cup D = \bigcup\{A_\alpha \cup (\bigcup\{D_\beta : \omega \leq \beta \leq \alpha\}) : \omega \leq \alpha < \kappa\}$. For any $P \in \mathcal{W}(A \cup D)$, there exists some $\alpha < \kappa$ such that $P \in \mathcal{W}(A_\alpha \cup (\bigcup\{D_\beta : \omega \leq \beta \leq \alpha\}))$. The set $A_\alpha \cup (\bigcup\{D_\beta : \omega \leq \beta \leq \alpha\})$ is U_α -saturated. Thus $P^* \subset U' \cup \phi(A_\alpha \cup (\bigcup\{D_\beta : \omega \leq \beta \leq \alpha\})) \subset U \cup \phi(A \cup D)$. Thus $A \cup D$ is U -saturated and $|D| \leq \max\{|A|, \omega\}$. \square

Corollary 13. *Let X be a space and ϕ be a neighborhood assignment for X . If X is a weakly monotonically monolithic space and A is a closed discrete subspace of X , then there is a closed discrete subspace $D \subset X \setminus \phi(A)$ such that $A \cup D$ is saturated.*

Theorem 14. *If X is a weakly monotonically monolithic space, then X is a D -space.*

PROOF: Let $|X| = \kappa$ and $X = \{x_\alpha : \alpha \in \kappa\}$. Let ϕ be any neighborhood assignment for X and $D_0 = \{x_0\}$. Then by Corollary 13 there is a closed discrete subspace $D'_1 \subset X \setminus \phi(D_0)$ such that $D_0 \cup D'_1$ is saturated. Let $D_1 = D_0 \cup D'_1$ if $x_1 \in \phi(D_0 \cup D'_1)$, otherwise $D_1 = \{x_1\} \cup D_0 \cup D'_1$.

Let $0 < \alpha < \kappa$ and assume we have closed discrete subspaces D_β and D'_β for each $\beta < \alpha$ such that:

- (1) $D'_0 = \emptyset$;
- (2) $D_{\beta_1} \subset D_{\beta_2}$ if $\beta_1 < \beta_2 < \beta$;
- (3) $x_\beta \in \phi(D_\beta)$;
- (4) $\bigcup\{D_\gamma : \gamma < \beta\}$ is closed discrete and $D'_\beta \subset X \setminus \phi(\bigcup\{D_\gamma : \gamma < \beta\})$ is such that $\bigcup\{D_\gamma : \gamma < \beta\} \cup D'_\beta$ is saturated;
- (5) $D_\beta = \bigcup\{D_\gamma : \gamma < \beta\} \cup D'_\beta$ if $x_\beta \in \phi(\bigcup\{D_\gamma : \gamma < \beta\} \cup D'_\beta)$, otherwise $D_\beta = \bigcup\{D_\gamma : \gamma < \beta\} \cup D'_\beta \cup \{x_\beta\}$.

In what follows, we will show that $\bigcup\{D_\beta : \beta < \alpha\}$ is a closed discrete subspace of X .

If $\alpha = \gamma + 1$ for some γ then $\bigcup\{D_\beta : \beta < \alpha\} = \bigcup\{D_\beta : \beta < \gamma\} \cup D_\gamma$ is a closed discrete subspace of X . If α is a limit ordinal and $\bigcup\{D_\beta : \beta < \alpha\}$ is not closed, then there is some $x \in \overline{\bigcup\{D_\beta : \beta < \alpha\}} \setminus \bigcup\{D_\beta : \beta < \alpha\}$ such that $\mathcal{W}(\bigcup\{D_\beta : \beta < \alpha\})$ is an external network of the point x . So there is some $P \in \mathcal{W}(\bigcup\{D_\beta : \beta < \alpha\})$ such that $x \in P \subset \phi(x)$ and hence $x \in P^*$. We know that $\mathcal{W}(\bigcup\{D_\beta : \beta < \alpha\}) = \bigcup\{\mathcal{W}(D_\beta) : \beta < \alpha\}$. So $P \in \mathcal{W}(D_{\beta_1})$ for some $\beta_1 < \alpha$. The set $D_{\beta_1} \cup D'_{\beta_1+1}$ is saturated. Thus $P^* \subset \phi(D_{\beta_1} \cup D'_{\beta_1+1})$ and hence $x \in \phi(D_{\beta_1+1})$. Thus $\phi(D_{\beta_1+1}) \cap D'_\gamma = \emptyset$ for any $\gamma \in (\beta_1 + 1, \kappa)$. The set D_{β_1+1} is a closed discrete subspace of X , so $x \notin \overline{\bigcup\{D_\beta : \beta < \alpha\}}$. This contradicts $x \in \overline{\bigcup\{D_\beta : \beta < \alpha\}}$. So $\bigcup\{D_\beta : \beta < \alpha\}$ is a closed discrete subspace of X .

Thus there is a closed discrete subspace $D'_\alpha \subset X \setminus \phi(\bigcup\{D_\beta : \beta < \alpha\})$ such that $\bigcup\{D_\beta : \beta < \alpha\} \cup D'_\alpha$ is saturated by Corollary 13. Let $D_\alpha = \bigcup\{D_\beta : \beta < \alpha\} \cup D'_\alpha$ if $x_\alpha \in \phi(\bigcup\{D_\beta : \beta < \alpha\} \cup D'_\alpha)$, otherwise $D_\alpha = \bigcup\{D_\beta : \beta < \alpha\} \cup D'_\alpha \cup \{x_\alpha\}$.

Let $D = \bigcup\{D_\alpha : \alpha < \kappa\}$. We see that $X = \bigcup\{\phi(d) : d \in D\}$. For each $x \in X$, let $\alpha_x = \min\{\alpha : x \in \phi(D_\alpha)\}$. Thus $\phi(D_{\alpha_x}) \cap D'_\beta = \emptyset$ for each $\beta \in (\alpha_x, \kappa)$. Let O_x be an open set of x such that $|O_x \cap D_{\alpha_x}| \leq 1$. Thus $|(O_x \cap \phi(D_{\alpha_x})) \cap D| \leq 1$. Thus D is a closed discrete subspace of X . So X is a D -space. \square

By Lemma 9 and Theorem 14, we have:

Corollary 15. *If a regular space X is sequential and has a point-countable wcs^* -network, then X is a D -space.*

Theorem 16. *Let X be a space and $\mathcal{F}(x)$ be a countable family of subsets of X for each $x \in X$. If for any non-closed subset $A \subset X$ there exists some point $x \in \overline{A} \setminus A$ such that for every open neighborhood U of x there exists some $y \in A$ and some $F \in \mathcal{F}(y)$ such that $x \in F \subset U$, then X is a D -space.*

PROOF: For any $A \subset X$, we let $\mathcal{W}(A) = \bigcup\{\mathcal{F}(a) : a \in A\}$. We can see that \mathcal{W} witnesses weak monotonic monolithicity of X . Thus X is a weakly monotonically monolithic space and hence X is a D -space by Theorem 14. \square

Recall that a space X satisfies *open* (G) if each point $x \in X$ has a countable neighborhood base \mathcal{B}_x such that whenever $x \in \overline{A}$ and $N(x)$ is a neighborhood of x , then there is an $a \in A$ and $B \in \mathcal{B}_a$ for which $x \in B \subset N(x)$.

By Theorem 16, we have:

Corollary 17 (cf. [9]). *Any space satisfying open (G) is a D -space.*

Corollary 18 (cf. [16]). *Let a space X have a point-countable family \mathcal{F} of subsets of X , such that for any non-closed subset $A \subset X$ there exists some point $x \in \overline{A} \setminus A$ such that for every open neighborhood U of x there exists some $F \in \mathcal{F}$ with $x \in F \subset U$ and $F \cap A \neq \emptyset$. Then X is a D -space.*

If X is a sequential space and $x \in W \subset X$ we say that W is a *weak-neighborhood* of x if whenever a sequence $\{x_n\}_{n \in \mathbb{N}}$ converges to x then $\{x_n\}_{n \in \mathbb{N}}$ is eventually in W . A collection \mathcal{W} of subsets of a sequential space X is said to be a \mathcal{W} -system for the topology if whenever $x \in U \subset X$, with U open, there exists a subcollection $\mathcal{V} \subset \mathcal{W}$ such that $x \in \bigcap \mathcal{V}$, $\bigcup \mathcal{V}$ is a weak-neighborhood of x and $\bigcup \mathcal{V} \subset U$ (cf. [4]).

Corollary 19 (cf. [4]). *A sequential space with a point-countable \mathcal{W} -system is a D -space.*

Corollary 20 (cf. [4] and [15]). *If X has a point-countable weak base, then X is a D -space.*

By Lemma 5 and Theorem 14, we have:

Corollary 21 (cf. [16]). *If X is a sequential space with a point-countable cs^* -network, then X is a D -space.*

Every k -network of X is a wcs^* -network of X , so by Lemma 9 and Theorem 14, we have:

Corollary 22 (cf. [17]). *If a regular space X is sequential and has a point-countable k -network, then X is a D -space.*

Every monotonically monolithic space is a weakly monotonically monolithic space and every subspace of monotonically monolithic space is monotonically monolithic. So we have:

Corollary 23 (cf. [19]). *If X is a monotonically monolithic space, then X is hereditarily a D -space.*

Acknowledgement. The proofs of the main results (Lemma 11, Lemma 12 and Theorem 14) depended heavily on the proofs from [19]. The author would like to thank the referee for his (or her) valuable remarks and suggestions which greatly improved the paper.

REFERENCES

- [1] Arhangel'skii A.V., *D-spaces and finite unions*, Proc. Amer. Math. Soc. **132.7** (2004), 2163–2170.
- [2] Arhangel'skii A.V., Buzyakova R.Z., *Addition theorems and D-spaces*, Comment. Math. Univ. Carolin. **43.4** (2002), 653–663.
- [3] Borges C.R., Wehrly A.C., *A study of D-spaces*, Topology Proc. **16** (1991), 7–15.
- [4] Burke D.K., *Weak-base and D-space*, Comment. Math. Univ. Carolin. **48.2** (2007), 281–289.
- [5] Buzyakova R.Z., *Hereditary D-property of function spaces over compacta*, Proc. Amer. Math. Soc. **132.11** (2004), 3433–3439.
- [6] van Douwen E.K., Pfeffer W.F., *Some properties of the Sorgenfrey line and related spaces*, Pacific J. Math. **81.2** (1979), 371–377.
- [7] Engelking R., *General Topology*, Sigma Series in Pure Mathematics, 6, Heldermann, Berlin, revised ed., 1989.
- [8] Fleissner W.G., Stanley A.M., *D-spaces*, Topology Appl. **114.3** (2001), 261–271.
- [9] Gruenhage G., *A note on D-spaces*, Topology Appl. **153** (2006), 229–240.
- [10] Gruenhage G., *Generalized metric spaces*, in Handbook of Set-theoretic Topology, K. Kunen and J. Vaughan (Eds), North-Holland, Amsterdam, 1984, pp. 423–501.
- [11] Gruenhage G., Michael E., Tanaka Y., *Spaces determined by point-countable covers*, Pacific J. Math. **113.2** (1984), 303–332.
- [12] Lin S., *Point-countable Covers and Sequence-covering Mappings*, Chinese Science Press, Beijing, 2002.
- [13] Lin S., Liu C., *On spaces with point-countable cs^* -networks*, Topology Appl. **74** (1996), 51–60.
- [14] Lin S., Tanaka Y., *Point-countable k -network, closed maps and related results*, Topology Appl. **59** (1994), 79–86.
- [15] Peng L.-X., *The D-property of some Lindelöf spaces and related conclusions*, Topology Appl. **154** (2007), 469–475.
- [16] Peng L.-X., *A special point-countable family that makes a space to be a D-space*, Adv. Math. (China) **37.6** (2008), 724–728.
- [17] Peng L.-X., *A note on D-spaces and infinite unions*, Topology Appl. **154** (2007), 2223–2227.

- [18] Steen L.A., Seebach J.A., Jr., *Counterexamples in Topology*, second edition, Springer, New York-Heidelberg, 1978.
- [19] Tkachuk V.V., *Monolithic spaces and D-spaces revised*, *Topology Appl.* **156** (2009), 840–846.

COLLEGE OF APPLIED SCIENCE, BEIJING UNIVERSITY OF TECHNOLOGY,
BEIJING 100124, CHINA

Email: pengliangxue@bjut.edu.cn

(Received January 7, 2009, revised January 26, 2010)