

Continua with unique symmetric product

JOSÉ G. ANAYA, ENRIQUE CASTAÑEDA-ALVARADO, ALEJANDRO ILLANES

Abstract. Let X be a metric continuum. Let $F_n(X)$ denote the hyperspace of nonempty subsets of X with at most n elements. We say that the continuum X has *unique hyperspace* $F_n(X)$ provided that the following implication holds: if Y is a continuum and $F_n(X)$ is homeomorphic to $F_n(Y)$, then X is homeomorphic to Y . In this paper we prove the following results: (1) if X is an indecomposable continuum such that each nondegenerate proper subcontinuum of X is an arc, then X has unique hyperspace $F_2(X)$, and (2) let X be an arcwise connected continuum for which there exists a unique point $v \in X$ such that v is the vertex of a simple triod. Then X has unique hyperspace $F_2(X)$.

Keywords: arc continuum; continuum; indecomposable; symmetric product; unique hyperspace

Classification: Primary 54B20; Secondary 54F15

1. Introduction

A *continuum* is a compact connected metric space with more than one point. Given a continuum X and a positive integer n , we consider the following hyperspaces of X :

$$\begin{aligned} 2^X &= \{A \subset X : A \text{ is closed and nonempty}\}, \\ C(X) &= \{A \in 2^X : A \text{ is connected}\}, \\ C_n(X) &= \{A \in 2^X : A \text{ has at most } n \text{ components}\}, \\ F_n(X) &= \{A \in 2^X : A \text{ has at most } n \text{ points}\}. \end{aligned}$$

All these hyperspaces are considered with the Hausdorff metric H_X [13, Theorem 2.2].

The hyperspace $F_n(X)$ is called the *n^{th} -symmetric product of X* .

Let $\mathcal{H}(X)$ denote one of the hyperspaces 2^X , $C(X)$, $C_n(X)$ or $F_n(X)$. We say that the continuum X has *unique hyperspace* $\mathcal{H}(X)$ provided that the following implication holds: if Y is a continuum and $\mathcal{H}(X)$ is homeomorphic to $\mathcal{H}(Y)$, then X is homeomorphic to Y .

The topic of this paper is inserted in the following general problem.

Problem 1. Find conditions on the continuum X in order that X has unique hyperspace $\mathcal{H}(X)$.

Problem 1 has been widely studied by a number of authors. The recently published paper [10] contains a detailed survey on this subject. For $n \geq 4$, the most general result on uniqueness of symmetric products is Theorem 5 of [5] that says that if $n \geq 4$ and X is a wired continuum, then X has unique hyperspace $F_n(X)$. Since the class of wired continua includes finite graphs, dendrites with closed set of end points, fans, compactifications of the ray $[0, \infty)$, compactifications of the real line and indecomposable arc continua, we have that Theorem 5 of [5] generalizes previous results in [1], [3], [6], [7], [9] and [12].

A continuum X is *indecomposable* provided that X is not the union of two proper subcontinua. An *arc continuum* is a continuum such that all its nondegenerate proper subcontinua are arcs. A *simple triod* is a continuum Y such that $Y = L_1 \cup L_2 \cup L_3$ where each L_i is an arc, and there exists a point v in Y such that v is an end point of each L_i and if $i \neq j$, then $L_i \cap L_j = \{v\}$. The point v is called the *vertex* of Y .

In [10, Question 42] it is asked if indecomposable arc continua have unique hyperspaces $F_2(X)$ and $F_3(X)$ (see also [5, Question 27]).

In this paper we prove the following theorems.

Theorem. *Let X be an indecomposable arc continuum. Then X has unique hyperspace $F_2(X)$.*

Theorem. *Let X be an arcwise connected continuum for which there exists a unique point $v \in X$ which is the vertex of a simple triod. Then X has unique hyperspace $F_2(X)$.*

The following questions remain open [10, Question 42].

Question 2. *Let X be an indecomposable arc continuum. Does X have unique hyperspace $F_3(X)$?*

Question 3. *Let X be an arcwise connected continuum for which there exists a unique point $v \in X$ which is the vertex of a simple triod. Does X have unique hyperspace $F_3(X)$?*

2. Indecomposable arc continua

We need the following conventions.

Given a continuum X , $A \subset X$, $x \in X$ and $\varepsilon > 0$, let $B(x, \varepsilon)$ be the open ε -ball around x in X and $N(A, \varepsilon) = \bigcup\{B(x, \varepsilon) \subset X : x \in A\}$.

Given a topological space Y and a point $e \in Y$. The point e is an *end point* of Y provided that e is an end point of each arc in Y containing e . Let

$$F_1(Y) = \{\{y\} : y \in Y\},$$

$$E(Y) = \{e \in Y : e \text{ is an end point of } Y\} \text{ and}$$

$$\mathcal{AC}(Y) = \{K \subset Y : K \text{ is an arc component of } Y\}.$$

We consider the following families of continua:

$$\begin{aligned} \mathfrak{A} &= \{X : X \text{ is a continuum and there exists } \varepsilon > 0 \text{ such that if } A \text{ is} \\ &\quad \text{a subcontinuum of } X \text{ and } 0 < \text{diameter}(A) < \varepsilon, \text{ then } A \text{ is an arc}\}, \\ \mathfrak{DC} &= \{X : X \text{ is a continuum and for each } K \in \mathcal{AC}(X), K \text{ is dense in } X\}, \\ \mathfrak{W} &= \{X \in \mathfrak{A} \cap \mathfrak{DC} : X \text{ has uncountably many arc components}\}. \end{aligned}$$

Given $X \in \mathfrak{W}$, let

$$\mathcal{N}\partial F_2(X) = \{A \in F_2(X) : A \text{ does not belong to the manifold interior of a 2-cell } \mathcal{M} \text{ that is contained in } F_2(X)\}.$$

By [14, Theorem 11.15], it follows that each indecomposable arc continuum belongs to \mathfrak{W} .

Given a continuum X and subsets J, L of X , let

$$\langle J, L \rangle = \{A \in F_2(X) : A \subset J \cup L, A \cap J \neq \emptyset \text{ and } A \cap L \neq \emptyset\}.$$

Lemma 4. *Let X be a continuum. Then $\mathcal{AC}(F_2(X)) = \{\langle J, L \rangle : J, L \in \mathcal{AC}(X)\}$.*

PROOF: (Compare with [5, Lemma 22]). Let $J, L \in \mathcal{AC}(X)$. We take $\{x, y\}, \{u, v\} \in \langle J, L \rangle$, where $x, u \in J$ and $y, v \in L$ (here, it is possible that $J = L, x = y$ or $u = v$). Since J and L are arc components of X , there exist arcs α and β (possibly degenerate) such that $x, u \in \alpha \subset J$ and $y, v \in \beta \subset L$. Notice that $\{\{x, r\} : r \in \beta\} \subset \langle J, L \rangle$ and $\{\{s, v\} : s \in \alpha\} \subset \langle J, L \rangle$ are homeomorphic to β and α , respectively and the element $\{x, v\}$ belongs to both sets. Thus, $\{\{x, r\} : r \in \beta\} \cup \{\{s, v\} : s \in \alpha\}$ is an arcwise connected subset of $\langle J, L \rangle$ that contains $\{x, y\}$ and $\{u, v\}$. We have shown that $\langle J, L \rangle$ is arcwise connected.

Let \mathcal{D} be the arc component of $F_2(X)$ containing $\langle J, L \rangle$. Fix an element $\{x, y\} \in \langle J, L \rangle$, where $x \in J$ and $y \in L$; and let $\{u, v\} \in \mathcal{D}$. Let Δ be an (possibly degenerate) arc in $F_2(X)$ such that $\{x, y\}, \{u, v\} \in \Delta$. Let $B = \bigcup\{A : A \in \Delta\}$. By [5, Lemma 1], B has at most two components. Moreover, if $B = B_1 \cup B_2$, where B_1 and B_2 are the components of B (possibly $B_1 = B_2$), we may assume that $x, u \in B_1$ and $y, v \in B_2$. By [4, Lemma 2.2], B_1 and B_2 are locally connected continua and then they are arcwise connected. Thus, $B_1 \subset J$ and $B_2 \subset L$. This proves that $\{u, v\} \in \langle J, L \rangle$. We have shown that $\mathcal{D} = \langle J, L \rangle$. Therefore, $\langle J, L \rangle$ is an arc component of $F_2(X)$.

Clearly, $F_2(X) = \bigcup\{\langle J, L \rangle : J, L \in \mathcal{AC}(X)\}$. Therefore, $\mathcal{AC}(F_2(X)) = \{\langle J, L \rangle : J, L \in \mathcal{AC}(X)\}$. □

Corollary 5. *Let X be a continuum. Then X is arcwise connected if and only if $F_2(X)$ is arcwise connected.*

Lemma 6. *Let $X \in \mathfrak{W}$ and $A \in F_2(X)$. Then $A \in \mathcal{N}\partial F_2(X)$ if and only if $A \in F_1(X)$ or $A \cap E(X) \neq \emptyset$.*

PROOF: (Necessity). Suppose that $A = \{x, y\}$, where $x \neq y$ and $A \cap E(X) = \emptyset$. Then there exist arcs E and F in X such that $x \in E$, $y \in F$, x is not an end point of E and y is not an end point of F . Shortening E and F , if necessary, we may assume that $E \cap F = \emptyset$. It is easy to show that the map $\varphi : E \times F \rightarrow \langle E, F \rangle$ given by $\varphi(u, v) = \{u, v\}$ is a homeomorphism. Thus $\langle E, F \rangle$ is a 2-cell containing $\{x, y\}$ in its interior as manifold. This proves that $A \notin \mathcal{N}\partial F_2(X)$ and completes the proof of the necessity.

(Sufficiency). Let d be a metric for X . Since $X \in \mathfrak{A}$, there exists $\varepsilon > 0$ such that for each subcontinuum B of X with $0 < \text{diameter}(B) < \varepsilon$, we have that B is an arc. Suppose that $A \notin \mathcal{N}\partial F_2(X)$. Then there exists a 2-cell \mathcal{M} in $F_2(X)$ such that A is in the manifold interior of \mathcal{M} . Taking a smaller 2-cell contained in \mathcal{M} , if necessary, we may assume that $\text{diameter}(\mathcal{M}) < \frac{\varepsilon}{2}$. Let $B = \bigcup\{D \in F_2(X) : D \in \mathcal{M}\}$. We need to show that $A \notin F_1(X)$ and $A \cap E(X) = \emptyset$.

Suppose, first, that $A \in F_1(X)$. Then $A = \{x\}$ for some $x \in X$. By [5, Lemma 1], B is a subcontinuum of X . Since $\text{diameter}(\mathcal{M}) < \frac{\varepsilon}{2}$, $H_X(D, \{x\}) < \frac{\varepsilon}{2}$ for each $D \in \mathcal{M}$. Thus, $B \subset N(\{x\}, \frac{\varepsilon}{2})$ and $\text{diameter}(B) < \varepsilon$. Hence, B is an arc. Thus, (see [11, Section 13]) there exists a homeomorphism $h : F_2(B) \rightarrow [0, 1] \times [0, 1]$ such that $h(\{x\}) = (0, 0)$. Notice that $\mathcal{M} \subset F_2(B)$, so $h(\mathcal{M}) \subset [0, 1] \times [0, 1]$ and $h(\mathcal{M})$ is a 2-cell containing $(0, 0)$ in its manifold interior. This contradicts the Invariance of Domain Theorem [8, Theorem VI 9] and completes the proof that $A \notin F_1(X)$.

Now, suppose that $A \cap E(X) \neq \emptyset$. Then $A = \{v, x\}$ for some $v \in E(X)$ and $v \neq x$. Let $\delta = \min\{\frac{d(v,x)}{2}, \frac{\varepsilon}{2}\} > 0$. In this case, we may assume that $\text{diameter}(\mathcal{M}) < \delta$. Since $B \subset N(A, \delta) = B(v, \delta) \cup B(x, \delta)$ and $v, x \in B$, by [5, Lemma 1], we have that B has exactly two components E and F , where $E \subset B(v, \delta)$ and $F \subset B(x, \delta)$ and, for each $D \in \mathcal{M}$, $D \cap E \neq \emptyset$ and $D \cap F \neq \emptyset$. This implies that $\mathcal{M} \subset \langle E, F \rangle$. By the choice of ε , E and F are (possibly degenerate) arcs. Since $A \in \mathcal{M}$, $A \in \langle E, F \rangle$, so $v \in E$ and $x \in F$. Thus, v is an end point of E . Let $g : E \times F \rightarrow \langle E, F \rangle$ be given by $g(s, t) = \{s, t\}$. Clearly, g is a homeomorphism. Hence, $\langle E, F \rangle$ is a 2-cell (or an arc) having $A = \{v, x\}$ in its manifold boundary and \mathcal{M} is a 2-cell such that A belongs to the manifold interior of \mathcal{M} and $\mathcal{M} \subset \langle E, F \rangle$. This contradicts again the Invariance of Domain Theorem [8, Theorem VI 9] and completes the proof of the sufficiency. \square

Lemma 7. *Let $X \in \mathfrak{W}$ and $\mathcal{K} \in \mathcal{AC}(\mathcal{N}\partial F_2(X))$. Then \mathcal{K} is one of the forms described in (1)–(4).*

- (1) $\mathcal{K} = F_1(L)$ for some $L \in \mathcal{AC}(X)$ such that $L \cap E(X) = \emptyset$.
- (2) $\mathcal{K} = F_1(L) \cup \{\{v, x\} \in F_2(X) : x \in L\}$ for some $L \in \mathcal{AC}(X)$ and $v \in E(X) \cap L$.
- (3) $\mathcal{K} = \{\{v, x\} \in F_2(X) : x \in L\}$ for some $L \in \mathcal{AC}(X)$ and $v \in E(X) - L$.
- (4) $\mathcal{K} = \{\{v, x\} \in F_2(X) : x \in J\} \cup \{\{w, x\} \in F_2(X) : x \in L\}$ for some $L, J \in \mathcal{AC}(X)$, $v \in E(X) \cap L$, $w \in E(X) \cap J$ and $L \neq J$.

PROOF: First, we will see that for each $J \in \mathcal{AC}(X)$, $|J \cap E(X)| \leq 1$. Suppose to the contrary that there exist two different elements $v, w \in J \cap E(X)$. Since

J is arcwise connected, there exists an arc $\alpha \subset J$ with end points v and w . We claim that $J = \alpha$. Suppose that there exists a point $x \in J - \alpha$. Since J is arcwise connected, there exists an arc β such that $\beta \cap \alpha = \{y\}$ for some $y \in J$. If $y = v$, then $\alpha \cup \beta$ is an arc in X , $v \in \alpha \cup \beta$ and v is not an end point of $\alpha \cup \beta$. This contradicts the fact that $v \in E(X)$. Thus, $y \neq v$. Similarly, $y \neq w$. Hence, $y \in \alpha - \{v, w\}$ and y is the vertex of the simple triod $\alpha \cup \beta$ in X . Since $X \in \mathfrak{W}$, there exists $\varepsilon > 0$ such that if $B \in C(X)$ and $0 < \text{diameter}(B) < \varepsilon$, then B is an arc. Since $\alpha \cup \beta$ contains simple triods of diameter less than ε , we obtain a contradiction. This completes the proof that $J = \alpha$. Since $X \in \mathfrak{W}$, J is dense in X , so $X = \alpha$. This contradicts the fact that X has uncountably many arc components ($X \in \mathfrak{W}$). We have shown that $|J \cap E(X)| \leq 1$.

By Lemma 4, $\mathcal{AC}(F_2(X)) = \{\langle J, L \rangle \subset F_2(X) : J, L \in \mathcal{AC}(X)\}$. Thus, there exist $J, L \in \mathcal{AC}(X)$ such that $\mathcal{K} \subset \langle J, L \rangle$. We consider five cases.

Case 1. $J = L$ and $J \cap E(X) = \emptyset$.

In this case, $\mathcal{K} \subset \langle J, L \rangle = F_2(J)$. By Lemma 6, $F_2(J) \cap \mathcal{N}\partial F_2(X) = F_1(J)$. Since $F_1(J)$ is homeomorphic to J , $F_1(J)$ is arcwise connected. Since $\mathcal{K} \subset F_2(J) \cap \mathcal{N}\partial F_2(X)$, we conclude that $\mathcal{K} = F_1(J)$ and \mathcal{K} is as in (1).

Case 2. $J = L$ and $J \cap E(X) \neq \emptyset$.

In this case, $\mathcal{K} \subset \langle J, L \rangle = F_2(J)$. Let $v \in J \cap E(X)$. Then $J \cap E(X) = \{v\}$. By Lemma 6, $F_2(J) \cap \mathcal{N}\partial F_2(X) = F_1(J) \cup \{\{v, x\} : x \in J\}$. Since the function $x \rightarrow \{v, x\}$, from J onto $\{\{v, x\} : x \in J\}$ is continuous, $\{\{v, x\} : x \in J\}$ is arcwise connected and intersects $F_1(J)$ in the element $\{v\}$. Thus, $F_1(J) \cup \{\{v, x\} : x \in J\}$ is arcwise connected and $\mathcal{K} = F_1(J) \cup \{\{v, x\} : x \in J\}$. Hence, \mathcal{K} is as in (2).

Case 3. $J \neq L$ and $(J \cup L) \cap E(X) = \emptyset$.

By Lemma 6, $\langle J, L \rangle \cap \mathcal{N}\partial F_2(X) = \emptyset$. Since $\mathcal{K} \subset \langle J, L \rangle \cap \mathcal{N}\partial F_2(X)$, we conclude that this case is impossible.

Case 4. $J \neq L$, $J \cap E(X) \neq \emptyset$ and $L \cap E(X) = \emptyset$.

Let $v \in J$ be such that $J \cap E(X) = \{v\}$. By Lemma 6, $\langle J, L \rangle \cap \mathcal{N}\partial F_2(X) = \{\{v, x\} : x \in L\}$. Since this set is arcwise connected, we obtain that $\mathcal{K} = \{\{v, x\} : x \in L\}$ and \mathcal{K} is as in (3).

Case 5. $J \neq L$, $J \cap E(X) \neq \emptyset$ and $L \cap E(X) \neq \emptyset$.

Let $v \in L$ and $w \in J$ be such that $L \cap E(X) = \{v\}$ and $J \cap E(X) = \{w\}$. By Lemma 6, $\langle J, L \rangle \cap \mathcal{N}\partial F_2(X) = \{\{v, x\} \in F_2(X) : x \in J\} \cup \{\{w, x\} \in F_2(X) : x \in L\}$. Since both sets in this union are arcwise connected and they meet in the element $\{v, w\}$, we obtain that $\mathcal{K} = \{\{v, x\} \in F_2(X) : x \in J\} \cup \{\{w, x\} \in F_2(X) : x \in L\}$ is as in (4).

This completes the proof of the lemma. □

Lemma 8. *Let X be an indecomposable arc continuum and let Y be a continuum such that $F_2(X)$ is homeomorphic to $F_2(Y)$. Then $Y \in \mathfrak{A}$.*

PROOF: Let d_Y be a metric for Y . Let $h : F_2(Y) \rightarrow F_2(X)$ be a homeomorphism. Let $\delta = \text{diameter}(X)$. Then there exists $\varepsilon > 0$ such that if $A, B \in F_2(Y)$ and $H_Y(A, B) < \varepsilon$, then $H_X(h(A), h(B)) < \frac{\delta}{5}$.

Let $Z \in C(Y)$ be such that $0 < \text{diameter}(Z) < \varepsilon$.

Fix an element $A_0 \in F_2(Z)$. Suppose that $h(A_0) = \{r, s\}$, where $r = s$ in the case that $h(A_0)$ is a one-point set. Given $B \in F_2(Z)$. For each $a \in A_0$ and each $b \in B$, we have that $a, b \in Z$ and $d_Y(a, b) < \varepsilon$. This implies that $H_Y(A_0, B) < \varepsilon$. Thus, $H_X(h(A_0), h(B)) < \frac{\delta}{5}$ and $h(B) \subset N(h(A_0), \frac{\delta}{5}) = B(r, \frac{\delta}{5}) \cup B(s, \frac{\delta}{5})$.

Let $D = \bigcup \{h(B) \in F_2(X) : B \in F_2(Z)\}$. By the previous paragraph, $D \subset B(r, \frac{\delta}{5}) \cup B(s, \frac{\delta}{5})$. By [5, Lemma 1], D has at most two components. Let D_1, D_2 be the components of D , where $D_1 = D_2$ in the case that D is connected. In the case that $D_1 \neq D_2$, each element of $h(F_2(Z))$ intersects both sets D_1 and D_2 ([5, Lemma 1]). So, in this case, $h(F_2(Z)) \subset \langle D_1, D_2 \rangle$. In the case that $D_1 = D_2$, $h(F_2(Z)) \subset F_2(D)$.

We claim that $D \neq X$. Suppose to the contrary that $D = X$. Then $X = B(r, \frac{\delta}{5}) \cup B(s, \frac{\delta}{5})$. By the connectedness of X , $B(r, \frac{\delta}{5}) \cap B(s, \frac{\delta}{5}) \neq \emptyset$. This implies that $\text{diameter}(X) \leq \frac{4\delta}{5} < \delta$, which is a contradiction. We have shown that $D \neq X$. Since X is an arc continuum, each set D_1 and D_2 is either an arc or a one-point set.

In the case that $D_1 = D_2$, $F_2(D)$ is a 2-cell ([11, Section 13]) and $F_2(Z)$ can be embedded in $F_2(D)$. In the case that $D_1 \neq D_2$, $\langle D_1, D_2 \rangle$ is homeomorphic to $D_1 \times D_2$, and then $\langle D_1, D_2 \rangle$ is either an arc or a 2-cell. In both cases, $F_2(Z)$ can be embedded in a 2-cell. By [2, Theorem 5], Z is an arc.

We have shown that $Y \in \mathfrak{A}$. □

Theorem 9. *Let X be an indecomposable arc continuum. Then X has unique hyperspace $F_2(X)$.*

PROOF: Let Y be a continuum and let $h : F_2(Y) \rightarrow F_2(X)$ be a homeomorphism.

By Lemma 4, $\mathcal{AC}(F_2(X)) = \{\langle J, L \rangle \subset F_2(X) : J, L \in \mathcal{AC}(X)\}$. By [14, Theorem 11.15] it follows that $\mathcal{AC}(X)$ is uncountable. This implies that $\mathcal{AC}(F_2(X))$ is uncountable. Since h is a homeomorphism, $\mathcal{AC}(F_2(Y)) = \{h^{-1}(K) : K \in \mathcal{AC}(F_2(X))\}$. Thus, $\mathcal{AC}(F_2(Y)) = \{\langle J, L \rangle \subset F_2(Y) : J, L \in \mathcal{AC}(Y)\}$ is uncountable. Hence, $\mathcal{AC}(Y)$ is uncountable.

Given $K \in \mathcal{AC}(Y)$, we will see that K is dense in Y . Let U be a nonempty open subset of Y . By Lemma 4, $\langle K \rangle$ is an arc component of $F_2(Y)$. Thus, $h(\langle K \rangle)$ is an arc component of $F_2(X)$. Applying Lemma 4 again, we have that there exist $J, L \in \mathcal{AC}(X)$ such that $h(\langle K \rangle) = \langle J, L \rangle$. Since $\langle U \rangle$ is nonempty and open in $F_2(Y)$, $h(\langle U \rangle)$ is nonempty and open in $F_2(X)$. Since $F_2(X) - F_1(X)$ is dense in $F_2(X)$, there exist $u, x \in X$ such that $\{u, x\} \in h(\langle U \rangle)$ and $u \neq x$. Then there exists $\varepsilon > 0$ such that if $A \in F_2(X)$ and $H_X(A, \{u, x\}) < \varepsilon$, then $A \in h(\langle U \rangle)$ and $\varepsilon < \frac{d(u, x)}{2}$. Since J and L are dense in X , there exist points $r \in B(u, \varepsilon) \cap J$ and $s \in B(x, \varepsilon) \cap L$. Then $H_X(\{r, s\}, \{u, x\}) < \varepsilon$ and $\{r, s\} \in h(\langle U \rangle) \cap \langle J, L \rangle$. Thus, $h^{-1}(\{r, s\}) \in \langle U \rangle \cap \langle K \rangle$ and $h^{-1}(\{r, s\}) \subset U \cap K$. Hence, $U \cap K \neq \emptyset$. Therefore, K is dense in Y .

By Lemma 8, we conclude that $Y \in \mathfrak{W}$.

Since $\mathcal{N}\partial F_2(X)$ is defined in terms of topological properties of $F_2(X)$, we obtain that $h(\mathcal{N}\partial F_2(Y)) = \mathcal{N}\partial F_2(X)$. Hence, for each $\mathcal{K} \in \mathcal{AC}(\mathcal{N}\partial F_2(Y))$, $h(\mathcal{K}) \in \mathcal{AC}(\mathcal{N}\partial F_2(X))$ and $h(\text{cl}_{F_2(Y)}(\mathcal{K})) = \text{cl}_{F_2(X)}(h(\mathcal{K}))$.

Given $L \in \mathcal{AC}(Y)$ and $v \in E(X)$, since L is dense in Y , $\text{cl}_{F_2(Y)}(F_1(L)) = F_1(Y)$ and $\text{cl}_{F_2(Y)}(\{\{v, x\} \in F_2(Y) : x \in L\}) = \{\{v, x\} \in F_2(Y) : x \in Y\}$ is homeomorphic to Y .

Notice that an element Z of \mathfrak{W} does not have cut points since a cut point belongs to each dense arc component and elements in \mathfrak{W} have uncountably many dense arc components.

Given $\mathcal{K} \in \mathcal{AC}(\mathcal{N}\partial F_2(Y))$, we have that \mathcal{K} is of one of the forms described in Lemma 7. In the case that \mathcal{K} is of the form (1), then $\text{cl}_{F_2(Y)}(\mathcal{K})$ is homeomorphic to Y ; if \mathcal{K} is of the form (2), then $\text{cl}_{F_2(Y)}(\mathcal{K})$ is homeomorphic to two copies of Y joined by a point (the element $\{v\}$); if \mathcal{K} is of the form (3), then $\text{cl}_{F_2(Y)}(\mathcal{K})$ is homeomorphic to Y ; and if \mathcal{K} is of the form (4), then $\text{cl}_{F_2(Y)}(\mathcal{K})$ is homeomorphic to two copies of Y joined by a point (the element $\{v, w\}$).

Fix $\mathcal{K} \in \mathcal{AC}(\mathcal{N}\partial F_2(Y))$. Then $\text{cl}_{F_2(Y)}(\mathcal{K})$ is homeomorphic to $\text{cl}_{F_2(X)}(h(\mathcal{K}))$. By the previous paragraph, $\text{cl}_{F_2(Y)}(\mathcal{K})$ (resp., $\text{cl}_{F_2(X)}(h(\mathcal{K}))$) is either homeomorphic to Y (resp., X) or homeomorphic to two copies of Y (resp., X) joined by a point. Since elements in \mathfrak{W} does not have cut points, it is not possible that $\text{cl}_{F_2(Y)}(\mathcal{K})$ is homeomorphic to Y and $\text{cl}_{F_2(X)}(h(\mathcal{K}))$ is homeomorphic to two copies of X joined by a point; and it is not possible that $\text{cl}_{F_2(X)}(h(\mathcal{K}))$ is homeomorphic to X and $\text{cl}_{F_2(Y)}(\mathcal{K})$ is homeomorphic to two copies of Y joined by a point.

Therefore, we only have two possibilities:

- (a) $\text{cl}_{F_2(Y)}(\mathcal{K})$ is homeomorphic to Y and $\text{cl}_{F_2(X)}(h(\mathcal{K}))$ is homeomorphic to X ; or
- (b) $\text{cl}_{F_2(Y)}(\mathcal{K})$ is homeomorphic to two copies of Y joined by a point and $\text{cl}_{F_2(X)}(h(\mathcal{K}))$ is homeomorphic to two copies of X joined by a point.

Clearly, each of the statements (a) and (b) implies that X is homeomorphic to Y . □

Corollary 10. *The Buckethandle continuum X has unique hyperspace $F_2(X)$.*

Corollary 11. *Each solenoid X has unique hyperspace $F_2(X)$.*

3. Arcwise connected continua

Theorem 12. *Let X be an arcwise connected continuum for which there exists a unique point $v_0 \in X$ such that v_0 is the vertex of a simple triod. Then X has unique hyperspace $F_2(X)$.*

PROOF: Let Y be a continuum and let $h : F_2(X) \rightarrow F_2(Y)$ be a homeomorphism. Define

$\mathcal{A}(X) = \{A \in F_2(X) : \text{for each neighborhood } \mathcal{U} \text{ of } A \text{ in } F_2(X) \text{ there exists a locally connected subcontinuum } \mathcal{M} \text{ of } F_2(X) \text{ such that } \mathcal{M} \subset \mathcal{U} \text{ and } \mathcal{M} \text{ is not embeddable in a 2-manifold}\}.$

Since $\mathcal{A}(X)$ is defined only using topological properties, we conclude that $h(\mathcal{A}(X)) = \mathcal{A}(Y)$.

Given an arcwise connected continuum Z , let $V(Z) = \{z \in Z : z \text{ is the vertex of a simple triod contained in } Z\}$.

Claim 1. Let Z be an arcwise connected continuum. Then

- (a) $\{A \in F_2(Z) : A \cap V(Z) \neq \emptyset\} \subset \mathcal{A}(Z)$,
- (b) if $A \in F_2(Z)$ and $A \cap \text{cl}_Z(V(Z)) = \emptyset$, then $A \notin \mathcal{A}(Z)$.

In order to prove (a), take $A \in F_2(Z)$ and $v \in V(Z)$ such that $v \in A$. Let \mathcal{U} be a neighborhood of A in $F_2(Z)$. If $A = \{v\}$, we can take an element $z \in Z - \{v\}$ such that $B = \{v, z\} \in \text{int}_{F_2(Z)}(\mathcal{U})$. In the case that $A \neq \{v\}$, $A = \{v, z\}$ for some $z \in Z - \{v\}$ and put $B = A$. In both cases, there exists $B = \{v, z\} \in \text{int}_{F_2(Z)}(\mathcal{U})$, where $v \neq z$. Let T be a simple triod in Z such that v is the vertex of T and $z \notin T$. Since Z is arcwise connected, there exists an arc J in Z such that $z \in J$ and $T \cap J = \emptyset$. Shortening T and J , if necessary, we may assume that the set $\mathcal{M} = \{\{a, b\} \in F_2(Z) : a \in T \text{ and } b \in J\}$ is contained in \mathcal{U} . Notice that \mathcal{M} is homeomorphic to $T \times J$ and thus \mathcal{M} is not embeddable in a 2-manifold (this follows from the Invariance of Domain Theorem [8, Theorem VI 9]). Hence, $A \in \mathcal{A}(Z)$.

To prove (b), take $A \in F_2(Z)$ such that $A \cap \text{cl}_Z(V(Z)) = \emptyset$. Let \mathcal{U} be a closed neighborhood of A in $F_2(Z)$ such that for each $B \in \mathcal{U}$, $B \cap \text{cl}_Z(V(Z)) = \emptyset$. Let \mathcal{M} be a locally connected subcontinuum of $F_2(Z)$ such that $\mathcal{M} \subset \mathcal{U}$. Let $M = \bigcup\{B : B \in \mathcal{M}\}$. By [5, Lemma 1] and [4, Lemma 2.2], M has at most two components and each one of them is a locally connected continuum. Then $M = M_1 \cup M_2$, where M_1 and M_2 are the components of M and it is possible that $M_1 = M_2$. Notice that no point of M is the vertex of a simple triod in Z . Thus, M does not contain simple triods. Hence, M_1 and M_2 are locally connected continua without simple triods. Therefore, each M_i is either an arc or a simple closed curve. In the case that $M_1 = M_2$, M is an arc or a simple closed curve, so ([11, Section 13]) $F_2(M)$ is a 2-cell or a Moebius strip. Since $\mathcal{M} \subset F_2(M)$, \mathcal{M} is embeddable in the Klein Bottle. In the case that $M_1 \neq M_2$, by [5, Lemma 1], each element $B \in \mathcal{M}$ intersects both sets M_1 and M_2 . Thus, \mathcal{M} is contained in the set $\mathcal{N} = \langle M_1, M_2 \rangle$ and \mathcal{N} is homeomorphic to $M_1 \times M_2$ which is homeomorphic to some of the following continua: (a) $[0, 1]^2$, (b) $S^1 \times [0, 1]$ or (c) $S^1 \times S^1$. In any case, \mathcal{N} is embeddable in $S^1 \times S^1$. Since $\mathcal{M} \subset \mathcal{N}$, we conclude that $A \notin \mathcal{A}(Z)$.

As a consequence of Claim 1, we obtain the following.

Claim 2. $\mathcal{A}(X) = \{A \in F_2(X) : v_0 \in A\}$.

Since the map $f : X \rightarrow F_2(X)$ given by $f(x) = \{v_0, x\}$ is an embedding and $f(X) = \mathcal{A}(X)$, we obtain that X , $\mathcal{A}(X)$ and $\mathcal{A}(Y)$ are homeomorphic.

Claim 3. $V(Y)$ is a one-point set.

We prove Claim 3. If $V(Y) = \emptyset$, by Claim 1(b), $\mathcal{A}(Y) = \emptyset$. This is a contradiction since $\mathcal{A}(Y)$ is homeomorphic to X . If $V(Y)$ contains two different elements v_1 and v_2 , by Claim 1(a), $\mathcal{R} = \{A \in F_2(Y) : v_1 \in A\}$ is contained in $\mathcal{A}(Y)$. Since the map $g : Y \rightarrow \mathcal{R}$ given by $g(y) = \{v_1, y\}$ is a homeomorphism and $V(Y)$ has two elements, we obtain that $V(\mathcal{R})$ has two elements. Thus, \mathcal{R} is a subcontinuum of $\mathcal{A}(Y)$ such that $|V(\mathcal{R})| \geq 2$. Since $\mathcal{A}(Y)$ is homeomorphic to X , we obtain that X contains a subcontinuum R such that $|V(R)| \geq 2$. Since $|V(X)| \geq |V(R)| \geq 2$, we obtain a contradiction. This completes the proof of Claim 3.

By Claim 3, $V(Y) = \{w\}$, for some $w \in Y$. By Claim 1, $\mathcal{A}(Y) = \{\{w, y\} : y \in Y\}$ which is homeomorphic to Y . Therefore, X is homeomorphic to Y . \square

Corollary 13. *If X belongs to one of the following families of continua, then X has unique hyperspace $F_2(X)$.*

- (a) fans,
- (b) cones over compact metric spaces containing no arcs,
- (c) cones over hereditarily indecomposable continua.

By Corollary 13 and [5, Theorem 5], each fan has unique hyperspace for all $n \neq 3$.

Question 14 ([10, Question 41]). *Let X be a fan, does X have unique hyperspace $F_3(X)$?*

Question 15. *Let X be an arcwise connected continuum for which there exists a unique point $v_0 \in X$ such that v_0 is the vertex of a simple triod. Does X have unique hyperspace $F_3(X)$?*

The most important question on the topic of this paper is the following.

Question 16 ([10, Question 43]). *Does there exist a finite-dimensional continuum X such that X does not have unique hyperspace $F_n(X)$ for some $n > 1$?*

Acknowledgment. The authors wish to thank the participants in the Sixth Workshop on Continua and Hyperspaces, celebrated in Puebla City during the Summer of 2012, for useful discussions. Particularly, V. Córdova Salazar, V. Galicia Aldama, L.A. Guerrero Méndez, D. Herrera Carrasco, M. de J. López Toriz, F. Macías Romero, D. Maya Escudero and F. Vázquez Juárez.

This paper was partially supported by the project “Hiperespacios topológicos (0128584)” of Consejo Nacional de Ciencia y Tecnología (CONACYT), 2009 and the project “Teoría de Continuos, Hiperespacios y Sistemas Dinámicos” (IN104613) of PAPIIT, DGAPA, UNAM.

REFERENCES

[1] Acosta G., Hernández-Gutiérrez R., Martínez-de-la-Vega V., *Dendrites and symmetric products*, Glas. Mat. Ser. III 44 (64) (2009), 195–210.

- [2] Castañeda E., *Embedding symmetric products in Euclidean spaces*, Continuum Theory, Lectures Notes in Pure and Applied Mathematics, 230, Marcel Dekker, New York, 2002, pp. 67–79.
- [3] Castañeda E., Illanes A., *Finite graphs have unique symmetric products*, Topology Appl. **153** (2006), 1434–1450.
- [4] Curtis D., Nhu N.T., *Hyperspaces of finite subsets which are homeomorphic to \aleph_0 -dimensional linear metric spaces*, Topology Appl. **19** (1985), 251–260.
- [5] Hernández-Gutiérrez R., Martínez-de-la-Vega V., *Rigidity of symmetric products*, preprint.
- [6] Herrera-Carrasco D., López M. de J., Macías-Romero F., *Dendrites with unique symmetric products*, Topology Proc. **34** (2009), 175–190.
- [7] Herrera-Carrasco D., Macías-Romero F., Vázquez-Juárez F., *Peano continua with unique symmetric products*, J. Math. Res. **4** (2012), 1–9.
- [8] Hurewicz W., Wallman H., *Dimension Theory*, Princeton University Press, Princeton, 1969.
- [9] Illanes A., *Dendrites with unique hyperspace $F_2(X)$* , JP J. Geom. Topol. **2** (2002), 75–96.
- [10] Illanes A., *Uniqueness of hyperspaces*, Questions Answers Gen. Topology **30** (2012), 21–44.
- [11] Illanes A., *Models of hyperspaces*, Topology Proc. **41** (2013), 39–64.
- [12] Illanes A., Martínez-Montejano J.M., *Compactifications of $[0, \infty)$ with unique hyperspace $F_n(X)$* , Glas. Mat. Ser. III 44 (64) (2009), 457–478.
- [13] Illanes A., Nadler S.B., Jr., *Hyperspaces; Fundamentals and Recent Advances*, Monographs and Textbooks in Pure and Applied Mathematics, 216, Marcel Dekker, New York, 1999.
- [14] Nadler S.B., Jr., *Continuum Theory. An Introduction*, Monographs and Textbooks in Pure and Applied Mathematics, 158, Marcel Dekker, New York, 1992.

José G. Anaya, Enrique Castañeda-Alvarado:

FACULTAD DE CIENCIAS, UNIVERSIDAD AUTÓNOMA DEL ESTADO DE MÉXICO, TOLUCA, MÉXICO, C.P. 50000

E-mail: jgao@uaemex.mx
eca@uaemex.mx

Alejandro Illanes:

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CIRCUITO EXTERIOR, CIUDAD UNIVERSITARIA, MÉXICO, 04510, D.F., MÉXICO

E-mail: illanes@matem.unam.mx

(Received January 9, 2013)