

Images of some functions and functional spaces under the Dunkl-Hermite semigroup

NÉJIB BEN SALEM, WALID NEFZI

Abstract. We propose the study of some questions related to the Dunkl-Hermite semigroup. Essentially, we characterize the images of the Dunkl-Hermite-Sobolev space, $\mathcal{S}(\mathbb{R})$ and $L^p_\alpha(\mathbb{R})$, $1 < p < \infty$, under the Dunkl-Hermite semigroup. Also, we consider the image of the space of tempered distributions and we give Paley-Wiener type theorems for the transforms given by the Dunkl-Hermite semigroup.

Keywords: Dunkl-Hermite functions; Dunkl-Hermite semigroup; Dunkl-Hermite-Sobolev space

Classification: 42B25, 46E35, 47B38, 47D03

1. Introduction and statement of the results

Let D_α , $\alpha \geq -\frac{1}{2}$, be the Dunkl operator on the real line defined by

$$D_\alpha f(x) = f'(x) + \frac{2\alpha + 1}{x} \left[\frac{f(x) - f(-x)}{2} \right], \quad f \in C^1(\mathbb{R}).$$

To this operator is associated the Dunkl-Hermite operator

$$\mathcal{H}_\alpha = -D_\alpha^2 + x^2.$$

Its spectral decomposition is given by the Dunkl-Hermite functions h_n^α defined by

$$h_n^\alpha(x) = e^{-\frac{x^2}{2}} H_n^\alpha(x), \quad n \in \mathbb{N},$$

namely we have (see [11])

$$\mathcal{H}_\alpha h_n^\alpha(x) = (2n + 2\alpha + 2)h_n^\alpha(x).$$

Here H_n^α is the Dunkl-Hermite polynomial given by

$$H_n^\alpha(x) = 2^{-\frac{n}{2}} \sqrt{\frac{b_n(\alpha)}{\Gamma(\alpha + 1)}} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^k}{k! b_{n-2k}(\alpha)} (2x)^{n-2k},$$

where $b_n(\alpha)$ is the generalized factorial defined by Rosenblum in [10],

$$b_n(\alpha) = \frac{2^n (\lfloor \frac{n}{2} \rfloor)!}{\Gamma(\alpha + 1)} \Gamma\left(\left[\frac{n+1}{2}\right] + \alpha + 1\right),$$

$[n/2]$ denotes the integral part of $n/2$. More precisely, these polynomials are expressed in terms of the Laguerre polynomials,

$$H_n^\alpha(x) = \frac{(-1)^{[\frac{n}{2}]}}{\sqrt{\Gamma(\alpha + 1)}} \frac{2^{\frac{n}{2}} ([\frac{n}{2}]!)!}{\sqrt{b_n(\alpha)}} x^{\theta_n} L_{[\frac{n}{2}]^{\alpha+\theta_n}}(x^2),$$

where θ_n is defined to be 0 if n is even and 1 if n is odd.

Hereafter, $L_\alpha^p(\mathbb{R}) = L^p(\mathbb{R}, |x|^{2\alpha+1} dx)$, $1 \leq p < +\infty$, denotes the space of measurable functions on \mathbb{R} satisfying

$$\|f\|_{\alpha,p} := \left(\int_{\mathbb{R}} |f(x)|^p |x|^{2\alpha+1} dx \right)^{\frac{1}{p}} < +\infty.$$

It is known that $\{h_n^\alpha, n \in \mathbb{N}\}$ forms an orthonormal basis of $L_\alpha^2(\mathbb{R})$. So for $f \in L_\alpha^2(\mathbb{R})$

$$\mathcal{H}_\alpha f = \sum_{n=0}^{\infty} (2n + 2\alpha + 2) a_n^\alpha(f) h_n^\alpha$$

with $a_n^\alpha(f) = \int_{\mathbb{R}} f(x) h_n^\alpha(x) |x|^{2\alpha+1} dx$.

Then, for a non-negative integer m , the Dunkl-Hermite-Sobolev space $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ is defined to be the image of $L_\alpha^2(\mathbb{R})$ under $(\mathcal{H}_\alpha)^{-m}$. We remark that $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ is a Hilbert space under the inner product

$$\langle f, g \rangle_{\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}} = \sum_{n=0}^{\infty} (2n + 2\alpha + 2)^{2m} a_n^\alpha(f) \overline{a_n^\alpha(g)}.$$

The Dunkl-Hermite semigroup denoted by $e^{-t\mathcal{H}_\alpha}$, $t > 0$, is defined by

$$e^{-t\mathcal{H}_\alpha} f = \sum_{n=0}^{\infty} e^{-(2n+2\alpha+2)t} a_n^\alpha(f) h_n^\alpha$$

for $f \in L_\alpha^2(\mathbb{R})$ and $f = \sum_{n=0}^{\infty} a_n^\alpha(f) h_n^\alpha$.

Using the Mehler formula for the Dunkl-Hermite polynomials H_n^α (see [10]), we can write $e^{-t\mathcal{H}_\alpha}$, on a dense subspace of $L_\alpha^2(\mathbb{R})$, as an integral operator with kernel $\mathcal{M}_t^\alpha(x, y)$

$$(1) \quad [e^{-t\mathcal{H}_\alpha} f](x) = \int_{\mathbb{R}} f(y) \mathcal{M}_t^\alpha(x, y) |y|^{2\alpha+1} dy.$$

The kernel $\mathcal{M}_t^\alpha(x, y)$ can be explicitly written as

$$\mathcal{M}_t^\alpha(x, y) = \frac{1}{\Gamma(\alpha + 1)(2 \sinh(2t))^{\alpha+1}} e^{-\frac{1}{2} \coth(2t)(x^2+y^2)} E_\alpha\left(\frac{x}{\sinh(2t)}, y\right),$$

where $E_\alpha(\xi, x)$ is the Dunkl kernel given by

$$E_\alpha(\xi, x) = j_\alpha(\xi x) + \frac{\xi x}{2(\alpha + 1)} j_{\alpha+1}(\xi x),$$

j_β being the spherical Bessel function of order β given by

$$j_\beta(t) = \Gamma(\beta + 1) \sum_{n=0}^\infty \frac{1}{n! \Gamma(n + \beta + 1)} \left(\frac{t}{2}\right)^{2n}.$$

We define the holomorphic Dunkl-Sobolev space $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$ as the image of $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ under $e^{-t\mathcal{H}_\alpha}$. It can be viewed as a Hilbert space simply by transferring the Hilbert space structure of $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$. In what follows, we give a characterization of the space $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$.

Using the reproducing kernel property, we show that if F is a holomorphic function on \mathbb{C} , then there exists a function $f \in \mathcal{S}(\mathbb{R})$ (the Schwartz space) such that $F = e^{-t\mathcal{H}_\alpha} f$ if and only if F satisfies

$$|F(z)|^2 \leq C_{t,\alpha,m} \frac{e^{-\tanh(2t)x^2 + \coth(2t)y^2}}{(1 + x^2 + y^2)^{2m}}, \quad z = x + iy,$$

for some constant $C_{t,\alpha,m}$ $m = 1, 2, 3, \dots$

The formula (1) permits to extend $e^{-t\mathcal{H}_\alpha}$ on the spaces $L_\alpha^p(\mathbb{R})$. We establish that if $f \in L_\alpha^p(\mathbb{R})$ for $1 < p < \infty$ then $e^{-t\mathcal{H}_\alpha}(f)$ is holomorphic and $e^{-t\mathcal{H}_\alpha}(f) \in L_\alpha^s(\mathbb{C}, V_{t,\frac{p}{2}}^{\frac{s+\epsilon}{2}})$ for every $\epsilon > 0$ and any $1 \leq s < \infty$, where

$$V_{t,\frac{p}{2}}^r(x + iy) = \exp\left(-2r\left(\frac{p}{(p-1)\sinh 4t}x^2 + \frac{\coth 2t}{2}y^2\right)\right).$$

Next, we consider the space of tempered distributions. For $S \in \mathcal{S}'(\mathbb{R})$, we show that $e^{-t\mathcal{H}_\alpha}$ is given by a function defined by

$$e^{-t\mathcal{H}_\alpha} S(x) = e^{-\frac{1}{2}\left(\frac{\cosh 2t-1}{\sinh 2t}\right)x^2} \left(e^{-\frac{1}{2}\left(\frac{\cosh 2t-1}{\sinh 2t}\right)y^2} S *_{\alpha} q_{\frac{\sinh 2t}{2}}\right)(x),$$

where $q_t, t > 0$, denotes the heat kernel associated with the Dunkl operator D_α , given by

$$q_t(x) = \frac{1}{\Gamma(\alpha + 1)} (4t)^{-(\alpha+1)} e^{-\frac{x^2}{4t}},$$

and $*_\alpha$ is the generalized convolution product associated with the Dunkl operator D_α (see [13]). Moreover, $e^{-t\mathcal{H}_\alpha} S$ is a \mathcal{C}^∞ function on \mathbb{R} .

These results permit us to characterize the image of tempered distributions on \mathbb{R} under the Dunkl-Hermite semigroup. We establish that if F is a holomorphic function on \mathbb{C} , then there exists a distribution $f \in \mathcal{S}'(\mathbb{R})$ with $F = e^{-t\mathcal{H}_\alpha} f$ if and

only if F satisfies

$$|F(z)|^2 \leq C_{t,\alpha}(1 + |z|^2)^{2m} \exp(-\tanh(2t)x^2 + \coth(2t)y^2),$$

for some non-negative integer m .

Next, we define the transform \mathcal{T}_a^α , for $a > 0$, by

$$\mathcal{T}_a^\alpha(S)(x) = \langle S, e^{-\frac{1}{2}a(\cdot)^2} E_\alpha(-ix, \cdot) \rangle, \quad S \in \mathcal{S}'(\mathbb{R}).$$

We prove that this transform is related to the Dunkl-Hermite semigroup and we establish a Paley-Wiener theorem for $\mathcal{T}_a^\alpha f$. For any $a > 0$ the transform \mathcal{T}_a^α of a tempered distribution f on \mathbb{R} extends to \mathbb{C} as an entire function which satisfies the estimate

$$|\mathcal{T}_a^\alpha f(z)| \leq C_\alpha(1 + x^2 + y^2)^m e^{\frac{1}{2}a^{-1}y^2}$$

for some non-negative integer m . Conversely, if an entire function F satisfies such an estimate, then $F = \mathcal{T}_a^\alpha f$ for some tempered distribution f .

Again relating the Dunkl-Hermite semigroup and the Dunkl transform, we obtain a characterization of the image of compactly supported distributions under the Dunkl-Hermite semigroup. If f is a distribution supported in a ball of radius R centered at the origin then for any $t > 0$ the function $e^{-t\mathcal{H}_\alpha} f$ extends to \mathbb{C} as an entire function which satisfies

$$|e^{-t\mathcal{H}_\alpha} f(z)| \leq C e^{-\frac{1}{2} \coth 2t(x^2 - y^2)} e^{\frac{R|x|}{\sinh 2t}}.$$

Conversely, any entire function F satisfying the above condition is of the form $e^{-t\mathcal{H}_\alpha} f$, where f is supported inside a ball of radius R centered at the origin.

We point out that the results of this paper extend naturally those established in [8] by R. Radha and S. Thangavelu.

We conclude this introduction by giving the organization of this paper. In the next section, we define the Dunkl-Hermite-Sobolev space and we characterize its images under the Dunkl-Hermite semigroup. The third section deals with a characterization of the image of $\mathcal{S}(\mathbb{R})$ and $L_\alpha^p(\mathbb{R})$ under the Dunkl-Hermite semigroup. In the last section we establish Paley-Wiener type theorems for the tempered distributions and the compactly supported distributions under the Dunkl-Hermite semigroup.

2. Holomorphic Dunkl-Sobolev spaces

We have established in [1] that every element in the range of the operator $e^{-t\mathcal{H}_\alpha}$ defined on L_α^2 can be analytically extended to the complex plane \mathbb{C} , hence we shall consider the operator $e^{-t\mathcal{H}_\alpha}$ as a linear operator from L_α^2 into an entire function space and the entire extension will be simply denoted by $e^{-t\mathcal{H}_\alpha} f(z)$, $z = x + iy$.

In this section, we introduce the Dunkl-Hermite-Sobolev space and we give a characterization of its images under the Dunkl-Hermite semigroup.

Notation 1. Let $U_{t,e}^\alpha(z) = \frac{2}{\pi \sinh(4t)} K_\alpha\left(\frac{|z|^2}{\sinh(4t)}\right) \exp\{\coth(4t)(x^2 - y^2)\} |z|^{2\alpha+2}$ and $U_{t,o}^\alpha(z) = \frac{2}{\pi \sinh(4t)} K_{\alpha+1}\left(\frac{|z|^2}{\sinh(4t)}\right) \exp\{\coth(4t)(x^2 - y^2)\} |z|^{2\alpha+2}$. We have

$$U_{t,o}^\alpha(z) = \frac{U_{t,e}^{\alpha+1}(z)}{|z|^2}.$$

Here K_ν is the Macdonald function defined in [4] by:

$$K_\nu(z) = \frac{\pi}{2} \frac{I_{-\nu}(z) - I_\nu(z)}{\sin(\nu\pi)}, \quad \nu \in \mathbb{C} \setminus \mathbb{Z}, \quad |\arg(z)| < \pi$$

where

$$I_\nu(z) = \frac{1}{\Gamma(\nu + 1)} \left(\frac{z}{2}\right)^\nu j_\nu(z)$$

and for an integer n ,

$$K_n(z) = \lim_{\nu \rightarrow n} K_\nu(z).$$

Let $\mathcal{H}_{t,e}^\alpha(\mathbb{C})$ denote the Hilbert space of all even entire functions on \mathbb{C} which are square integrable with respect to the weight function $U_{t,e}^\alpha$, equipped with the inner product defined by

$$\langle f, g \rangle_{\alpha,e} = \int_{\mathbb{C}} f(z) \overline{g(z)} U_{t,e}^\alpha(z) dz.$$

Let $\mathcal{H}_{t,o}^\alpha(\mathbb{C})$ denote the Hilbert space of all odd entire functions on \mathbb{C} which are square integrable with respect to the weight function $U_{t,o}^\alpha$, equipped with the inner product defined by

$$\langle f, g \rangle_{\alpha,o} = \int_{\mathbb{C}} f(z) \overline{g(z)} U_{t,o}^\alpha(z) dz.$$

Let \mathcal{H}_t^α denote the direct sum of $\mathcal{H}_{t,e}^\alpha$ and $\mathcal{H}_{t,o}^\alpha$ admitting the inner product

$$\langle f, g \rangle_{\alpha,t} = \langle f_e, g_e \rangle_{\alpha,e} + \langle f_o, g_o \rangle_{\alpha,o},$$

where $f_e(z) = \frac{f(z)+f(-z)}{2}$ and $f_o(z) = \frac{f(z)-f(-z)}{2}$.

We recall the following results proved in [1].

Theorem 1. *The image of $L_\alpha^2(\mathbb{R})$ under the Dunkl-Hermite semigroup is the Fock type space \mathcal{H}_t^α . The Dunkl-Hermite semigroup $e^{-t\mathcal{H}_\alpha}$ is an isometric isomorphism from $L_\alpha^2(\mathbb{R})$ into $\mathcal{H}_t^\alpha(\mathbb{C})$.*

Also we have the orthogonality property

$$\begin{aligned} (2) \quad \langle h_n^\alpha, h_m^\alpha \rangle_{\alpha,t} &= \int_{\mathbb{C}} h_{n,e}^\alpha(z) \overline{h_{m,e}^\alpha(z)} U_{t,e}^\alpha(z) dz + \int_{\mathbb{C}} h_{n,o}^\alpha(z) \overline{h_{m,o}^\alpha(z)} U_{t,o}^\alpha(z) dz \\ &= e^{2(2n+2\alpha+2)t} \delta_{n,m}, \end{aligned}$$

where $h_n^\alpha(z)$ is the extension of the Dunkl-Hermite function $h_n^\alpha(x)$ to \mathbb{C} as an entire function.

Let $\widetilde{h}_n^\alpha(z) = e^{-(2n+2\alpha+2)t}h_n^\alpha(z)$, then $\{\widetilde{h}_n^\alpha, n \in \mathbb{N}\}$ forms an orthonormal basis for $\mathcal{H}_t^\alpha(\mathbb{C})$. Thus any $F \in \mathcal{H}_t^\alpha(\mathbb{C})$ can be written as

$$F = \sum_{n=0}^{\infty} \langle F, \widetilde{h}_n^\alpha \rangle_{\alpha,t} \widetilde{h}_n^\alpha.$$

Definition 1. Let m be a non-negative integer. The Dunkl-Hermite-Sobolev space $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ is defined to be the image of $L_\alpha^2(\mathbb{R})$ under $(\mathcal{H}_\alpha)^{-m}$.

Remark 1. We remark that $f \in \mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ if and only if $\sum_{n=0}^{\infty} (2n + 2\alpha + 2)^{2m} |a_n^\alpha(f)|^2 < \infty$. The Sobolev space $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ is an Hilbert space under the inner product

$$\langle f, g \rangle_{\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}} = \sum_{n=0}^{\infty} (2n + 2\alpha + 2)^{2m} a_n^\alpha(f) \overline{a_n^\alpha(g)}.$$

As $(\mathcal{H}_\alpha)^m f = \sum_{n=0}^{\infty} (2n + 2\alpha + 2)^m a_n^\alpha(f) h_n^\alpha$ then

$$\langle f, g \rangle_{\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}} = \langle (\mathcal{H}_\alpha)^m f, (\mathcal{H}_\alpha)^m g \rangle_{L_\alpha^2}.$$

Definition 2. We define the holomorphic Dunkl-Sobolev space $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$ to be the image of $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ under $e^{-t\mathcal{H}_\alpha}$.

Remark 2. It is clear that by transferring the Hilbert space structure of $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ to $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$, the space $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$ becomes a Hilbert space. The Dunkl-Hermite semigroup $e^{-t\mathcal{H}_\alpha}$ is an isometric isomorphism from $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ onto $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$. Then we can write

$$\langle F, G \rangle_{\mathcal{W}_{t,\alpha}^{m,2}} = \sum_{n=0}^{\infty} (2n + 2\alpha + 2)^{2m} a_n^\alpha(f) \overline{a_n^\alpha(g)}$$

whenever $F = e^{-t\mathcal{H}_\alpha} f$ and $G = e^{-t\mathcal{H}_\alpha} g$.

Notation 2. We denote by $\mathcal{O}(\mathbb{C})$ the set of all holomorphic functions on \mathbb{C} . Let $\mathcal{F}_{t,e}^{m,\alpha}(\mathbb{C})$ be the space of all even functions in $\mathcal{O}(\mathbb{C})$ which are square integrable with respect to the measure $|\frac{d^{2m}}{dt^{2m}} U_{t,e}^\alpha(z)| dz$. We equip $\mathcal{F}_{t,e}^{m,\alpha}(\mathbb{C})$ with the sesquilinear form

$$\langle F, G \rangle_{m,e} = \int_{\mathbb{C}} F(z) \overline{G(z)} \frac{d^{2m}}{dt^{2m}} U_{t,e}^\alpha(z) dz.$$

Let $\mathcal{F}_{t,o}^{m,\alpha}(\mathbb{C})$ be the space of all odd functions in $\mathcal{O}(\mathbb{C})$ which are square integrable with respect to the measure $|\frac{d^{2m}}{dt^{2m}} U_{t,o}^\alpha(z)| dz$. We equip $\mathcal{F}_{t,o}^{m,\alpha}(\mathbb{C})$ with the

sesquilinear form

$$\langle F, G \rangle_{m,o} = \int_{\mathbb{C}} F(z) \overline{G(z)} \frac{d^{2m}}{dt^{2m}} U_{t,o}^\alpha(z) dz.$$

Let $\mathcal{F}_t^{m,\alpha}(\mathbb{C})$ be the direct sum of $\mathcal{F}_{t,e}^{m,\alpha}(\mathbb{C})$ and $\mathcal{F}_{t,o}^{m,\alpha}(\mathbb{C})$ admitting the sesquilinear form

$$\langle F, G \rangle_{m,\alpha} = \langle F_e, G_e \rangle_{m,e} + \langle F_o, G_o \rangle_{m,o}.$$

We shall show below that this defines a pre-Hilbert space structure on $\mathcal{F}_t^{m,\alpha}(\mathbb{C}) \cap \mathcal{H}_t^\alpha(\mathbb{C})$.

Let $\mathcal{B}_t^{m,\alpha}(\mathbb{C})$ denote the completion of $\mathcal{F}_t^{m,\alpha}(\mathbb{C}) \cap \mathcal{H}_t^\alpha(\mathbb{C})$ with respect to the norm induced by the above inner product. In the following proposition, we also show that $\|F\|_{m,\alpha}$ and $\|F\|_{\mathcal{W}_{t,\alpha}^{m,2}}$ coincide up to a constant multiple.

Proposition 1. *The sesquilinear form $\langle F, G \rangle_{m,\alpha}$, for a non-negative integer m , is an inner product on $\mathcal{F}_t^{m,\alpha}(\mathbb{C}) \cap \mathcal{H}_t^\alpha(\mathbb{C})$ and hence induces a norm $\|F\|_{m,\alpha}^2 = \langle F, F \rangle_{m,\alpha}$. We also have*

$$\|F\|_{m,\alpha}^2 = 2^{2m} \|F\|_{\mathcal{W}_{t,\alpha}^{m,2}}^2$$

for all functions $F = e^{-t\mathcal{H}_\alpha} f$ with $f \in \mathcal{S}(\mathbb{R})$.

PROOF: Let F be in $\mathcal{F}_t^{m,\alpha}(\mathbb{C}) \cap \mathcal{H}_t^\alpha(\mathbb{C})$. We expand the restriction of F to \mathbb{R} into an orthogonal expansion in terms of h_n^α (see [1]), and we can write

$$F(x + iy) = \sum_n \langle F, h_n^\alpha \rangle_{2,\alpha} h_n^\alpha(x + iy),$$

so we have that

$$\begin{aligned} I_t^\alpha &:= \int_{\mathbb{C}} |F_e(x + iy)|^2 U_{t,e}^\alpha(z) dz + \int_{\mathbb{C}} |F_o(x + iy)|^2 U_{t,o}^\alpha(z) dz \\ &= \left\langle \sum_n \langle F, h_n^\alpha \rangle_{2,\alpha} h_n^\alpha, \sum_q \langle F, h_q^\alpha \rangle_{2,\alpha} h_q^\alpha \right\rangle_{\alpha,t}. \end{aligned}$$

Using the orthogonality relation (2), we can show that

$$I_t^\alpha = \sum_n |\langle F, h_n^\alpha \rangle_{2,\alpha}|^2 e^{2(2n+2\alpha+2)t}.$$

By definition, for a nonnegative integer m , we have

$$\begin{aligned} \langle F, F \rangle_{m,\alpha} &= \int_{\mathbb{C}} |F_e(z)|^2 \frac{d^{2m}}{dt^{2m}} U_{t,e}^\alpha(z) dz + \int_{\mathbb{C}} |F_o(z)|^2 \frac{d^{2m}}{dt^{2m}} U_{t,o}^\alpha(z) dz \\ &= \frac{d^{2m}}{dt^{2m}} I_t^\alpha \\ &= 2^{2m} \sum_n (2n + 2\alpha + 2)^{2m} |\langle F, h_n^\alpha \rangle_{2,\alpha}|^2 e^{2(2n+2\alpha+2)t}. \end{aligned}$$

Thus it follows that the sesquilinear form defined above is positive definite and induces the norm $\|F\|_{m,\alpha}$.

On the other hand, we have the expansion

$$F(z) = \sum_{m=0}^\infty \langle F, \widetilde{h}_m^\alpha \rangle_{\alpha,t} \widetilde{h}_m^\alpha(z)$$

and

$$F = e^{-t\mathcal{H}_\alpha} f \quad \text{with } f \in L_\alpha^2(\mathbb{R}).$$

Thus we have

$$\begin{aligned} \langle F, h_n^\alpha \rangle_{2,\alpha} &= \int_{\mathbb{R}} \sum_{m=0}^\infty \langle F, \widetilde{h}_m^\alpha \rangle_{\alpha,t} \widetilde{h}_m^\alpha(x) h_n^\alpha(x) |x|^{2\alpha+1} dx \\ &= \int_{\mathbb{R}} \sum_{m=0}^\infty \langle f, h_m^\alpha \rangle_{2,\alpha} e^{-(2m+2\alpha+2)t} h_m^\alpha(x) h_n^\alpha(x) |x|^{2\alpha+1} dx \\ &= \sum_{m=0}^\infty \langle f, h_m^\alpha \rangle_{2,\alpha} e^{-(2m+2\alpha+2)t} \int_{\mathbb{R}} h_m^\alpha(x) h_n^\alpha(x) |x|^{2\alpha+1} dx \\ &= \langle f, h_n^\alpha \rangle_{2,\alpha} e^{-(2n+2\alpha+2)t}. \end{aligned}$$

Interchanging the order of summation and integration is justified by Lebesgue's dominated convergence theorem and limiting behavior of $\|h_n^\alpha\|_{\alpha,p}$ given in [2]. Again using the orthogonality relation (2), we get

$$\begin{aligned} \|F\|_{m,\alpha}^2 &= 2^{2m} \sum_n (2n + 2\alpha + 2)^{2m} |\langle F, h_n^\alpha \rangle_{2,\alpha}|^2 e^{2(2n+2\alpha+2)t} \\ &= 2^{2m} \sum_n (2n + 2\alpha + 2)^{2m} |\langle f, h_n^\alpha \rangle_{2,\alpha}|^2 \\ &= 2^{2m} \sum_n (2n + 2\alpha + 2)^{2m} |\langle F, \widetilde{h}_n^\alpha \rangle_{\alpha,t}|^2 \\ &= 2^{2m} \|F\|_{\mathcal{W}_{t,\alpha}^{m,2}}^2. \end{aligned}$$

□

Using this proposition we can easily prove the following result on the range of the Dunkl-Hermite-Sobolev spaces under the Dunkl-Hermite semigroup.

Theorem 2. *For every nonnegative integer m , $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$ coincides with $\mathcal{B}_t^{m,\alpha}(\mathbb{C})$ and the Dunkl-Hermite semigroup $e^{-t\mathcal{H}_\alpha}$ is an isometric isomorphism from $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ onto $\mathcal{B}_t^{m,\alpha}(\mathbb{C})$ up to a constant multiple.*

PROOF: Let $F \in \mathcal{F}_t^{m,\alpha}(\mathbb{C}) \cap \mathcal{H}_t^\alpha(\mathbb{C})$, hence F is of the form $e^{-t\mathcal{H}_\alpha} f$ with $f \in L_\alpha^2(\mathbb{R})$. Further, it follows from the above proposition, as the norms $\|F\|_{m,\alpha}$ and $\|F\|_{\mathcal{W}_{t,\alpha}^{m,2}}$ coincide, that $f \in \mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$. Consequently, $\mathcal{F}_t^{m,\alpha}(\mathbb{C}) \cap \mathcal{H}_t^\alpha(\mathbb{C})$ is contained in $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$. We have $\widetilde{h}_n^\alpha = e^{-t\mathcal{H}_\alpha} h_n^\alpha$, and

$$\begin{aligned} \|\widetilde{h}_n^\alpha\|_{m,\alpha}^2 &= 2^{2m} \|\widetilde{h}_n^\alpha\|_{\mathcal{W}_{t,\alpha}^{m,2}}^2 \\ &= 2^{2m} \|h_n^\alpha\|_{\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}}^2 \\ &= 2^{2m} (2n + 2\alpha + 2)^{2m} < \infty. \end{aligned}$$

So for all $n \in \mathbb{N}$, $\widetilde{h}_n^\alpha \in \mathcal{B}_t^{m,\alpha}(\mathbb{C})$. We have

$$\begin{aligned} \langle F, \widetilde{h}_n^\alpha \rangle_{\mathcal{W}_{t,\alpha}^{m,2}} &= \sum_{p=0}^{\infty} (2p + 2\alpha + 2)^{2m} \langle F, \widetilde{h}_p^\alpha \rangle_{\alpha,t} \langle \widetilde{h}_n^\alpha, \widetilde{h}_p^\alpha \rangle_{\alpha,t} \\ &= (2n + 2\alpha + 2)^{2m} \langle F, \widetilde{h}_n^\alpha \rangle_{\alpha,t}. \end{aligned}$$

Then it can be easily seen that if $\langle F, \widetilde{h}_n^\alpha \rangle_{\mathcal{W}_{t,\alpha}^{m,2}} = 0$ then $\langle F, \widetilde{h}_n^\alpha \rangle_{\alpha,t} = 0$. This gives that $F = 0$ because $\{\widetilde{h}_n^\alpha, n \in \mathbb{N}\}$ form an orthonormal basis for $\mathcal{H}_t^\alpha(\mathbb{C})$, so we have

$$\{\widetilde{h}_n^\alpha, n \in \mathbb{N}\} \subset \mathcal{B}_t^{m,\alpha}(\mathbb{C}) \subset \mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$$

and

$$\overline{\{\widetilde{h}_n^\alpha, n \in \mathbb{N}\}^{\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})}} = \mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C}).$$

Hence $\mathcal{F}_t^{m,\alpha}(\mathbb{C}) \cap \mathcal{H}_t^\alpha(\mathbb{C})$ is dense in $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$. □

3. The image of $\mathcal{S}(\mathbb{R})$ and $L_\alpha^p(\mathbb{R})$ under the Dunkl-Hermite semigroup

3.1 The image of $\mathcal{S}(\mathbb{R})$ under the Dunkl-Hermite semigroup. We begin by establishing that $\mathcal{S}(\mathbb{R})$ is stable under the Dunkl-Hermite semigroup.

First we recall that the heat kernel q_t , $t > 0$, associated with the Dunkl operators, see [12], is given by

$$q_t(x) = \frac{1}{\Gamma(\alpha + 1)} (4t)^{-(\alpha+1)} e^{-\frac{x^2}{4t}}.$$

This function belongs to $\mathcal{S}(\mathbb{R})$ and satisfies the following property

$$\tau_{-y}^\alpha q_t(x) = \frac{1}{\Gamma(\alpha + 1)} (4t)^{-(\alpha+1)} e^{-\frac{(x^2+y^2)}{4t}} E_\alpha\left(\frac{x}{2t}, y\right),$$

where τ_y^α is the generalized translation associated with the Dunkl operator D_α (see [13]).

Using the Mehler formula for the Dunkl-Hermite polynomials H_n^α (see [10]), we can write $e^{-t\mathcal{H}_\alpha}$ on $\mathcal{S}(\mathbb{R})$ as an integral operator with kernel $\mathcal{M}_t^\alpha(x, y)$

$$[e^{-t\mathcal{H}_\alpha} f](x) = \int_{\mathbb{R}} f(y) \mathcal{M}_t^\alpha(x, y) |y|^{2\alpha+1} dy.$$

The kernel $\mathcal{M}_t^\alpha(x, y)$ can be explicitly written as

$$\mathcal{M}_t^\alpha(x, y) = \frac{1}{\Gamma(\alpha + 1)(2 \sinh(2t))^{\alpha+1}} e^{-\frac{1}{2} \coth(2t)(x^2+y^2)} E_\alpha\left(\frac{x}{\sinh(2t)}, y\right),$$

where $E_\alpha(\xi, x)$ is the Dunkl kernel. We can see that the kernel $\mathcal{M}_t^\alpha(x, y)$ satisfies the following relation

$$\mathcal{M}_t^\alpha(x, y) = e^{-\frac{1}{2}(\frac{\cosh 2t - 1}{\sinh 2t})(x^2+y^2)} \tau_{-y}^\alpha q_{\frac{\sinh 2t}{2}}(x).$$

So for $\varphi \in \mathcal{S}(\mathbb{R})$, we have

$$e^{-t\mathcal{H}_\alpha} \varphi(y) = e^{-\frac{1}{2}(\frac{\cosh 2t - 1}{\sinh 2t})y^2} \left(e^{-\frac{1}{2}(\frac{\cosh 2t - 1}{\sinh 2t})x^2} \varphi *_\alpha q_{\frac{\sinh 2t}{2}} \right)(y),$$

where $*_\alpha$ is the generalized convolution product associated with the Dunkl operator D_α (see [13]).

As a consequence we have the following result.

Proposition 2. *The Dunkl-Hermite semigroup $e^{-t\mathcal{H}_\alpha}$ is a continuous transform from $\mathcal{S}(\mathbb{R})$ into $\mathcal{S}(\mathbb{R})$.*

In the following, we shall give a characterization of the image of the Schwartz space under the Dunkl-Hermite semigroup.

Let $F \in \mathcal{H}_t^\alpha(\mathbb{C})$ and for $z \in \mathbb{C}$, $F(z)$ be its entire extension. Since $F \rightarrow F(z)$ is a continuous linear functional on $\mathcal{H}_t^\alpha(\mathbb{C})$ for each $z \in \mathbb{C}$, Riesz representation theorem ensures that there exists a unique $\mathcal{N}_t^\alpha(z, \cdot) \in \mathcal{H}_t^\alpha(\mathbb{C})$ such that

$$F(z) = \langle F, \mathcal{N}_t^\alpha(z, \cdot) \rangle_{\alpha, t} = \langle F_e, \mathcal{N}_{t,e}^\alpha(z, \cdot) \rangle_{\alpha, e} + \langle F_o, \mathcal{N}_{t,o}^\alpha(z, \cdot) \rangle_{\alpha, o}.$$

The function $\mathcal{N}_t^\alpha(z, w)$ is called the reproducing kernel for $\mathcal{H}_t^\alpha(\mathbb{C})$. By expanding F in terms of \widetilde{h}_n^α , we can write

$$F(z) = \sum_{n=0}^{\infty} \langle F, \widetilde{h}_n^\alpha \rangle_{\alpha, t} \widetilde{h}_n^\alpha(z) = \langle F, \sum_{n=0}^{\infty} \widetilde{h}_n^\alpha(\cdot) \overline{\widetilde{h}_n^\alpha(z)} \rangle_{\alpha, t}.$$

So, we deduce that

$$\mathcal{N}_t^\alpha(z, w) = \sum_n e^{-(2n+2\alpha+2)2t} h_n^\alpha(w) \overline{h_n^\alpha(\bar{z})}.$$

Cauchy-Schwartz inequality gives us

$$|F(z)|^2 = |\langle F, \mathcal{N}_t^\alpha(z, \cdot) \rangle_{\alpha,t}|^2 \leq \|F\|_{\alpha,t}^2 \|\mathcal{N}_t^\alpha(z, \cdot)\|_{\alpha,t}^2 = \|F\|_{\alpha,t}^2 \mathcal{N}_t^\alpha(z, z).$$

Using Mehler’s formula, we can explicitly calculate $\mathcal{N}_t^\alpha(z, z)$, in fact, we get

$$\begin{aligned} \mathcal{N}_t^\alpha(z, z) &= \sum_n e^{-(2n+2\alpha+2)2t} h_n^\alpha(z) \overline{h_n^\alpha(\bar{z})} = e^{-(2\alpha+2)2t} \sum_n (e^{-4t})^n h_n^\alpha(z) \overline{h_n^\alpha(\bar{z})} \\ &= \frac{1}{2^{\alpha+1} \Gamma(\alpha+1)} (\sinh(4t))^{-(\alpha+1)} \exp\left(-\frac{1}{2} \coth(4t)(z^2 + \bar{z}^2)\right) E_\alpha\left(\frac{1}{\sinh(4t)}, z\bar{z}\right). \end{aligned}$$

If $z = x + iy$ we have that

$$\begin{aligned} |F(z)|^2 &\leq \frac{1}{2^{\alpha+1} \Gamma(\alpha+1)} (\sinh(4t))^{-(\alpha+1)} \exp(-\coth(4t)(x^2 - y^2)) \\ &\quad \times E_\alpha\left(\frac{1}{\sinh(4t)}, x^2 + y^2\right) \|F\|_{\alpha,t}^2. \end{aligned}$$

It is known that the kernel E_α satisfies the inequality below for all $x, y \in \mathbb{R}$ (see [3])

$$(3) \quad E_\alpha\left(\frac{1}{\sinh(4t)}, x^2 + y^2\right) \leq \exp\left(\frac{1}{\sinh(4t)}(x^2 + y^2)\right).$$

As

$$-\coth(4t)(x^2 - y^2) + \frac{1}{\sinh(4t)}(x^2 + y^2) = -\tanh(2t)x^2 + \coth(2t)y^2,$$

we deduce

$$|F(z)|^2 \leq \frac{1}{2^{\alpha+1} \Gamma(\alpha+1)} (\sinh(4t))^{-(\alpha+1)} \exp(-\tanh(2t)x^2 + \coth(2t)y^2) \|F\|_{\alpha,t}^2,$$

which gives a pointwise estimate for functions $F \in \mathcal{H}_t^\alpha(\mathbb{C})$.

Notation 3. We denote by $\mathcal{N}_t^{\alpha,2m}(z, w)$ the kernel defined by

$$\mathcal{N}_t^{\alpha,2m}(z, w) = \sum_n (2n + 2\alpha + 2)^{-2m} \widetilde{h}_n^\alpha(\bar{z}) \widetilde{h}_n^\alpha(w).$$

In order to obtain pointwise estimates for $F \in \mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$, we have to show the following result.

Proposition 3. $\mathcal{N}_t^{\alpha,2m}(z, w)$ is a reproducing kernel for $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$.

PROOF: For $z \in \mathbb{C}$, the function $w \rightarrow \mathcal{N}_t^{\alpha,2m}(z, w)$ belongs to $\mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$ because $\widetilde{h}_n^\alpha(w) \in \mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$ for all $w \in \mathbb{C}$. We show now the reproducing property. For $z \in \mathbb{C}$ and $F \in \mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$, we have

$$\begin{aligned} \langle F, \mathcal{N}_t^{\alpha,2m}(z, \cdot) \rangle_{\mathcal{W}_{t,\alpha}^{m,2}} &= \sum_{n=0}^{\infty} (2n + 2\alpha + 2)^{2m} \langle F, \widetilde{h}_n^\alpha \rangle_{\alpha,t} \overline{\langle \mathcal{N}_t^{\alpha,2m}(z, \cdot), \widetilde{h}_n^\alpha \rangle_{\alpha,t}} \\ &= \sum_{n=0}^{\infty} (2n + 2\alpha + 2)^{2m} \langle F, \widetilde{h}_n^\alpha \rangle_{\alpha,t} (2n + 2\alpha + 2)^{-2m} \widetilde{h}_n^\alpha(z) \\ &= \sum_{n=0}^{\infty} \langle F, \widetilde{h}_n^\alpha \rangle_{\alpha,t} \widetilde{h}_n^\alpha(z) = F(z). \end{aligned} \quad \square$$

The last kernel can be written as

$$\mathcal{N}_t^{\alpha,2m}(z, w) = \frac{2^{2m}}{(2m - 1)!} \int_0^{+\infty} s^{2m-1} \mathcal{N}_{s+t}^\alpha(z, w) ds.$$

Using the explicit formula for $\mathcal{N}_s^\alpha(z, z)$, we have

$$\begin{aligned} \mathcal{N}_t^{\alpha,2m}(z, z) &= \frac{2^{2m}}{(2m - 1)! 2^{\alpha+1} \Gamma(\alpha + 1)} \int_0^{+\infty} s^{2m-1} (\sinh 4(t + s))^{-(\alpha+1)} \\ &\quad \times \exp(-\coth 4(t + s)(x^2 - y^2)) \times E_\alpha\left(\frac{1}{\sinh 4(t + s)}, x^2 + y^2\right) ds. \end{aligned}$$

Theorem 3 (Dunkl-Sobolev-embedding theorem). *Let m be a nonnegative integer. Then every $F \in \mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$ satisfies the estimate*

$$|F(z)|^2 \leq C_{t,\alpha} (1 + x^2 + y^2)^{-2m} \exp(-\tanh(2t)x^2 + \coth(2t)y^2),$$

where $C_{t,\alpha}$ is a constant depending on t and α .

PROOF: We begin by estimating the integral appearing in the representation of the reproducing kernel $\mathcal{N}_t^{\alpha,2m}(z, z)$, using the inequality (3) we obtain

$$\begin{aligned} \mathcal{N}_t^{\alpha,2m}(z, z) &\leq \frac{2^{2m}}{(2m - 1)! 2^{\alpha+1} \Gamma(\alpha + 1)} \int_0^{+\infty} s^{2m-1} (\sinh 4(t + s))^{-(\alpha+1)} \\ &\quad \times e^{-\tanh 2(t+s)x^2 + \coth 2(t+s)y^2} ds. \end{aligned}$$

We rewrite this in the following form

$$\mathcal{N}_t^{\alpha,2m}(z, z) \leq \frac{2^{2m}}{(2m - 1)! 2^{\alpha+1} \Gamma(\alpha + 1)} e^{-\tanh(2t)x^2 + \coth(2t)y^2} J_t^\alpha,$$

where

$$J_t^\alpha = \int_0^{+\infty} s^{2m-1} (\sinh 4(t+s))^{-(\alpha+1)} \times e^{-x^2(\tanh 2(t+s)-\tanh(2t))} \times e^{y^2(\coth 2(t+s)-\coth(2t))} ds,$$

which after some simplification yields

$$J_t^\alpha = \int_0^{+\infty} s^{2m-1} (\sinh 4(t+s))^{-(\alpha+1)} \times \exp\left(-x^2\left(\frac{\sinh 2s}{\cosh 2(t+s)\cosh 2t}\right) - y^2\left(\frac{\sinh 2s}{\sinh 2(t+s)\sinh 2t}\right)\right) ds.$$

Thus we only need to show that the above integral is bounded by $C_{t,\alpha}(1+x^2+y^2)^{-2m}$.

To prove this estimate we break up the above integral into two parts. Using the elementary properties of the functions \sinh and \cosh , we see that

$$\int_0^t s^{2m-1} (\sinh 4(t+s))^{-(\alpha+1)} \times \exp\left(-x^2\left(\frac{\sinh 2s}{\cosh 2(t+s)\cosh 2t}\right) - y^2\left(\frac{\sinh 2s}{\sinh 2(t+s)\sinh 2t}\right)\right) ds$$

is bounded by

$$\begin{aligned} & \int_0^{+\infty} s^{2m-1} e^{-4(\alpha+1)s} \exp\left(-2\left(\frac{x^2}{\cosh^2 4t} + \frac{y^2}{\sinh^2 4t}\right)s\right) ds \\ &= (2m-1)! [2(2(\alpha+1) + \frac{x^2}{\cosh^2 4t} + \frac{y^2}{\sinh^2 4t})]^{-2m} \\ &\leq C_{t,\alpha,m}(1+x^2+y^2)^{-2m}. \end{aligned}$$

On the other hand the integral

$$\int_t^\infty s^{2m-1} (\sinh 4(t+s))^{-(\alpha+1)} \times \exp\left(-x^2\left(\frac{\sinh 2s}{\cosh 2(t+s)\cosh 2t}\right) - y^2\left(\frac{\sinh 2s}{\sinh 2(t+s)\sinh 2t}\right)\right) ds,$$

is bounded by

$$\frac{(2m-1)!}{(4(\alpha+1))^{2m}} \exp\left(-\left(\frac{\tanh 2t}{\cosh 4t}x^2 + \frac{1}{\sinh 4t}y^2\right)\right).$$

The above clearly gives the required estimate. □

Now we are in a position to prove the following result which characterizes the image of $\mathcal{S}(\mathbb{R})$ under $e^{-t\mathcal{H}_\alpha}$.

Theorem 4. *Let $t > 0$ be fixed, and F be a holomorphic function on \mathbb{C} . Then there exists a function $f \in \mathcal{S}(\mathbb{R})$ such that $F = e^{-t\mathcal{H}_\alpha} f$ if and only if F satisfies*

$$|F(z)|^2 \leq C_{t,\alpha,m} \frac{e^{-\tanh(2t)x^2 + \coth(2t)y^2}}{(1+x^2+y^2)^{2m}}$$

for some constants $C_{t,\alpha,m}$, $m = 1, 2, 3, \dots$

PROOF: If $f \in \mathcal{S}(\mathbb{R})$, then $(\mathcal{H}_\alpha)^m f \in L^2_\alpha(\mathbb{R})$ for all integer m , so $f \in \mathcal{W}^{m,2}_{\mathcal{H}_\alpha}(\mathbb{R})$ for all m , which implies that

$$F = e^{-t\mathcal{H}_\alpha} f \in \mathcal{W}^{m,2}_{t,\alpha}(\mathbb{C}) \text{ for all } m.$$

From Theorem 3, we have $|F(z)|^2$ is bounded by $C_{t,\alpha,m} \frac{e^{-\tanh(2t)x^2 + \coth(2t)y^2}}{(1+x^2+y^2)^{2m}}$ for all m .

Conversely, suppose F satisfies the necessity condition. Using [6, p.140],

$$(4) \quad K_\alpha(z) = \left(\frac{\pi}{2z}\right)^{\frac{1}{2}} \frac{e^{-z}}{\Gamma(\alpha + \frac{1}{2})} \int_0^{+\infty} e^{-s} s^{\alpha-\frac{1}{2}} \left(1 + \frac{s}{2z}\right)^{\alpha-\frac{1}{2}} ds$$

for $|\arg z| < \pi, \alpha > -\frac{1}{2}$,

then by choosing m large enough, we see that

$$\int_{\mathbb{C}} |F_e(z)|^2 U_{t,e}^\alpha(z) dz + \int_{\mathbb{C}} |F_o(z)|^2 U_{t,o}^\alpha(z) dz < +\infty,$$

from which it follows that $F \in \mathcal{H}_t^\alpha(\mathbb{C})$, thus there exists a function $f \in L^2_\alpha(\mathbb{R})$ such that $F = e^{-t\mathcal{H}_\alpha} f$.

We have

$$K_\alpha\left(\frac{|z|^2}{\sinh 4t}\right) \times |z|^{2\alpha+2} = \left(\frac{\pi \sinh 4t}{2}\right)^{\frac{1}{2}} \frac{|z|^2}{\Gamma(\alpha + \frac{1}{2})} \\ \times e^{-\frac{|z|^2}{\sinh 4t}} \int_0^{+\infty} e^{-s} s^{\alpha-\frac{1}{2}} \left(|z|^2 + \frac{s(\sinh 4t)}{2}\right)^{\alpha-\frac{1}{2}} ds,$$

so it is an easy matter to see that $\frac{d^{2m}}{dt^{2m}} U_{t,e}^\alpha(z)$ and $\frac{d^{2m}}{dt^{2m}} U_{t,o}^\alpha(z)$ are a sum of $(2m+1)$ terms times $e^{\tanh(2t)x^2 - \coth(2t)y^2}$, where each term is of the form

$$(p(t,\alpha)x^2 + q(t,\alpha)y^2 + c(t,\alpha))^k \leq C_{t,\alpha} (1+x^2+y^2)^{2m} \text{ with } k \leq 2m,$$

where $p(t,\alpha)$, $q(t,\alpha)$ and $c(t,\alpha)$ are real constants. In view of Theorem 2, it follows that $F \in \mathcal{B}_t^{m,\alpha}(\mathbb{C}) = \mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$. This leads to the fact that $F \in \mathcal{W}_{t,\alpha}^{m,2}(\mathbb{C})$

for all m . Consequently $f \in \mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ for all m . Since

$$\bigcap_m \mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R}) = \mathcal{S}(\mathbb{R}),$$

the result follows. □

3.2 The image of $L_\alpha^p(\mathbb{R})$ under the Dunkl-Hermite semigroup. We begin this subsection by recalling that in [2] the authors have proved that the Dunkl-Hermite semigroup initially defined on $L_\alpha^2 \cap L_\alpha^p(\mathbb{R})$ extends to the whole of L_α^p and we have

$$\|e^{-t\mathcal{H}_\alpha} f\|_{\alpha,p} \leq (\cosh(2t))^{-(\alpha+1)} \|f\|_{\alpha,p}.$$

In the following, we give a characterization of the image of L_α^p under the Dunkl-Hermite semigroup.

Theorem 5. Fix $t > 0$ and let $1 < p < \infty$. Then for all $f \in L_\alpha^p(\mathbb{R})$, we have

$$|e^{-t\mathcal{H}_\alpha} f(x + iy)| \leq C_{t,p,\alpha} \|f\|_{p,\alpha} \exp\left(\left(\frac{p}{(p-1)\sinh 4t} - \frac{\coth 2t}{2}\right)x^2 + \frac{\coth 2t}{2}y^2\right).$$

PROOF: As we have shown previously, we have

$$e^{-t\mathcal{H}_\alpha} f(z) = e^{-\frac{1}{2}\left(\frac{\cosh 2t - 1}{\sinh 2t}\right)z^2} \left(e^{-\frac{1}{2}\left(\frac{\cosh 2t - 1}{\sinh 2t}\right)x^2} f *_\alpha q_{\frac{\sinh 2t}{2}}\right)(z),$$

so

$$|e^{-t\mathcal{H}_\alpha} f(x + iy)| \leq \frac{1}{\Gamma(\alpha + 1)} (2 \sinh 2t)^{-(\alpha+1)} e^{-\frac{\coth 2t}{2}(x^2 - y^2)} I_{t,\alpha},$$

where

$$I_{t,\alpha} = \int_{\mathbb{R}} |f(s)| \left| e^{-\frac{\coth 2t}{2}s^2} E_\alpha\left(\frac{s}{\sinh 2t}, z\right) \right| |s|^{2\alpha+1} ds.$$

So by Hölder’s inequality, we have

$$I_{t,\alpha} \leq \|f\|_{p,\alpha} \left\| e^{-\frac{\coth 2t}{2}s^2} E_\alpha\left(\frac{s}{\sinh 2t}, z\right) \right\|_{p',\alpha},$$

where p' is such that $\frac{1}{p} + \frac{1}{p'} = 1$.

We know that

$$\left| E_\alpha\left(\frac{s}{\sinh 2t}, z\right) \right|^{p'} \leq e^{\frac{p'sx}{\sinh 2t}},$$

so

$$\left\| e^{-\frac{\coth 2t}{2}s^2} E_\alpha\left(\frac{s}{\sinh 2t}, z\right) \right\|_{p',\alpha}^{p'} \leq \int_{\mathbb{R}} e^{-\frac{\coth 2t}{2}p's^2} e^{\frac{p'sx}{\sinh 2t}} |s|^{2\alpha+1} ds.$$

We can easily verify that

$$e^{-\frac{\coth 2t}{2}p's^2} e^{\frac{p'sx}{\sinh 2t}} = e^{\frac{p'x^2}{\sinh 4t}} e^{-\frac{p'}{2}(\sqrt{\coth 2t}s - \sqrt{\frac{2}{\sinh 4t}}x)^2}$$

which completes the proof. □

Notation 4. We denote by $V_{t, \frac{p}{2}}(z)$ the function defined by

$$V_{t, \frac{p}{2}}(x + iy) = \exp \left(-2 \left(\frac{p}{(p-1) \sinh 4t} x^2 + \frac{\coth 2t}{2} y^2 \right) \right)$$

and by $V_{t, \frac{p}{2}}^s$, the s -th power of $V_{t, \frac{p}{2}}$.

We write $\mathcal{HL}_\alpha^p(\mathbb{C}, V_{t, \frac{p}{2}}(z))$ for the class of holomorphic functions in $L_\alpha^p(\mathbb{C}, V_{t, \frac{p}{2}}(z))$.

The next corollary follows from Theorem 5, by a straightforward computation.

Corollary 1. *Let $f \in L_\alpha^p(\mathbb{R})$, $1 < p < \infty$ and fix $t > 0$, then*

- (i) $e^{-t\mathcal{H}_\alpha}(f) \in \mathcal{HL}_\alpha^p(\mathbb{C}, V_{t, \frac{p}{2}}^{\frac{p+\epsilon}{2}})$, for $\epsilon > 0$.

So

$$e^{-t\mathcal{H}_\alpha}(f) \in \bigcap_{\epsilon > 0} \mathcal{HL}_\alpha^p(\mathbb{C}, V_{t, \frac{p}{2}}^{\frac{p+\epsilon}{2}}).$$

- (ii) $e^{-t\mathcal{H}_\alpha}(f) \in \mathcal{HL}'_\alpha(\mathbb{C}, V_{t, \frac{p}{2}}^{\frac{p+\epsilon}{2}})$, for $\epsilon > 0$, where $2 \leq p < \infty$ and $\frac{1}{p} + \frac{1}{p'} = 1$.

So

$$e^{-t\mathcal{H}_\alpha}(f) \in \bigcap_{\epsilon > 0} \mathcal{HL}'_\alpha(\mathbb{C}, V_{t, \frac{p}{2}}^{\frac{p+\epsilon}{2}}).$$

- (iii) $e^{-t\mathcal{H}_\alpha}(f) \in \mathcal{HL}_\alpha^s(\mathbb{C}, V_{t, \frac{p}{2}}^{\frac{s+\epsilon}{2}})$, for $\epsilon > 0$, where $1 \leq s < \infty$.

4. Paley Wiener type Theorems

In this section we establish Paley-Wiener type theorems for the tempered distributions and the compactly supported distributions under the Dunkl-Hermite semigroup.

Theorem 6. *Let m be a positive integer. Then every $F \in \mathcal{W}_{t, \alpha}^{-m, 2}(\mathbb{C})$ satisfies the estimate*

$$|F(z)|^2 \leq C_{t, \alpha} (1 + |z|^2)^{2m} \exp \left(-\tanh(2t)x^2 + \coth(2t)y^2 \right).$$

Conversely, if an entire function F satisfies the above estimate, then F belongs to $\mathcal{W}_{t, \alpha}^{-m-1, 2}(\mathbb{C})$.

PROOF: It is easy to see that the reproducing kernel for $\mathcal{W}_{t, \alpha}^{-m, 2}(\mathbb{C})$ is given by

$$\mathcal{N}_t^{\alpha, -2m}(z, w) = \sum_n (2n + 2\alpha + 2)^{2m} \widetilde{h}_n^\alpha(\bar{z}) \widetilde{h}_n^\alpha(w).$$

So we only need to estimate the $(2m)$ -th derivate of $\mathcal{N}_t^\alpha(z, z)$ with respect to t .

Thanks to inequality (3), we have

$$\frac{d^{2m}}{dt^{2m}} \mathcal{N}_t^\alpha(z, z) \leq C_{t, \alpha} (1 + |z|^2)^{2m} e^{-\tanh(2t)x^2 + \coth(2t)y^2}.$$

Then if $F \in \mathcal{W}_{t,\alpha}^{-m,2}(\mathbb{C})$

$$|F(z)|^2 \leq C_{t,\alpha}(1 + |z|^2)^{2m} e^{-\tanh(2t)x^2 + \coth(2t)y^2}.$$

To prove the converse, we need to make use of duality between $\mathcal{W}_{\mathcal{H}_\alpha}^{m,2}(\mathbb{R})$ and $\mathcal{W}_{\mathcal{H}_\alpha}^{-m,2}(\mathbb{R})$.

The duality bracket is given by

$$\langle F, G \rangle = \int_{\mathbb{C}} F_e(z) \overline{G_e(z)} U_{t,e}^\alpha(z) dz + \int_{\mathbb{C}} F_o(z) \overline{G_o(z)} U_{t,o}^\alpha(z) dz.$$

If F satisfies the given estimates then F_e and F_o satisfy them too, and for any $G \in \mathcal{W}_{t,\alpha}^{m+1,2}(\mathbb{C})$ the integral defining $\langle F, G \rangle$ converges and hence F defines a continuous linear functional on $\mathcal{W}_{t,\alpha}^{m+1,2}(\mathbb{C})$.

Consequently, F belongs to $\mathcal{W}_{t,\alpha}^{-m-1,2}(\mathbb{C})$ which proves the converse. □

We recall the following definition given in [14].

Definition 3. Let S be in $\mathcal{S}'(\mathbb{R})$ and φ in $\mathcal{S}(\mathbb{R})$, the Dunkl convolution product of S and φ is the function $S *_\alpha \varphi$ defined by

$$\forall x \in \mathbb{R}, S *_\alpha \varphi(x) = \langle S_y, \tau_{-y}^\alpha \varphi(x) \rangle,$$

where τ_y^α is the generalized translation associated with the Dunkl operator D_α (see [13]).

It was shown in [14] that $S *_\alpha \varphi$ is a \mathcal{C}^∞ function on \mathbb{R} and for all $n \in \mathbb{N}$, we have

$$D_\alpha^n (S *_\alpha \varphi) = S *_\alpha (D_\alpha^n \varphi) = (D_\alpha^n S) *_\alpha \varphi.$$

It can be obviously seen that for fixed $x \in \mathbb{R}$ and $t > 0$, the function

$$y \longrightarrow \mathcal{M}_t^\alpha(x, y) \in \mathcal{S}(\mathbb{R}).$$

Definition 4. The Dunkl-Hermite semigroup of a distribution S in $\mathcal{S}'(\mathbb{R})$ is defined by

$$e^{-t\mathcal{H}_\alpha}(S)(x) = \langle S_y, \mathcal{M}_t^\alpha(x, y) \rangle.$$

Remark 3. For $S \in \mathcal{S}'(\mathbb{R})$, we have

$$e^{-t\mathcal{H}_\alpha} S(x) = e^{-\frac{1}{2}(\frac{\cosh 2t - 1}{\sinh 2t})x^2} \left(e^{-\frac{1}{2}(\frac{\cosh 2t - 1}{\sinh 2t})y^2} S *_\alpha q_{\frac{\sinh 2t}{2}} \right)(x),$$

so $e^{-t\mathcal{H}_\alpha} S$ is a \mathcal{C}^∞ function on \mathbb{R} .

Theorem 7. Suppose F is a holomorphic function on \mathbb{C} . Then there exists a distribution $f \in \mathcal{S}'(\mathbb{R})$ with $F = e^{-t\mathcal{H}_\alpha} f$ if and only if F satisfies

$$|F(z)|^2 \leq C_{t,\alpha}(1 + |z|^2)^{2m} \exp(-\tanh(2t)x^2 + \coth(2t)y^2),$$

for some nonnegative integer m .

PROOF: Let $f \in \mathcal{S}'(\mathbb{R})$. Since the union of all $\mathcal{W}_{\mathcal{H}_\alpha}^{-m,2}(\mathbb{R})$ is $\mathcal{S}'(\mathbb{R})$, then there exists m such that $f \in \mathcal{W}_{\mathcal{H}_\alpha}^{-m,2}(\mathbb{R})$. Thus

$$e^{-t\mathcal{H}_\alpha} f \in \mathcal{W}_{t,\alpha}^{-m,2}(\mathbb{C}),$$

and from Theorem 6 we have the result.

Conversely, suppose that F satisfies the hypothesis, then F belongs to $\mathcal{W}_{t,\alpha}^{-m-1,2}(\mathbb{C})$ and $F = e^{-t\mathcal{H}_\alpha} f$ with $f \in \mathcal{W}_{\mathcal{H}_\alpha}^{-m-1,2}(\mathbb{R})$. Then $f \in \mathcal{S}'(\mathbb{R})$. \square

In [7], the authors introduced the generalized windowed transform associated with D_α as follows. Given a function g in the Schwartz space, the windowed Dunkl transform of a regular function f , with window g , is defined by

$$\mathcal{V}_g^\alpha(f)(x, y) = \int_{\mathbb{R}} f(u) \tau_{-y}^\alpha g(u) E_\alpha(-ix, u) |u|^{2\alpha+1} du.$$

Here we extend this definition to the tempered distribution.

Definition 5. The windowed Dunkl transform of a tempered distribution S with window $g \in \mathcal{S}(\mathbb{R})$ is defined by

$$\mathcal{V}_g^\alpha(S)(x, y) = \langle S, \tau_{-y}^\alpha g E_\alpha(-ix, \cdot) \rangle.$$

When S is given by the function $f|u|^{2\alpha+1}$, $S = S_{f|u|^{2\alpha+1}}$, then

$$\mathcal{V}_g^\alpha(S_{f|u|^{2\alpha+1}})(x, y) = \int_{\mathbb{R}} f(u) \tau_{-y}^\alpha g(u) E_\alpha(-ix, u) |u|^{2\alpha+1} du,$$

which we write simply $\mathcal{V}_g^\alpha(f)(x, y)$.

In the case where $g(x) = \varphi_a(x) = e^{-\frac{1}{2}ax^2}$, for $a > 0$, $\mathcal{V}_{\varphi_a}^\alpha f$ is called gaussian Dunkl windowed transform. In our context, we are interested in the case $y = 0$ and we denote

$$\mathcal{T}_a^\alpha f(x) = \mathcal{V}_{\varphi_a}^\alpha(f)(x, 0).$$

Hence, for $a > 0$, the transform \mathcal{T}_a^α is defined by

$$\mathcal{T}_a^\alpha(S)(x) = \langle S, e^{-\frac{1}{2}a(\cdot)^2} E_\alpha(-ix, \cdot) \rangle, \quad S \in \mathcal{S}'(\mathbb{R}).$$

If $f \in \mathcal{S}(\mathbb{R})$ we have

$$\mathcal{T}_a^\alpha(f)(x) = \int_{\mathbb{R}} f(u) e^{-\frac{1}{2}au^2} E_\alpha(-ix, u) |u|^{2\alpha+1} du.$$

We see that $\mathcal{T}_a^\alpha f$ extends to \mathbb{C} as an entire function even when f is in $\mathcal{S}'(\mathbb{R})$. This property of \mathcal{T}_a^α allows us to prove the following analogue of Paley-Wiener theorem given by Trimèche in [13].

Theorem 8. For any $a > 0$ the transform \mathcal{T}_a^α of a tempered distribution f on \mathbb{R} extends to \mathbb{C} as an entire function which satisfies the estimate

$$|\mathcal{T}_a^\alpha f(z)| \leq C_\alpha(1 + x^2 + y^2)^m e^{\frac{1}{2}a^{-1}y^2}$$

for some non-negative integer m .

Conversely, if an entire function F satisfies such an estimate, then $F = \mathcal{T}_a^\alpha f$ for some tempered distribution f .

PROOF: We relate the transform $\mathcal{T}_a^\alpha f$ to $e^{-t\mathcal{H}_\alpha} f$. Indeed, considering the case $a > 1$ first and writing $a = \coth 2t$ for some $t > 0$, we can easily verify that

$$e^{-t\mathcal{H}_\alpha} f(z) = \frac{1}{\Gamma(\alpha + 1)(2 \sinh 2t)^{\alpha+1}} e^{-\frac{1}{2} \coth 2t z^2} \mathcal{T}_a^\alpha f\left(\frac{iz}{\sinh 2t}\right) \quad \forall z \in \mathbb{C}.$$

We obtain the required estimate on $\mathcal{T}_a^\alpha f(z)$ by applying Theorem 7.

Conversely, if F satisfies the given estimates then again by Theorem 7 the function

$$G(z) = \frac{1}{\Gamma(\alpha + 1)(2 \sinh 2t)^{\alpha+1}} e^{-\frac{1}{2} \coth 2t z^2} F\left(\frac{iz}{\sinh 2t}\right)$$

should be of the form $e^{-t\mathcal{H}_\alpha} f(z)$ with a tempered distribution f .

When $a < 1$ we take $t > 0$ so that $a = \tanh 2t$ and the proof requires an analogue of Theorem 7 for functions of the form $e^{-(t+i\frac{\pi}{4})\mathcal{H}_\alpha} f$ (see [1]).

The image of tempered distributions under $e^{-(t+i\frac{\pi}{4})\mathcal{H}_\alpha}$ can be characterized in a similar way. The final estimates do not depend on the factor $e^{-i\frac{\pi}{4}\mathcal{H}_\alpha}$ which is just the Dunkl transform \mathcal{F}_D .

Here the Dunkl transform of a distribution f in $\mathcal{S}'(\mathbb{R})$ is defined by

$$\langle \mathcal{F}_D(f), \psi \rangle = \langle f, \mathcal{F}_D(\psi) \rangle, \quad \psi \in \mathcal{S}(\mathbb{R})$$

and for $f \in \mathcal{S}(\mathbb{R})$

$$\mathcal{F}_D(f)(x) = \int_{\mathbb{R}} f(y) E_\alpha(-ix, y) |y|^{2\alpha+1} dy.$$

We have

$$e^{-(t+i\frac{\pi}{4})\mathcal{H}_\alpha} f = e^{-t\mathcal{H}_\alpha} (e^{-i\frac{\pi}{4}\mathcal{H}_\alpha} f)$$

and

$$e^{-i\frac{\pi}{4}\mathcal{H}_\alpha} f = \frac{1}{2^{\alpha+1}\Gamma(\alpha + 1)} e^{(\alpha+1)i\frac{\pi}{2}} \mathcal{F}_D f.$$

We know that \mathcal{F}_D is an isomorphism from $\mathcal{S}'(\mathbb{R})$ onto $\mathcal{S}'(\mathbb{R})$ (see [13]), so we have the analogue of Theorem 7. □

Finally, we remark that we also have the following result which characterizes the image of compactly supported distributions under the Dunkl-Hermite semigroup.

Theorem 9. *Let f be a distribution supported in a ball of radius R centered at the origin. Then for any $t > 0$ the function $e^{-t\mathcal{H}_\alpha} f$ extends to \mathbb{C} as an entire function which satisfies*

$$|e^{-t\mathcal{H}_\alpha} f(z)| \leq C e^{-\frac{1}{2} \coth 2t(x^2 - y^2)} e^{\frac{R|x|}{\sinh 2t}},$$

with C being a positive constant.

Conversely, any entire function F satisfying the above estimate is of the form $e^{-t\mathcal{H}_\alpha} f$ where f is supported inside a ball of radius R centered at the origin.

PROOF: We have to relate the Dunkl-Hermite semigroup and the Dunkl transform in $\mathcal{E}'(\mathbb{R})$

$$e^{-t\mathcal{H}_\alpha} S(z) = \frac{1}{\Gamma(\alpha + 1)(2 \sinh(2t))^{\alpha+1}} e^{-\frac{1}{2} \coth 2tz^2} \mathcal{F}_D [S_y e^{-\frac{1}{2} \coth 2ty^2}] \left(\frac{iz}{\sinh 2t} \right).$$

Here the Dunkl transform of a distribution S in $\mathcal{E}'(\mathbb{R})$ is defined by

$$\forall y \in \mathbb{R}, \mathcal{F}_D(S)(y) = \langle S_x, E_\alpha(-iy, x) \rangle.$$

We obtain the necessity condition by appealing Theorem 5.3 given in [13], i.e., Paley-Wiener theorem for compactly supported distributions and the Dunkl transform.

Conversely, if F satisfies the given estimates then again by the same Theorem 5.3, the function

$$G(z) = \Gamma(\alpha + 1)(2 \sinh(2t))^{\alpha+1} e^{-\frac{1}{4} \sinh 4tz^2} F(-iz \sinh 2t)$$

should be of the form $\mathcal{F}_D(f)$ for a distribution f supported inside a ball of radius R centered at the origin and

$$F(z) = e^{-t\mathcal{H}_\alpha} (f(y)e^{\frac{1}{2} \coth 2ty^2})(z),$$

where $f(y)e^{\frac{1}{2} \coth 2ty^2}$ is also a distribution supported inside a ball of radius R centered at the origin. This completes the proof of the theorem. □

REFERENCES

- [1] Ben Salem N., Nefzi W., *Inversion of the Dunkl-Hermite semigroup*, Bull. Malays. Math. Sci. Soc. (2) **35** (2012), no. 2, 287–301.
- [2] Ben Salem N., Samaali T., *Hilbert transforms associated with Dunkl-Hermite polynomials*, SIGMA Symmetry Integrability Geom. Methods Appl. **5** (2009), Paper 037.
- [3] Dunkl C.F., *Integral kernels with reflection group invariance*, Canad. J. Math. **43** (1991), 1213–1227.
- [4] Erdelyi A. et al., *Higher Transcendental Functions, vol 2*, McGraw-Hill, New-York, 1953.
- [5] de Jeu M.F.E., *The Dunkl transform*, Invent. Math. **113** (1993), 147–162.
- [6] Lebedev N.N., *Special Functions and their Applications*, translated by R.A. Silverman, Dover, New York, 1972.

- [7] Maalaoui R., Trimèche K., *A family of generalized windowed transforms associated with the Dunkl operators on \mathbb{R}^d* , Integral Transforms Spec. Funct. **23** (2012), no. 3, 191–206.
- [8] Radha R., Thangavelu S., *Holomorphic Sobolev spaces, Hermite and special Hermite semigroups and a Paley-Wiener theorem for the windowed Fourier transform*, J. Math. Anal. Appl. **354** (2009), 564–574.
- [9] Radha R., Venku Naidu D., *Image of $L^p(\mathbb{R}^n)$ under the Hermite semigroup*. Int. J. Math. Math. Sci. (2008), Art. ID 287218, 13 pages.
- [10] Rosenblum M., *Generalized Hermite polynomials and the Bose-like oscillator calculus*, in Operator theory: Advances and Applications, Vol. 73, Birkhäuser, Basel, 1994, pp. 369–396.
- [11] Rösler M., *Generalized Hermite polynomials and the heat equation for Dunkl operators*, Comm. Math. Phys. **192** (1998), no. 3, 519–542.
- [12] Rösler M., Voit M., *Markov Processes related with Dunkl operators*, Adv. Appl. Math. **21** (1998), 575–643.
- [13] Trimèche K., *Paley-Wiener theorems for the Dunkl transform and Dunkl translation operators*, Integral Transforms Spec. Funct. **13** (2002), 17–38.
- [14] Trimèche K., *Hypoelliptic Dunkl convolution equations in the space of distributions on \mathbb{R}^d* , J. Fourier Anal. Appl. **12** (2006), 517–542.

FACULTY OF SCIENCES OF TUNIS, UNIVERSITY OF TUNIS EL MANAR, CAMPUS UNIVER-
SITAIRE, 2092 TUNIS, TUNISIA

E-mail: Nejib.BenSalem@fst.rnu.tn
walidahla@yahoo.fr

(Received November 16, 2012)