

Radon-Nikodym property

SURJIT SINGH KHURANA

Abstract. For a Banach space E and a probability space $(X, \mathcal{A}, \lambda)$, a new proof is given that a measure $\mu : \mathcal{A} \rightarrow E$, with $\mu \ll \lambda$, has RN derivative with respect to λ iff there is a compact or a weakly compact $C \subset E$ such that $|\mu|_C : \mathcal{A} \rightarrow [0, \infty]$ is a finite valued countably additive measure. Here we define $|\mu|_C(A) = \sup\{\sum_k |\langle \mu(A_k), f_k \rangle|\}$ where $\{A_k\}$ is a finite disjoint collection of elements from \mathcal{A} , each contained in A , and $\{f_k\} \subset E'$ satisfies $\sup_k |f_k(C)| \leq 1$. Then the result is extended to the case when E is a Frechet space.

Keywords: liftings; lifting topology; weakly compact sets; Radon-Nikodym derivative

Classification: Primary 46B22, 46G05, 46G10, 28A51; Secondary 60B05, 28B05, 28C05

1. Introduction and notations

In this paper K will always denote the field of real or complex numbers (we will call them scalars), \mathbb{R} the real numbers and \mathbb{N} the set of natural numbers. All locally convex space are assumed to be Hausdorff and are over K and notations and results of [8] will be used. Given a locally convex space E with E' its dual, for $x \in E$ and $f \in E'$, we will also write $\langle f, x \rangle = \langle x, f \rangle$ for $f(x)$; for an $A \subset E$, $\Gamma(A)$ will denote the absolute convex hull of A . Let $(X, \mathcal{A}, \lambda)$ be a complete probability space. By a measure we will always mean a countably additive measure. For a measure μ , $|\mu|$ will denote its total variation measure. For measures and vector measures we refer to [2]; see also [4], [5].

In [3], for a Banach space E , an interesting characterization is given for a vector measure $\mu : \mathcal{A} \rightarrow E$ of bounded variation, $\mu \ll \lambda$, to have a derivative. It is proved that μ has derivative iff there is a compact or weakly compact $C \subset E$ such that $|\mu|_C : \mathcal{A} \rightarrow [0, \infty]$ is a finite valued countably additive measure. Here we define $|\mu|_C(A) = \sup\{\sum_k |\langle \mu(A_k), f_k \rangle|\}$ where $\{A_k\}$ is a finite disjoint collection of elements from \mathcal{A} , each contained in A , and $\{f_k\} \subset E'$ satisfies $\sup_k |f_k(C)| \leq 1$. First we give a new proof of this result and then extend this to Frechet spaces.

2. Main results

Theorem 1. *Let \mathcal{A} be a σ -algebra of subsets of a set X , E a Frechet space and $\mu : \mathcal{A} \rightarrow E$ countably additive measure with $\mu \ll \lambda$. Suppose μ has finite variation with respect to every continuous semi-norm on E . Then μ has RN derivative relative to λ iff there is a compact or weakly compact $C \subset E$ such that $|\mu|_C : \mathcal{A} \rightarrow [0, \infty]$ is a finite-valued measure. Here we define $|\mu|_C(A) = \sup\{\sum_k |\langle \mu(A_k), f_k \rangle|\}$ where $\{A_k\}$ is a finite disjoint collection of elements from \mathcal{A} , each contained in A , and $\{f_k\} \subset E'$ satisfies $\sup_k |f_k(C)| \leq 1$.*

PROOF: First we consider E to be a Banach and give an entirely different proof than the one given in [3]; we will reduce it to a reflexive Banach subspace of E with a finer topology. Take an absolutely convex weakly compact $C \subset E$ with a countably additive measure $\nu = |\mu|_C \leq 1$. This implies $\mu(\mathcal{A}) \subset C$ ([3, Theorem 2.1(1)]). By [1], there is a reflexive Banach space $E_0 \subset E$ such that $C \subset E_0$, C is weakly compact in E_0 , and the identity mapping $E_0 \rightarrow E$ is continuous. Take an $f_0 \in E'_0$ with norm ≤ 1 and fix $c > 0$. If we consider C as a subset of E , $(f_0)|_C$ is an affine continuous function on C . It is proved in [7, Proposition 3.5, p.31] that $(E'_C + K)$ is uniformly dense in the space of all continuous affine functions on C (this is proved when $K = \mathbb{R}$ but easily extends to general K). Thus there is an $f \in E'$ and $r \in K$ such that $\sup |(f + r - f_0)(C)| \leq c$. Since C is absolutely convex (that implies $0 \in C$), we get $\sup |(f - f_0)(C)| \leq 2c$. Take a decreasing sequence $\{A_n\} \subset \mathcal{A}$ such that $A_n \downarrow \emptyset$. Now $f \circ \mu(A_n) \rightarrow 0$ and since $\sup |(f - f_0)(C)| \leq 2c$ and c is arbitrarily small, we get $f_0 \circ \mu(A_n) \rightarrow 0$ and so $\mu : \mathcal{A} \rightarrow E_0$ is countably additive. Now we will prove that $\mu : \mathcal{A} \rightarrow E_0$ is of bounded variation. Take $p > 0$ with $pC \subset B$ (the unit ball of E_0), a finite collection $\{f_i\}$ in the closed unit ball of E'_0 , and disjoint elements $\{A_i\} \subset \mathcal{A}$. We have $|pf_i(C)| \leq 1 \forall i$. As explained above, take $\{f'_i\} \subset E'$ with $\sup |(pf_i - f'_i)(C)| \leq \frac{1}{2^i} \forall i$. We get

$$\sum |\langle f_i, \mu(A_i) \rangle| \leq \frac{1}{p} \sum |(pf_i - f'_i)\mu(A_i)| + \frac{1}{p} \sum |(f'_i)\mu(A_i)| \leq \frac{1}{p} + \frac{1}{p} |\mu|_C(X) \leq \frac{2}{p}.$$

This proves $\mu : \mathcal{A} \rightarrow E_0$ is of bounded variation. Since E_0 is reflexive, there is an $h \in L_1(X, E_0)$ with $\mu = h\lambda$. From this it easily follows that $h \in L_1(X, E)$. The converse is same as for Frechet space which we will consider now.

Now we consider the case when E is a Frechet space. Suppose μ has RN derivative $\frac{d\mu}{d\lambda} = g \in L^1(\lambda, E) = L^1(\lambda) \hat{\otimes} E$ (the completion in projective tensor product). Thus $g = \sum_i \alpha_i g_i x_i$, $\{g_i\}$, $\{x_i\}$ being null sequences in $L^1(\lambda)$ and E respectively and $\{\alpha_i\} \in \ell_1$ ([8, Theorem 6.4, p. 94]); we can assume that $\int |g_i| d\lambda \leq 1 \forall i$. Let C be the closed, absolutely convex hull of $\{x_i\}$; C is compact. Take a finite, disjoint family $\{A_k\}$ of elements of \mathcal{A} and $\{f_k\}$ elements of E' with $\sup |f_k(C)| \leq 1 \forall k$. We have $\sum_k |\langle \mu(A_k), f_k \rangle| \leq \sum_k \sum_i \int_{A_k} |\alpha_i| |g_i| |f_k(x_i)| d\lambda \leq \alpha$ where $\alpha = \sum_i |\alpha_i|$. Thus $|\mu|_C$ is finite-valued. It is a routine verification that $|\mu|_C$ is countably additive ([2, p. 4], [3, Theorem 2.1(5), p. 142]).

Conversely suppose for an absolutely convex weakly compact $C \subset E$, $\nu = |\mu|_C$ is finite-valued. We have $\mu \ll |\mu|_C$. Also it follows from the definition of $|\mu|_C$ that, for an $f \in E'$ with $\sup |f(C)| \leq 1$, we have $|\mu|_C \geq |f \circ \mu|$. Denoting the completion of $|\mu|_C$ by $|\mu|_C$ again, we fix a lifting ρ_0 for this measure ([9]) and take the lifting topology \mathcal{T}_0 on X which has $\{\rho_0(A) : A \in \mathcal{A}\}$ as the base of open sets; we can assume this topology to be Hausdorff and denote by $C_b(X)$ all scalar-valued bounded continuous functions on X . For each $f \in E'$ there is a $\phi_f \in L_1(|\mu|_C)$ such that $f \circ \mu = \phi_f |\mu|_C$. Put $\sup |f(C)| = p$; we claim $|\phi_f| \leq p$ a.e. $[|\mu|_C]$. Suppose this is not true. Then there is a $c > 0$ such that $|\mu|_C(A) > 0$ where $A = \{x \in X : |\phi_f(x)| \geq p + c\}$. Now, since $|\frac{1}{p}f(C)| \leq 1$, we have $|\mu|_C(A) \geq \frac{1}{p}|f \circ \mu|(A) = \frac{1}{p} \int_A |\phi_f| d|\mu|_C \geq \frac{1}{p}(p + c)|\mu|_C(A)$ which is a contradiction. Thus there is a unique function in $C_b(X)$ which is equal to ϕ_f a.e. $[|\mu|_C]$; we denote this function also by ϕ_f .

Define $\phi : X \rightarrow K^{E'}$, $(\phi(x))_f = \phi_f(x)$. It is a simple verification that $\phi_{f_1+f_2} = \phi_{f_1} + \phi_{f_2}$ and $\phi_{rf} = r\phi_f$ for any $f_1, f_2, f \in E'$ and any $r \in K$. Also E , with weak topology, can be considered as a subspace of $K^{E'}$ with product topology. We claim that $\phi(X) \subset C$. If this is not true, by separation theorem ([8, 9.2, p. 65]), $\exists x_0 \in X$ and $f \in E'$ such that $p = \sup |f(C)| < Rl(\phi_f(x_0)) = p + 3\eta$ for some $\eta > 0$ (note C is absolutely convex and so $\sup(Rl(f(C))) = \sup |f(C)|$). Now $\nu(A) > 0$ where $A = \{x \in X : |\phi_f(x)| > p + 2\eta\}$. Since $|\phi_f| \leq p$, we have $p\nu(A) \geq \int_A |\phi_f| d\nu \geq (p + 2\eta)\nu(A)$, a contradiction. By [6, Theorem 2, p. 389], ϕ is weakly equivalent to a function ϕ_0 such that $\phi_0(X)$ is contained in a separable weakly compact subset of E ; thus ϕ_0 is bounded. Now it is well-known that if a weakly measurable function has a separable range in E then it is strongly measurable ([2, Theorem 2, p. 42]; it is proved for a Banach space but thus easily extends to a Fréchet space). Now being bounded and strongly measurable, $\phi_0 \in L^1(\nu, E)$. Since $|\mu| \ll \nu$ and $\nu \ll \lambda$, the result follows. \square

Acknowledgment. We are very thankful to the referee for pointing out several typographical errors and also making some very useful suggestions which has improved the paper.

REFERENCES

- [1] Davis W.J., Figiel T., Johnson W.B., Pelczynski A., *Factoring weakly compact operators*, J. Funct. Anal. **17** (1974), 311–327.
- [2] Diestel J., Uhl J.J., *Vector Measures*, Amer. Math. Soc. Surveys, 15, American Mathematical Society, Providence, RI, 1977.
- [3] Gruenwald M.E., Wheeler R.F., *A strict representation of $L_1(\mu, X)$* , J. Math. Anal. Appl. **155** (1991), 140–155.
- [4] Khurana S.S., *Topologies on spaces of continuous vector-valued functions*, Trans Amer. Math. Soc. **241** (1978), 195–211.
- [5] Khurana S.S., *Topologies on spaces of continuous vector-valued functions II*, Math. Ann. **234** (1978), 159–166.
- [6] Khurana S.S., *Pointwise compactness and measurability*, Pacific J. Math. **83** (1979), 387–391.

- [7] Phelps R.R., *Lectures on Choquet's Theorem*, D. van Nostrand Company, Inc., Princeton, N.J.-Toronto, Ont.-London, 1966.
- [8] Schaefer H.H., *Topological Vector Spaces*, Springer, 1986.
- [9] Ionescu Tulcea A., Ionescu Tulcea C., *Topics in the theory of lifting*, Springer, New York, 1969.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOWA, IOWA CITY, IOWA 52242,
U.S.A.

E-mail: khurana@math.uiowa.edu

(Received June 13, 2017, revised August 23, 2017)