

Resolvability in c.c.c. generic extensions

LAJOS SOUKUP, ADRIENNE STANLEY

Abstract. Every crowded space X is ω -resolvable in the c.c.c. generic extension $V^{\text{Fn}(|X|,2)}$ of the ground model.

We investigate what we can say about λ -resolvability in c.c.c. generic extensions for $\lambda > \omega$.

A topological space is *monotonically ω_1 -resolvable* if there is a function $f : X \rightarrow \omega_1$ such that

$$\{x \in X : f(x) \geq \alpha\} \subset^{dense} X$$

for each $\alpha < \omega_1$.

We show that given a T_1 space X the following statements are equivalent:

- (1) X is ω_1 -resolvable in some c.c.c. generic extension;
- (2) X is monotonically ω_1 -resolvable;
- (3) X is ω_1 -resolvable in the Cohen-generic extension $V^{\text{Fn}(\omega_1,2)}$.

We investigate which spaces are monotonically ω_1 -resolvable. We show that if a topological space X is c.c.c., and $\omega_1 \leq \Delta(X) \leq |X| < \omega_\omega$, where $\Delta(X) = \min\{|G| : G \neq \emptyset \text{ open}\}$, then X is monotonically ω_1 -resolvable.

On the other hand, it is also consistent, modulo the existence of a measurable cardinal, that there is a space Y with $|Y| = \Delta(Y) = \aleph_\omega$ which is not monotonically ω_1 -resolvable.

The characterization of ω_1 -resolvability in c.c.c. generic extension raises the following question: is it true that crowded spaces from the ground model are ω -resolvable in $V^{\text{Fn}(\omega,2)}$?

We show that (i) if $V = L$ then every crowded c.c.c. space X is ω -resolvable in $V^{\text{Fn}(\omega,2)}$, (ii) if there are no weakly inaccessible cardinals, then every crowded space X is ω -resolvable in $V^{\text{Fn}(\omega_1,2)}$.

Moreover, it is also consistent, modulo a measurable cardinal, that there is a crowded space X with $|X| = \Delta(X) = \omega_1$ such that X remains irresolvable after adding a single Cohen real.

Keywords: resolvable; monotonically ω_1 -resolvable; measurable cardinal

Classification: 54A35, 03E35, 54A25

1. Introduction

Notion of resolvability was introduced and studied first by E. Hewitt [4], in 1943. A topological space X is κ -resolvable if it can be partitioned into κ many dense subspaces. X is *resolvable* iff it is 2-resolvable, and *irresolvable* otherwise.

DOI 10.14712/1213-7243.2015.226

The second author was supported by the Fulbright Scholar Program.

The preparation of this paper was supported by OTKA grant no. K113047.

Irresolvable spaces with many interesting extra properties were constructed, but there are no “absolute” examples for crowded irresolvable spaces, because if X is a crowded space, then clearly

$$V^{\text{Fn}(|X|,2)} \models X \text{ is } \omega\text{-resolvable.}$$

In this paper we investigate what we can say about λ -resolvability in c.c.c. generic extensions for $\lambda > \omega$.

To characterize spaces which are ω_1 -resolvable in some c.c.c. generic extension we introduce the notion of monotone κ -resolvability.

Definition 1.1. Let κ be an infinite cardinal. A topological space X is *monotonically κ -resolvable*[†] if there is a function $f : X \rightarrow \kappa$ such that

$$\{x \in X : f(x) \geq \alpha\} \subset^{\text{dense}} X$$

for each $\alpha < \kappa$. We will say that f *witnesses* that X is monotonically κ -resolvable.

Clearly a space X is monotonically κ -resolvable iff X has a partition $\{X_\zeta : \zeta < \kappa\}$ of X such that

$$\text{int} \left(\bigcup \{X_\zeta : \zeta < \xi\} \right) = \emptyset$$

for all $\xi < \kappa$.

Theorem 1.2. *Let X be a T_1 topological space. The following statements are equivalent:*

- (1) X is ω_1 -resolvable in some c.c.c. generic extension,
- (2) X is monotonically ω_1 -resolvable,
- (3) X is ω_1 -resolvable in the Cohen generic extension $V^{\text{Fn}(\omega_1,2)}$.

Which spaces are monotonically ω_1 -resolvable?

Theorem 1.3. *If a topological space X is c.c.c., and $\omega_1 \leq \Delta(X) \leq |X| < \omega_\omega$, then X is monotonically ω_1 -resolvable.*

Theorem 1.4. *If κ is a measurable cardinal, then there is a space X with $|X| = \Delta(X) = \kappa$ which is not monotonically ω_1 -resolvable.*

What about spaces of cardinality ω_ω ?

Theorem 1.5. *It is consistent, modulo the existence of a measurable cardinal, that there is a space X with $|X| = \Delta(X) = \omega_\omega$ which is not monotonically ω_1 -resolvable.*

Do we really need to add $|X|$ -many Cohen reals to make X resolvable?

[†]In [13] a “monotonically ω -resolvable” space is called “almost- ω -resolvable”. However, in [12] a space X is *almost- κ -resolvable* if it contains a family of κ dense sets with pairwise nowhere dense intersections.

Theorem 1.6. (1) *It is consistent, modulo a measurable cardinal, that there is a crowded space X with $|X| = \Delta(X) = \omega_1$ (so X is monotonically ω_1 -resolvable) such that*

$$V^{\text{Fn}(\omega,2)} \models \text{“}X \text{ is irresolvable.”}$$

(2) *If $V = L$, then every crowded space with $|X| = \Delta(X) = \text{cf}(|X|)$ is monotonically ω -resolvable, and so it is ω -resolvable in $V^{\text{Fn}(\omega,2)}$.*

(3) *If the cardinality of a crowded c.c.c. space X is less than the first weakly inaccessible cardinal, then X is ω -resolvable in $V^{\text{Fn}(\omega_1,2)}$ §.*

The almost resolvability of c.c.c. spaces was investigated by Pavlov in [11]: on page 53 Pavlov writes that mimicking Malykhin’s method, by using Ulam matrices, he showed that every crowded c.c.c. space of cardinality ω_1 is almost resolvable. In [3, Theorem 2.22] a stronger result was proved: a crowded c.c.c. space is almost resolvable if its cardinality is less than the first weakly inaccessible cardinal. Theorem 1.6(2) is a further improvement of this result because monotone ω -resolvability implies almost resolvability.

In [1, 3.12 Problem (2)] the authors ask *if every space with countable cellularity and cardinality less than the first inaccessible non-countable cardinal is almost- ω -resolvable*. As we will see Theorem 1.6(3) gives a positive answer to a weakening of this question.

2. Characterization of ω_1 -resolvability in c.c.c. extensions

Instead of Theorem 1.2 we prove the following stronger result. We say that a function $g : X \rightarrow \kappa$ witnesses that X is κ -resolvable if

$$\{x \in X : g(x) = \alpha\} \subset^{dense} X$$

for each $\alpha < \kappa$.

Theorem 2.1. *Assume that X is a crowded topological space and κ is an infinite cardinal. If $\kappa = \text{cf}([\kappa]^\omega, \subset)$ then the following statements are equivalent:*

- (1) *X is κ -resolvable in some c.c.c. generic extension;*
- (2) *there is a function $h : X \rightarrow [\kappa]^\omega$ such that $\bigcup h''U = \kappa$ for each non-empty open $U \subset X$;*
- (3) *X is κ -resolvable in the Cohen-generic extension $V^{\text{Fn}(\kappa,2)}$.*

PROOF: First we show that (1) \rightarrow (2). Assume that \mathbb{P} is a c.c.c. poset such that there is a function $g \in V^{\mathbb{P}}$ witnessing the κ -resolvability of X .

For each $x \in X$ define

$$h(x) = \{\alpha < \kappa : \exists p_\alpha^x \in \mathbb{P}(p_\alpha^x \Vdash \dot{g}(\check{x}) = \check{\alpha})\}.$$

Since the conditions $\{p_\alpha^x : \alpha \in h(x)\}$ are pairwise incomparable and \mathbb{P} is c.c.c., the set $h(x)$ is countable.

§ ω_1 is not a misprint here.

We now show that the function h defined above satisfies (2). Fix $\alpha < \kappa$ and U an open subset of X . We need to show that there exists $x \in U$ such that $\alpha \in h(x)$. Since

$$V^{\mathbb{P}} \models g^{-1}(\{\alpha\}) \subset^{dense} X$$

it follows that there is $x \in U$ such that

$$V^{\mathbb{P}} \models g(x) = \alpha.$$

Thus, there exists $p \in \mathbb{P}$ such that

$$p \Vdash \dot{g}(\dot{x}) = \check{\alpha}.$$

Then $\alpha \in h(x)$.

Next we show that (2) \rightarrow (3). Let \mathcal{A} be a cofinal subset of $[\kappa]^\omega$ with $|\mathcal{A}| = \kappa$. Let $\{A_\alpha : \alpha < \kappa\}$ be an enumeration of \mathcal{A} , and for each $x \in X$ pick

$$h^*(x) \in \mathcal{A} \text{ such that } h^*(x) \supset \bigcup_{\alpha \in h(x)} A_\alpha.$$

Then for all non-empty open U

$$(+) \quad \{h^*(x) : x \in U\} \text{ is cofinal in } [\kappa]^\omega.$$

Next we note that forcing with $\text{Fn}(\kappa, 2)$ is the same as forcing with $\text{Fn}(\kappa, \omega)$. Further, $\text{Fn}(\kappa, \omega)$ is isomorphic to

$$\mathbb{P} = \{p \in \text{Fn}(\mathcal{A}, \kappa) : \forall A \in \text{dom}(p) p(A) \in A\}.$$

Indeed, for each $A \in \mathcal{A}$ fix a bijection $\rho_A : \omega \rightarrow A$, and then for $q \in \text{Fn}(\kappa, \omega)$ define $\varphi(q) \in \mathbb{P}$ as follows:

- (i) $\text{dom}(\varphi(q)) = \{A_\alpha : \alpha \in \text{dom}(q)\}$, and
- (ii) $\varphi(q)(A_\alpha) = \rho_{A_\alpha}(q(\alpha))$ for $A_\alpha \in \text{dom}(\varphi(q))$.

Then φ is clearly an isomorphism between $\text{Fn}(\kappa, \omega)$ and \mathbb{P} .

We will proceed using \mathbb{P} .

Let G be a \mathbb{P} -generic filter, and let $g = \bigcup G$. Then $g \in V^{\mathbb{P}}$ and $g : \mathcal{A} \rightarrow \kappa$ is such that $g(A) \in A$.

We claim that $f = g \circ h^*$ witnesses that X is κ -resolvable.

Fix $\alpha < \kappa$ and an open $U \subset X$.

Let $q \in \mathbb{P}$ be arbitrary. Then, by (+), there is $x \in U$ such that

$$\{\alpha\} \cup \bigcup \text{dom}(q) \subsetneq h^*(x).$$

Then $h^*(x) \notin \text{dom}(q)$, and $\alpha \in h^*(x)$, so

$$p = q \cup \{\langle h^*(x), \alpha \rangle\} \in \mathbb{P}_1,$$

and

$$p \Vdash (g \circ h^*)(\check{x}) = \check{\alpha}.$$

Thus, by genericity, there is $p \in G$ and $x \in U$ such that

$$p \Vdash (g \circ h^*)(\dot{x}) = \check{\alpha}.$$

Hence

$$V^{\mathbb{P}} \models X \text{ is } \kappa\text{-resolvable.}$$

Finally (3) \rightarrow (1) is trivial. □

Problem 2.2. *Can we drop the assumption $\kappa = \text{cf}([\kappa]^\omega, \subset)$ from Theorem 2.1?*

3. On monotone ω_1 -resolvability of c.c.c. spaces

We start with an easy to prove observation.

Lemma 3.1. *Let X be a topological space and $\mathcal{B} \subset \mathcal{P}(X)$. If every $B \in \mathcal{B}$ is monotonically κ -resolvable, then so is $\overline{\bigcup \mathcal{B}}$. So every space contains a subspace which is the greatest monotonically κ -resolvable subspace (this subspace can be empty, of course).*

Corollary 3.2. *Let X be a topological space. Let Z be a dense subset of X . If Z is monotonically κ -resolvable, then X is also monotonically κ -resolvable.*

Before proving Theorem 1.3 we prove the following “stepping-down” theorem. The proof uses ideas from [8].

Theorem 3.3. *If X is a κ -c.c., monotonically κ^+ -resolvable space, then X is monotonically κ -resolvable as well.*

PROOF: Since an open subspace of a κ -c.c., monotonically κ^+ -resolvable space is also κ -c.c. and monotonically κ^+ -resolvable, by Lemma 3.1 it is enough to show that

- (*) every κ -c.c., monotonically κ^+ -resolvable space X has a monotonically κ -resolvable non-empty open subset.

Ulam [14] proved that there is a “matrix”

$$\langle M_{\alpha,\zeta} : \alpha < \kappa^+, \zeta < \kappa \rangle \subset \mathcal{P}(\kappa^+)$$

such that

- (i) $M_{\alpha,\xi} \cap M_{\beta,\xi} = \emptyset$ for $\{\alpha, \beta\} \in [\kappa^+]^2$ and $\xi \in \kappa$,
- (ii) $M_{\alpha,\xi} \cap M_{\alpha,\zeta} = \emptyset$ for $\alpha \in \kappa^+$ and $\{\xi, \zeta\} \in [\kappa]^2$, and
- (iii) $|M_{\alpha}^-| \leq \kappa$, where $M_{\alpha}^- = \kappa^+ \setminus \bigcup_{\zeta < \kappa} M_{\alpha,\zeta}$ for $\alpha < \kappa^+$.

Fix a partition $\{Y_\eta : \eta < \kappa^+\}$ witnessing that X is monotonically κ^+ -resolvable.

Let

$$Z_{\alpha,\zeta} = \bigcup \{Y_\eta : \eta \in M_{\alpha,\zeta}\}$$

for $\alpha < \kappa^+$ and $\zeta < \kappa$, and let

$$Z_\alpha = \bigcup_{\zeta < \kappa} Z_{\alpha, \zeta}.$$

Since $Z_\alpha = \bigcup \{Y_\eta : \eta \in \kappa^+ \setminus M_\alpha^-\}$, assumption (iii) implies that every Z_α is dense in X .

Case 1. There is $\alpha < \kappa^+$ such that for all $\zeta < \kappa$

$$\bigcup_{\zeta \leq \xi} Z_{\alpha, \zeta} \subset^{dense} Z_\alpha.$$

Then $(Z_{\alpha, \zeta})_{\zeta < \kappa}$ witnesses that Z_α is monotonically κ -resolvable and so by Corollary 3.2, X is also monotonically κ -resolvable.

Case 2. For all $\alpha < \kappa^+$ there is $\zeta_\alpha < \kappa$ and there is a non-empty open set $U_\alpha \in \tau_X$ such that

$$(\dagger) \quad \bigcup_{\zeta_\alpha \leq \xi} Z_{\alpha, \zeta} \cap U_\alpha = \emptyset.$$

Then there is a set $I \in [\kappa^+]^{\kappa^+}$ and there is an ordinal $\zeta < \kappa$ such that $\zeta_\alpha = \zeta$ for all $\alpha \in I$.

Fix an arbitrary $K \in [I]^\kappa$. By (iii) we can find $\rho < \kappa^+$ such that

$$\bigcup_{\alpha \in K} M_\alpha^- \subset \rho.$$

Let $Z = \bigcup_{\rho < \eta} Y_\eta$. Then $Z \subset^{dense} X$ and $Z \subset Z_\alpha$ for all $\alpha \in K$.

Claim. If $L \in [K]^\kappa$ then

$$\bigcap_{\alpha \in L} U_\alpha \cap Z = \emptyset.$$

PROOF OF THE CLAIM: Assume on the contrary that $z \in \bigcap_{\alpha \in L} U_\alpha \cap Z$. Then $z \in Y_\eta$ for some $\rho < \eta$.

Let $\alpha \in L$. Then $\eta \in \kappa^+ \setminus \rho \subset \bigcup_{\xi < \kappa} M_{\alpha, \xi}$. Pick $\xi_\alpha < \kappa$ with $\eta \in M_{\alpha, \xi_\alpha}$. Then $Y_\eta \subset Z_{\alpha, \xi_\alpha}$, so $Z_{\alpha, \xi_\alpha} \cap U_\alpha \neq \emptyset$, so $\xi_\alpha < \zeta_\alpha = \zeta$ by (\dagger) .

Since $\zeta < \kappa = |L|$, there are $\alpha \neq \beta \in [L]^2$ such that $\xi_\alpha = \xi_\beta$. Thus $\eta \in M_{\alpha, \xi_\alpha} \cap M_{\beta, \xi_\beta}$ which contradicts (i) because $\xi_\alpha = \xi_\beta$. \square

Fix an enumeration $K = \{\chi_\xi : \xi < \kappa\}$, and let $V_\zeta = \bigcup_{\zeta < \xi} U_{\chi_\xi}$. Then the sequence $\langle V_\zeta : \zeta < \kappa \rangle$ is decreasing and

$$\bigcap_{\zeta < \kappa} V_\zeta \cap Z = \emptyset$$

by the Claim.

Since X is κ -c.c. there is $\xi < \kappa$ such that $\overline{V_\zeta} = \overline{V_\xi}$ for all $\xi < \zeta < \kappa$.

We can assume that $\xi = 0$. Let

$$T_\zeta = \begin{cases} V_0 \setminus Z & \text{if } \zeta = 0, \\ ((\bigcap_{\xi < \zeta} V_\xi) \setminus V_\zeta) \cap Z & \text{if } \zeta > 0. \end{cases}$$

Then

$$\bigcup_{\xi < \zeta} T_\zeta \supset V_\xi \cap Z \subset^{dense} V.$$

Thus the partition $\{T_\zeta : \zeta < \kappa\}$ witnesses that V is monotonically κ -resolvable. □

PROOF OF THEOREM 1.3: Let $\mathcal{Y} = \{Y \in \tau_X : |Y| = \Delta(Y)\}$.

Then $\bigcup \mathcal{Y}$ is dense in X , and every open subset of every $Y \in \mathcal{Y}$ is also in \mathcal{Y} . Thus by Lemma 3.1 it is enough to prove that a c.c.c. space Y with $\omega_1 \leq |Y| = \Delta(Y) < \omega_\omega$ is monotonically ω_1 -resolvable.

Let $Y \in \mathcal{Y}$ such that $\omega_n = |Y|$. Clearly, Y is monotonically ω_n -resolvable because $|Y| = \Delta(Y) = \omega_n$. Since Y is c.c.c. then Y is ω_{n-1} -c.c. By Theorem 3.3, Y is monotonically ω_{n-1} -resolvable. By repeating the application of Theorem 3.3 $n - 2$ times we conclude that Y is monotonically ω_1 -resolvable. □

Problem 3.4. *Is it true that every crowded c.c.c. space with $\Delta(X) \geq \omega_1$ is monotonically ω_1 -resolvable?*

4. Spaces which are not monotonically ω_1 -resolvable

If X is a topological space, and $\mathcal{D} \subset \mathcal{P}(X)$, we write

$$\overline{\overline{\mathcal{D}}} = \{\overline{D} : D \in \mathcal{D}\}.$$

Lemma 4.1. *Let X be a topological space. Assume that $\overline{\overline{\mathcal{D}}}$ is point-countable for each point-countable family $\mathcal{D} \subset \mathcal{P}(X)$. Then X does not contain any monotonically ω_1 -resolvable subspace Y .*

PROOF: Assume that $\{Y_\zeta : \zeta < \omega_1\}$ is a partition of Y . Let $D_\xi = \bigcup \{Y_\zeta : \xi < \zeta\}$ for $\xi < \omega_1$. Then the family $\mathcal{D} = \{D_\xi : \xi < \omega_1\}$ is point-countable. Hence $\overline{\overline{\mathcal{D}}}$ is also point-countable. So D_ξ is not dense in Y for all but countably many ξ . Therefore the partition $\{Y_\zeta : \zeta < \omega_1\}$ does not witness that Y is monotonically ω_1 -resolvable. □

To prove Theorems 1.4 and 1.5 we recall some definitions and results from [6] and [5].

Definition 4.2 ([6, Definition 3.1]). Let κ be an infinite cardinal, and let \mathcal{F} be a filter on κ . Let T be the tree $\kappa^{<\omega}$. A topology $\tau_{\mathcal{F}}$ is defined on T by

$$\tau_{\mathcal{F}} = \{V \subset T : \forall t \in V \{\alpha \in \kappa : t \frown \alpha \in V\} \in \mathcal{F}\},$$

and the space $\langle T, \tau_{\mathcal{F}} \rangle$ is denoted by $X(\mathcal{F})$.

PROOF OF THEOREM 1.4: Let \mathcal{U} be a κ -complete non-principal ultrafilter on κ .

The space $X = X(\mathcal{U})$ is monotonically normal by [6, Theorem 3.1].

An ultrafilter \mathcal{U} is λ -*descendingly complete* if $\bigcap\{U_{\zeta} : \zeta < \lambda\} \neq \emptyset$ for each decreasing sequence $\{U_{\zeta} : \zeta < \lambda\} \subset \mathcal{U}$.

A σ -complete ultrafilter is clearly ω -descendingly-complete. In the proof of [6, Theorem 3.5] the authors prove Lemma 3.6 which claims that $\overline{\overline{\mathcal{D}}}$ is point-countable for each point-countable family $\mathcal{D} \subset \mathcal{P}(X(\mathcal{F}))$ provided that \mathcal{F} is a ω -descendingly complete ultrafilter. So $\overline{\overline{\mathcal{D}}}$ is point-countable for each point-countable family $\mathcal{D} \subset \mathcal{P}(X)$, and so X is not monotonically ω_1 -resolvable by Lemma 4.1. \square

Instead of Theorem 1.5 we prove the following theorem which is a slight improvement of [5, Theorem 5].

Theorem 4.3. *If it is consistent that there is a measurable cardinal, then it is also consistent that there is an ω -resolvable monotonically normal space X with $|X| = \Delta(X) = \omega_{\omega}$ such that if a family $\mathcal{D} \subset \mathcal{P}(X)$ is point-countable, then the family $\overline{\overline{\mathcal{D}}} = \{\overline{\overline{D}} : D \in \mathcal{D}\}$ is also point countable. Hence X does not contain any monotonically ω_1 -resolvable subspace.*

PROOF: In [5, p.665] the authors write that “starting from one measurable, Woodin ([15]) constructed a model in which \aleph_{ω} carries an ω_1 -descendingly complete uniform ultrafilter. Woodin’s model V_1 can be embedded into a bigger ZFC model V_2 so that the pair of models (V_1, V_2) with $\kappa = \aleph_{\omega}$ satisfies the two models situation”, i.e.

- (1) $\omega_1^{V_1} = \omega_1^{V_2}$,
- (2) there is a countable subset A of ω_{ω} in V_2 such that no $B \in V_1$ of cardinality $< \omega_{\omega}$ covers A ,
- (3) for the filter \mathcal{G} on ω_{ω} defined in V_2 by $B \in \mathcal{G}$ iff $A \setminus B$ is finite, we have $\mathcal{G} \cap V_1 \in V_1$.

(The “two model situation” is defined in [5, Theorem 4.5]).

Let $\mathcal{F} = \mathcal{G} \cap V_1$ and consider the space $X = X(\mathcal{F})$. As it was observed in [6], spaces obtained as $X(\mathcal{H})$ from some filter \mathcal{H} are monotonically normal and ω -resolvable.

In [5, Theorem 4.1] Juhász and Magidor showed that the space $X(\mathcal{F})$ is actually hereditarily ω_1 -irresolvable. They proved the following lemma:

Lemma 4.2 ([5]). *For any $D \subset X(F)$ and $t \in \overline{\overline{D}}$ there is a finite sequence s of members of A such that $t \frown s \in D$.*

Using this lemma we show that $\overline{\mathcal{D}}$ is point-countable for each point-countable family $\mathcal{D} \subset \mathcal{P}(X)$, and so X is not monotonically ω_1 -resolvable by Lemma 4.1.

Indeed, let $\mathcal{D} \subset \mathcal{P}(X)$ be an uncountable family such that $t \in \bigcap_{D \in \mathcal{D}} \overline{D}$. Then, by [5, Lemma 4.3], for each $D \in \mathcal{D}$ we can pick a finite sequence s_D of members of A such that $t \frown s_D \in D$. Since there are only countable many finite sequences of elements of A there is s such that $s_D = s$ for uncountably many $D \in \mathcal{D}$. Then $t \frown s$ is in uncountably many elements of \mathcal{D} , so \mathcal{D} is not point-countable.

We have thus proved that no subspace of X is monotonically ω_1 -resolvable. \square

5. ω -resolvability after adding a single Cohen real

Before proving Theorem 1.6 we need some preparation.

The notion of almost resolvability was introduced by Bolstein [2] in 1973: a topological space is *almost-resolvable* if it is a countable union of sets with empty interiors. The notion of monotone ω -resolvability was first considered in [13] under the name almost- ω -resolvability.

Clearly almost ω -resolvable (i.e. monotonically ω -resolvable) spaces are almost resolvable.

Lemma 5.1. *Let X be a crowded topological space.*

(1) *If X is monotonically ω -resolvable, then X is ω -resolvable in $V^{\text{Fn}(\omega, 2)}$.*

(2) *If X is resolvable in $V^{\text{Fn}(\omega, 2)}$, then X is almost-resolvable.*

PROOF: (1) Assume that the function $f : X \rightarrow \omega$ witnesses the monotone ω -resolvability of X .

If \mathcal{G} is the V -generic filter in $\text{Fn}(\omega, \omega)$, and $g = \bigcup \mathcal{G}$, then the function $h = g \circ f$ witnesses that X is ω -resolvable.

We need to show that $\{y \in X : (g \circ f)(y) = n\}$ is dense in X .

Indeed, let $p \in \text{Fn}(\omega, \omega)$ and $\emptyset \neq U \in \tau_X$. Since $f : X \rightarrow \omega$ witnesses the monotone ω -resolvability of X there is $y \in U$ such that

$$f(y) > \max \text{dom}(p).$$

Let

$$q = p \cup \{\langle f(y), n \rangle\}.$$

Then $q \leq p$ and

$$q \Vdash (g \circ f)(y) = n.$$

So we proved that $g \circ f$ witnesses that X is ω -resolvable in the generic extension.

(2) Assume that

$$V^{\text{Fn}(\omega, 2)} \models \text{“}X \text{ has a partition } \{D_0, D_1\} \text{ into dense subsets.”}$$

For all $p \in \text{Fn}(\omega, 2)$ and $i < 2$ let

$$D_i^p = \{x \in X : p \Vdash x \in \dot{D}_i\}.$$

Then $X = \bigcup \{D_i^p : p \in \text{Fn}(\omega, 2), i < 2\}$, and we claim that $\text{int } D_i^p = \emptyset$ for each $p \in \text{Fn}(\omega, 2)$, and $i < 2$.

Indeed, fix p and i and let U be an arbitrary non-empty open subset. Then $p \Vdash U \cap D_{1-i} \neq \emptyset$, so there is $q \leq p$ and $y \in U$ such that $q \Vdash y \in D_{1-i}$. Then $q \Vdash y \notin \dot{D}_i$, hence $p \nVdash y \in \dot{D}_i$, and so $y \notin D_i^p$. Thus $U \not\subset D_i^p$. Since U was arbitrary, we proved $\text{int } D_i^p = \emptyset$. \square

After this preparation we can prove Theorem 1.6.

PROOF OF THEOREM 1.6: (1) Kunen [7] proved that it is consistent, modulo a measurable cardinal, that there is a maximal independent family $\mathcal{A} \subset \mathcal{P}(\omega_1)$ which is also σ -independent.

In [9, Theorems 3.1 and 3.2] the authors proved that if there is a maximal independent family $\mathcal{A} \subset \mathcal{P}(\omega_1)$ which is also σ -independent, then there is a Baire space X with $|X| = \Delta(X) = \omega_1$ such that every open subspace of X is irresolvable, i.e. the space X is *OHI*.

It is well-known that a crowded OHI Baire space X is not almost resolvable: if $X = \bigcup_{n \in \omega} X_n$, then $\text{int } X_n \neq \emptyset$ for some $n \in \omega$.

Indeed, if $\text{int } X_n = \emptyset$, then $X \setminus X_n$ is dense, so $U_n = \text{int}(X \setminus X_n)$ is dense in X because every open subset of X is irresolvable. Thus $\bigcap_{n \in \omega} U_n \neq \emptyset$ because X is Baire. However

$$\bigcap_{n \in \omega} U_n \subset \bigcap_{n \in \omega} (X \setminus X_n) = X \setminus \bigcup_{n \in \omega} X_n = \emptyset,$$

which is a contradiction.

Thus X is not almost resolvable, so it is not ω -resolvable in the model $V^{\text{Fn}(\omega, 2)}$ by Lemma 5.1(2).

(2) In [10] the authors proved that if $V = L$, then there are no crowded Baire irresolvable spaces. Hence, by [13], if $V = L$, then every crowded space X is almost- ω -resolvable (i.e. monotonically ω -resolvable).

So these spaces are ω -resolvable in the model $V^{\text{Fn}(\omega, 2)}$ by Lemma 5.1(1).

(3) Let X be a crowded c.c.c. space.

We can assume that $|X| = \Delta(X)$.

By induction we define a strictly decreasing sequence of cardinals:

$$\kappa_0, \kappa_1, \dots, \kappa_n \dots$$

as follows.

- (i) $\kappa_0 = \Delta(X)$,
- (ii) if κ_i is singular, then $\kappa_{i+1} = \text{cf}(\kappa_i)$,
- (iii) if $\kappa_i > \omega$ is regular, then $\kappa_i = \lambda^+$ (because $|X|$ is below the first weakly inaccessible cardinal,) and let $\kappa_{i+1} = \lambda$,
- (iv) if $\kappa_i = \omega$ or $\kappa_i = \omega_1$, then we stop.

Assume that the construction stopped in the n th step.

Then we can prove, by finite induction, that X is monotonically κ_i -resolvable for all $i \leq n$ by Theorem 3.3. Thus X is monotonically ω -resolvable or monotonically ω_1 -resolvable, and so either X is ω -resolvable in $V^{\text{Fn}(\omega,2)}$ by Lemma 5.1(1), or X is ω_1 -resolvable in $V^{\text{Fn}(\omega_1,2)}$ by Theorem 2.1. \square

Problem 5.2 ([13, Questions 5.2.]). *Are almost resolvability and almost- ω -resolvability equivalent in the class of irresolvable spaces?*

Problem 5.3. *Is there, in ZFC, a crowded topological space X which is irresolvable in the Cohen generic extension $V^{\text{Fn}(\omega,2)}$?*

REFERENCES

- [1] Angoa J., Ibarra M., Tamariz-Mascarúa A., *On ω -resolvable and almost- ω -resolvable spaces*, Comment. Math. Univ. Carolin. **49** (2008), no. 3, 485–508.
- [2] Bolstein R., *Sets of points of discontinuity*, Proc. Amer. Math. Soc. **38** (1973), no. 1, 193–197.
- [3] Dorantes-Aldama A., *Baire irresolvable spaces with countable Souslin number*, Topology Appl. **188** (2015), 16–26.
- [4] Hewitt E., *A problem of set theoretic topology*, Duke Math. J. **10** (1943), 309–333.
- [5] Juhász I., Magidor M., *On the maximal resolvability of monotonically normal spaces*, Israel J. Math. **192** (2012), 637–666.
- [6] Juhász I., Soukup L., Szentmiklóssy Z., *Resolvability and monotone normality*, Israel J. Math. **166** (2008), 1–16.
- [7] Kunen K., *Maximal σ -independent families*, Fund. Math. **117** (1983), no. 1, 75–80.
- [8] Kunen K., Prikry K., *On descendingly incomplete ultrafilters*, J. Symbolic Logic **36** (1971), no. 4, 650–652.
- [9] Kunen K., Szymanski A., Tall F., *Naire irresolvable spaces and ideal theory*, Ann. Math. Sil. no. 14 (1986), 98–107.
- [10] Kunen K., Tall F., *On the consistency of the non-existence of Baire irresolvable spaces*, manuscript privately circulated, Topology Atlas, 1998, <http://at.yorku.ca/v/a/a/a/27.htm>
- [11] Pavlov O., *Problems on (ir)resolvability*, in Open Problems in Topology II, Elsevier, 2007, pp. 51–59.
- [12] Gruenhage G., *Generalized metrizable spaces*, in Recent Progress in General Topology, III, Springer Science & Business Media, 2013, pp. 471–505.
- [13] Tamariz-Mascarúa A., Villegas-Rodríguez H., *Spaces of continuous functions, box products and almost- ω -resolvable spaces*, Comment. Math. Univ. Carolin. **43** (2002), no. 4, 687–705.
- [14] Ulam S., *Zur Masstheorie in der allgemeinen Mengenlehre*, Fund. Math. **16** (1930), 140–150.
- [15] Woodin W.H., *Descendingly complete ultrafilter on \aleph_ω* , personal communication.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES,
HUNGARY

E-mail: soukup@renyi.hu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTHERN IOWA, USA

E-mail: adrienne.stanley@uni.edu

(Received May 23, 2017, revised August 29, 2017)