

Hölder continuity of bounded generalized solutions for some degenerated quasilinear elliptic equations with natural growth terms

SALVATORE BONAFEDE

Abstract. We prove the local Hölder continuity of bounded generalized solutions of the Dirichlet problem associated to the equation

$$\sum_{i=1}^m \frac{\partial}{\partial x_i} a_i(x, u, \nabla u) - c_0 |u|^{p-2} u = f(x, u, \nabla u),$$

assuming that the principal part of the equation satisfies the following degenerate ellipticity condition

$$\lambda(|u|) \sum_{i=1}^m a_i(x, u, \eta) \eta_i \geq \nu(x) |\eta|^p,$$

and the lower-order term f has a natural growth with respect to ∇u .

Keywords: elliptic equations; weight function; regularity of solutions

Classification: 35J15, 35J70, 35B65

1. Introduction

Consider the equation

$$(1.1) \quad \sum_{i=1}^m \frac{\partial}{\partial x_i} a_i(x, u, \nabla u) - c_0 |u|^{p-2} u = f(x, u, \nabla u) \quad \text{in } \Omega,$$

where Ω is a bounded open set of \mathbb{R}^m , $m \geq 2$, c_0 is a positive constant, ∇u is the gradient of unknown function u and f is a nonlinear function which has the growth of rate p , $1 < p < m$, with respect to the gradient ∇u . We shall suppose that the following degenerate ellipticity condition is satisfied:

$$(1.2) \quad \lambda(|u|) \sum_{i=1}^m a_i(x, u, \eta) \eta_i \geq \nu(x) |\eta|^p,$$

where $\eta = (\eta_1, \eta_2, \dots, \eta_m) \in \mathbb{R}^m$, $|\eta| = (\eta_1^2 + \dots + \eta_m^2)^{1/2}$, and $\nu: \Omega \rightarrow (0, \infty)$, $\lambda: [0, \infty) \rightarrow [1, \infty)$ are functions with properties to be specified later on.

In the present article, we prove the local Hölder continuity in Ω of every bounded generalized solution of equation (1.1) under the condition (1.2). We want to emphasize that the study of the quasilinear equations where the lower-order term has natural p -growth deserves special attention because to obtain the existence of the solutions, if $1 < p \leq m$, it is not possible to directly apply the standard theory of the pseudomonotone operators; moreover, the solution in general is unbounded. In this regard, we refer, for instance, to [1], [8], [20].

Existence and L^∞ -estimates of bounded solutions for quasilinear elliptic equations with natural growth of lower-order terms, in nondegenerate case, were established, for instance, in [2], [3], [7], [29], [30], [31], and the Hölder continuity on compact subsets of Ω of solutions was proved in [19, Chapter IX, Section 2], [32]. Similar results for elliptic equations and variational inequalities without the natural growth were obtained in [23], [25], [26] for the nondegenerate case, and also in [15]–[18], [22], [27], [28] for the degenerate case. We also mention the articles [10], [33] where the linear case is studied with weights belonging to Muckenhoupt's class.

Assuming the degenerate ellipticity condition (1.2), Drábek and Nicolosi in [9] obtained the existence of bounded generalized solutions of equation (1.1) establishing more general results than those obtained from Boccardo, Murat and Puel in [2], [3]. On the related topic and in degenerate-case, we also refer to [4]–[6] and [12], [13]. The results obtained in [9] are the starting point for this research.

The present paper is organized as follows. In Section 2 we formulate the hypotheses, we state our problem and the main results. Section 3 consists of preliminary lemmas which are sufficient in the proof of our main results. In Section 4 we prove local Hölder continuity of solutions of Dirichlet problem associated to equation (1.1). For the proof, we use an analogue of Moser's method (see [21]) proposed in [25] and modified in [32] for equations with natural growth terms. In Section 5 we give examples where all our assumptions are satisfied.

2. Hypotheses and formulation of the main results

We shall suppose that \mathbb{R}^m , $m \geq 2$, is the m -dimensional euclidean space with elements $x = (x_1, x_2, \dots, x_m)$. Let Ω be an open bounded nonempty subset of \mathbb{R}^m .

Let p be a real number such that $1 < p < m$.

Hypothesis 2.1. Let $\nu: \Omega \rightarrow (0, \infty)$ be a measurable function such that

$$\nu(x) \in L^1_{\text{loc}}(\Omega), \quad \left(\frac{1}{\nu(x)}\right)^{1/(p-1)} \in L^1_{\text{loc}}(\Omega).$$

We denote $W^{1,p}(\nu, \Omega)$ as the set of all functions $u \in L^p(\Omega)$ having for every $i = 1, \dots, m$ the weak derivative $\partial u / \partial x_i$ with the property $\nu |\partial u / \partial x_i|^p \in L^1(\Omega)$.

The space $W^{1,p}(\nu, \Omega)$ is a Banach space with respect to the norm

$$\|u\|_{1,p} = \left[\int_{\Omega} (|u|^p + \nu |\nabla u|^p) dx \right]^{1/p}.$$

The space $\mathring{W}^{1,p}(\nu, \Omega)$ is the closure of $C_0^\infty(\Omega)$ in $W^{1,p}(\nu, \Omega)$. Put $W = \mathring{W}^{1,p}(\nu, \Omega) \cap L^\infty(\Omega)$.

Hypothesis 2.2. We assume that

$$\frac{1}{\nu(x)} \in L^s(\Omega)$$

with $s > \max(1/(p-1), m/p)$.

We set $\tilde{p} = mp/(m-p+m/s)$. Then, we have $W^{1,p}(\nu, \Omega) \subset L^{\tilde{p}}(\Omega)$ and there exists $\hat{c} > 0$ depending only on m, p, s , and Ω such that for every $u \in \mathring{W}^{1,p}(\nu, \Omega)$

$$\left(\int_{\Omega} |u|^{\tilde{p}} dx \right)^{1/\tilde{p}} \leq \hat{c} \left(\int_{\text{supp}u} \left(\frac{1}{\nu} \right)^s dx \right)^{1/(ps)} \left(\int_{\Omega} \sum_{i=1}^m \nu \left| \frac{\partial u}{\partial x_i} \right|^p dx \right)^{1/p}.$$

In this connection see, for instance, [13] and [22].

Hypothesis 2.3. The functions $f(x, u, \eta)$, $a_i(x, u, \eta)$, $i = 1, 2, \dots, m$, are Carathéodory functions in $\Omega \times \mathbb{R} \times \mathbb{R}^m$, i.e., measurable with respect to x for every $(u, \eta) \in \mathbb{R} \times \mathbb{R}^m$ and continuous with respect to (u, η) for almost every $x \in \Omega$.

Hypothesis 2.4. There exist a number σ and a function $f^*(x)$ such that

$$\max \left(0, \frac{2-p}{2} \right) < \sigma < 1, \quad f^* \in L^1(\Omega)$$

and

$$(2.1) \quad |f(x, u, \eta)| \leq \lambda(|u|)[f^*(x) + |u|^{p-1+\sigma} + (\nu^{1/p}(x)|\eta|)^{p-1+\sigma} + \nu(x)|\eta|^p]$$

holds for almost every $x \in \Omega$ and for all real numbers $u, \eta_1, \eta_2, \dots, \eta_m$.

Hypothesis 2.5. There exist a nonnegative number $c_1 < c_0$ and a function $f_0(x) \in L^\infty(\Omega)$ such that for almost all $x \in \Omega$ and for all real numbers $u, \eta_1, \eta_2, \dots, \eta_m$ the inequality

$$(2.2) \quad u f(x, u, \eta) + c_1 |u|^p + \lambda(|u|)\nu(x)|\eta|^p + f_0(x) \geq 0$$

holds.

Hypothesis 2.6. There exists a function $a^* \in L^{p/(p-1)}(\Omega)$ such that for almost every $x \in \Omega$ and for any real numbers $u, \eta_1, \eta_2, \dots, \eta_m$ the inequality

$$(2.3) \quad \frac{|a_i(x, u, \eta)|}{\nu^{1/p}(x)} \leq \lambda(|u|)[a^*(x) + |u|^{p-1} + \nu^{(p-1)/p}(x)|\eta|^{p-1}]$$

holds.

Hypothesis 2.7. The condition (1.2) is satisfied for almost every $x \in \Omega$ and for all real numbers $u, \eta_1, \eta_2, \dots, \eta_m$; the function $\lambda = \lambda(z)$ is monotone and non-decreasing.

Hypothesis 2.8. For almost all $x \in \Omega$ and for every real $u, \eta_1, \eta_2, \dots, \eta_m, \tau_1, \tau_2, \dots, \tau_m$ the inequality

$$\sum_{i=1}^m [a_i(x, u, \eta) - a_i(x, u, \tau)](\eta_i - \tau_i) \geq 0$$

holds while the inequality holds if and only if $\eta \neq \tau$.

Assumptions 2.1–2.4, 2.6 and 2.7 provide the correctness of the following definition.

Definition. A generalized solution of equation (1.1) is a function $u \in W$ such that for every function $w \in W$,

$$(2.4) \quad \int_{\Omega} \left\{ \sum_{i=1}^m a_i(x, u, \nabla u) \frac{\partial w}{\partial x_i} + c_0 |u|^{p-2} u w + f(x, u, \nabla u) w \right\} dx = 0.$$

Note that if in addition to Assumptions 2.1–2.4, 2.6 and 2.7 they hold Assumptions 2.5 and 2.8, then there exists a generalized solution of equation (1.1). This follows from Theorem 2.1 of [9].

We will need the following further hypotheses on weight function.

Hypothesis 2.9. There exists a real number $\bar{t} > ms/(ps - m)$ such that

$$\nu(x) \in L^{\bar{t}}(\Omega).$$

For every $y \in \mathbb{R}^m$ and $R > 0$ we denote

$$B_R(y) = \{x \in \mathbb{R}^m : |x - y| < R\};$$

when not important, or clear from the context, we shall omit denoting the center as follows: $B_R = B_R(y)$.

Hypothesis 2.10. There exists $c > 0$ such that for every $y \in \Omega$ and $R > 0$ with $B_R(y) \subset \Omega$ the following inequality holds

$$\left\{ R^{-m} \int_{B_R(y)} \left(\frac{1}{\nu} \right)^s dx \right\}^{1/s} \left\{ R^{-m} \int_{B_R(y)} \nu^{\bar{t}} dx \right\}^{1/\bar{t}} \leq c.$$

As for Hypotheses 2.9 and 2.10 see, for example, [5]. Such kind of assumptions on weight function, introduced in [22], [27], [28], have been used in recent articles, see, for example, [16] and [18].

The main result of the present paper is a theorem on the local Hölder continuity of any generalized solution $u \in W$ of equation (1.1). More precisely, we prove the following

Theorem 2.11. *Assume that Hypotheses 2.1–2.4, 2.6, 2.7, 2.9 and 2.10 are satisfied with the functions $|a^*|^{p/(p-1)}$, f^* belonging to $L^\tau(\Omega)$ with $\tau > ms/(ps - m)$.*

Let $u \in W$ be a generalized solution of equation (1.1) and $M = \|u\|_{L^\infty(\Omega)}$.

Then there exist positive constants C and σ' such that for every open set Ω' , $\overline{\Omega'} \subset \Omega$ and every $x', x'' \in \Omega'$

$$|u(x') - u(x'')| \leq C[\text{dist}(\Omega', \partial\Omega)]^{-\sigma'} |x' - x''|^{\sigma'},$$

where $\sigma' = \sigma'(\mathbf{data}) < 1$, $C = C(\mathbf{data})$ and $\mathbf{data} = (m, p, \tau, \bar{t}, s, M, \lambda(z), \|f^*\|_{L^\tau(\Omega)}, \| |a^*|^{p/(p-1)} \|_{L^\tau(\Omega)}, \sigma, \text{meas } \Omega)$.

3. Auxiliary results

Lemma 3.1. *Let $f \in W^{1,q}(B_R)$, $q \geq 1$. Suppose there exist a measurable subset $G \subset B_R$ and positive constants C' and C'' such that*

$$\text{meas } G \geq C' R^m, \quad \max_G |f| \leq C''.$$

Then

$$\int_{B_R} |f|^q dx \leq CR^q \left(\sum_{i=1}^m \int_{B_R} \left| \frac{\partial f}{\partial x_i} \right|^q dx + R^{m-q} \right),$$

where C is a positive constant depending only on m, q, C', C'' .

The proof of this lemma is given in [24, Chapter 1, Section 2, Lemma 4].

The following lemma is due to John and Nirenberg (see [14], and see also [11, Theorem 7.21]).

Lemma 3.2. *Let $f \in W^{1,1}(\mathcal{O})$ where \mathcal{O} is a convex domain in \mathbb{R}^m . Suppose there exists a positive constant K such that*

$$\sum_{i=1}^m \int_{\mathcal{O} \cap B_\varrho} \left| \frac{\partial f}{\partial x_i} \right| dx \leq K \varrho^{m-1} \quad \text{for all balls } B_\varrho.$$

Then there exist positive constants σ_0 and C depending only on m such that

$$\int_{\mathcal{O}} \exp \left(\frac{\sigma}{K} |f - (f)_\mathcal{O}| \right) dx \leq C(\text{diam } \mathcal{O})^m,$$

where $\sigma = \sigma_0(\text{meas } \mathcal{O})(\text{diam } \mathcal{O})^{-m}$, $(f)_\mathcal{O} = \frac{1}{\text{meas } \mathcal{O}} \int_{\mathcal{O}} f dx$.

The following result is discussed in [11, Lemma 8.23] (see also [19, Lemma 4.8]).

Lemma 3.3. *Let ω be a non-decreasing function on an interval $(0, R_0]$ satisfying for all $R \leq R_0$ the inequality*

$$\omega(\vartheta R) \leq \theta \omega(R) + \varphi(R),$$

where φ is also non-decreasing function and $0 < \vartheta, \theta < 1$. Then for any $\delta \in (0, 1)$ and $R \leq R_0$ we have

$$\omega(R) \leq C \left(\left(\frac{R}{R_0} \right)^\epsilon \omega(R_0) + \varphi(R^\delta R_0^{1-\delta}) \right),$$

where $C = C(\vartheta, \theta)$ and $\epsilon = \epsilon(\vartheta, \theta, \delta)$ are positive constants.

4. Proof of Theorem 2.11

Suppose that Hypotheses 2.1–2.4, 2.6, 2.7, 2.9 and 2.10 hold with the functions $|a^*|^{p/(p-1)}$, $f^* \in L^\tau(\Omega)$, $\tau > m s / (ps - m)$. Let $u \in W$ be a generalized solution of equation (1.1). We set $M = \|u\|_\infty$, thus

$$(4.1) \quad |u| \leq M < \infty \quad \text{on } \Omega.$$

By d_i , $i = 1, 2, \dots$, we denote positive constants depending only on **data**.

Furthermore, let Ω' be an arbitrary open subset of Ω such that $\overline{\Omega'} \subset \Omega$ and $d = \text{dist}(\Omega', \partial\Omega)$. We fix $x_0 \in \Omega'$. For every $R \in (0, \min\{1, d/4\})$, we set

$$\omega_1(R) = \min_{B_R(x_0)} u, \quad \omega_2(R) = \max_{B_R(x_0)} u, \quad \omega(R) = \omega_2(R) - \omega_1(R).$$

Here the symbols min and max of course stands for *essential infimum* and *supremum*.

We fix a positive number r such that

$$(4.2) \quad r = 1 - \frac{m}{pt_\star} - \frac{m}{ps},$$

where $t_\star = \min(\tau, \bar{t})$.

For every $R \in (0, \min\{1, d/4\})$, we shall establish the inequality

$$(4.3) \quad \omega(R) \leq \alpha \omega(2R) + R^r$$

with a constant $\alpha \in (0, 1)$ depending only on **data**. This inequality and Lemma 3.3 imply the validity of Theorem 2.11.

To prove (4.3), we fix R such that $0 < R < \min\{1, d/4\}$. If $\omega(2R) < R^r$, then inequality (4.3) is evident. Therefore, we shall suppose that

$$(4.4) \quad \omega(2R) \geq R^r.$$

We shall also assume that

$$(4.5) \quad \text{meas } G(R) \geq \frac{1}{2} \text{meas } B_{3R/2}(x_0),$$

where $G(R) = \{x \in B_{3R/2}(x_0) : u(x) \leq (\omega_1(2R) + \omega_2(2R))/2\}$.

Let $F_1 : \Omega \rightarrow \mathbb{R}$ be the function such that

$$F_1 = \begin{cases} \frac{2e\omega(2R)}{\omega_2(2R) - u + R^r} & \text{in } B_{2R}(x_0), \\ e & \text{in } \Omega \setminus B_{2R}(x_0). \end{cases}$$

Due to (4.4) we have $F_1 \geq e$ in Ω .

Now, we need some integral estimates of solution u .

Lemma 4.1. *Let $B_\varrho \subset \Omega$ and let $\zeta \in C_0^\infty(\Omega)$ be a function such that*

$$(4.6) \quad \zeta = 0 \quad \text{in } \Omega \setminus B_\varrho \quad \text{and} \quad 0 \leq \zeta \leq 1.$$

Then there exist positive constants d_1, d_2 such that

$$(4.7) \quad \int_{B_\varrho} \nu |\nabla u|^p \zeta^p \, dx \leq d_1 \varrho^{m(\tau-1)/\tau} + d_2 \max_{B_\varrho} |\nabla \zeta|^p \left(\int_{B_\varrho} \nu^{\bar{\tau}} \, dx \right)^{1/\bar{\tau}} \varrho^{m(\bar{\tau}-1)/\bar{\tau}}.$$

PROOF: For every $x \in \Omega$ we set $v_1(x) = e^{\lambda_1 u(x)} \zeta^p(x)$ where

$$(4.8) \quad \lambda_1 = 3\lambda^2(M).$$

Simple calculations show that $v_1 \in \dot{W}^{1,p}(\nu, \Omega) \cap L^\infty(\Omega)$ and the following assertion holds:

(a) for every $i = 1, 2, \dots, m$,

$$\frac{\partial v_1}{\partial x_i} = \lambda_1 e^{\lambda_1 u} \zeta^p \frac{\partial u}{\partial x_i} + p e^{\lambda_1 u} \zeta^{p-1} \frac{\partial \zeta}{\partial x_i} \quad \text{a.e. in } \Omega.$$

Since $v_1 \in \dot{W}^{1,p}(\nu, \Omega) \cap L^\infty(\Omega)$, by virtue of (2.4), we have

$$\int_{\Omega} \left\{ \sum_{i=1}^m a_i(x, u, \nabla u) \frac{\partial v_1}{\partial x_i} + c_0 |u|^{p-2} u v_1 + f(x, u, \nabla u) v_1 \right\} dx = 0.$$

From this equality, using (1.2), (2.1), (4.8) and assertion (a), we deduce that

$$(4.9) \quad \lambda(M) \int_{B_\varrho} \nu |\nabla u|^p e^{\lambda_1 u} \zeta^p \, dx \leq I_\varrho + e^{\lambda_1 M} \int_{B_\varrho} g^*(x) \, dx,$$

where

$$I_\varrho = p \sum_{i=1}^m \int_{B_\varrho} |a_i(x, u, \nabla u)| \left| \frac{\partial \zeta}{\partial x_i} \right| e^{\lambda_1 u} \zeta^{p-1} dx,$$

$$g^*(x) = c_0 M^{p-1} + 2\lambda(M)[|f^*(x)| + M^{p-1+\sigma} + 1].$$

Let us obtain suitable estimate for the first addend in the right-hand side of (4.9). *Estimate of I_ϱ .* Using the Young's inequality with the exponents $p/(p-1)$ and p , (2.3), (4.1) and (4.6), we obtain

$$(4.10) \quad I_\varrho \leq \frac{\lambda(M)}{2} \int_{B_\varrho} \nu |\nabla u|^p e^{\lambda_1 u} \zeta^p dx$$

$$+ d_3 e^{\lambda_1 M} \int_{B_\varrho} g_1(x) dx + d_4 e^{\lambda_1 M} \int_{B_\varrho} \nu |\nabla \zeta|^p dx,$$

where

$$g_1(x) = 4^{1/(p-1)} (\lambda(M))^{p/(p-1)} [|a^*(x)|^{p/(p-1)} + M^p].$$

From (4.9), (4.10) it follows that

$$(4.11) \quad \frac{\lambda(M)}{2} \int_{B_\varrho} \nu |\nabla u|^p e^{\lambda_1 u} \zeta^p dx \leq d_5 \int_{B_\varrho} (g^* + g_1) dx + d_6 \int_{B_\varrho} \nu |\nabla \zeta|^p dx.$$

By Hölder's inequality and the inequality $\tau > ms/(ps - m)$ we have

$$\int_{B_\varrho} (g^* + g_1) dx \leq \|g^* + g_1\|_\tau |B_\varrho|^{(\tau-1)/\tau} \leq d_7 \varrho^{m(\tau-1)/\tau}.$$

Moreover

$$\int_{B_\varrho} \nu |\nabla \zeta|^p dx \leq d_8 \max_{B_\varrho} |\nabla \zeta|^p \varrho^{m(\bar{t}-1)/\bar{t}} \left(\int_{B_\varrho} \nu^{\bar{t}} dx \right)^{1/\bar{t}}.$$

The last two inequalities and (4.11) imply inequality (4.7).

The lemma is proved. \square

Lemma 4.2. *Let $B_\varrho \subset B_{2R}(x_0)$ and let $\zeta \in C_0^\infty(\Omega)$ be a function such that condition (4.6) is satisfied. Then there exist positive constants d_9, d_{10}, d_{11} such that*

$$(4.12) \quad \int_{B_\varrho} \frac{\nu |\nabla u|^p \zeta^p dx}{(\omega_2(2R) - u + R^r)^p} \leq d_9 \varrho^{m-p+m/s} + d_{10} \max_{B_\varrho} |\nabla \zeta|^p \left(\int_{B_\varrho} \nu^{\bar{t}} dx \right)^{1/\bar{t}}$$

$$\times \varrho^{m(\bar{t}-1)/\bar{t}} + d_{11} \varrho^{m(\tau-1)/\tau}.$$

PROOF: For every $x \in B_{2R}(x_0)$, we set $U(x) = \omega_2(2R) - u(x) + R^r$,

$$v_2(x) = \begin{cases} \zeta^p(x)[U(x)]^{1-p} & \text{if } x \in B_{2R}(x_0), \\ 0 & \text{if } x \in \Omega \setminus B_{2R}(x_0). \end{cases}$$

Simple calculations show that

$$v_2 \in \mathring{W}^{1,p}(\nu, \Omega) \cap L^\infty(\Omega)$$

and the following assertion holds:

(b) for every $i = 1, 2, \dots, m$

$$\frac{\partial v_2}{\partial x_i} = pU^{1-p}\zeta^{p-1}\frac{\partial \zeta}{\partial x_i} + (p-1)U^{-p}\zeta^p\frac{\partial u}{\partial x_i} \quad \text{a.e. in } B_{2R}.$$

Putting the function v_2 into (2.4) instead of w and using (1.2), (2.1), (4.1) and assertion (b), we obtain

$$(4.13) \quad \frac{1}{\lambda(M)} \int_{B_\varrho} \nu |\nabla u|^p U^{-p} \zeta^p \, dx \leq I_{2,\varrho} + I_{3,\varrho} + \frac{(2M+1)}{p-1} \int_{B_\varrho} g^* U^{-p} \, dx,$$

where

$$I_{2,\varrho} = p \sum_{i=1}^m \int_{B_\varrho} |a_i(x, u, \nabla u)| \left| \frac{\partial \zeta}{\partial x_i} \right| U^{1-p} \zeta^{p-1} \, dx,$$

$$I_{3,\varrho} = \frac{2\lambda(M)}{p-1} \int_{B_\varrho} \nu |\nabla u|^p U^{1-p} \zeta^p \, dx.$$

Let us obtain suitable estimates for $I_{2,\varrho}$, $I_{3,\varrho}$.

Estimate of $I_{2,\varrho}$. Using the Young's inequality with the exponents $p/(p-1)$ and p , (2.3), (4.1) and (4.6), we obtain

$$(4.14) \quad I_{2,\varrho} \leq \frac{1}{4\lambda(M)} \int_{B_\varrho} \nu |\nabla u|^p U^{-p} \zeta^p \, dx$$

$$+ d_{12} \int_{B_\varrho} g_1 U^{-p} \, dx + d_{13} \max_{B_\varrho} |\nabla \zeta|^p \left(\int_{B_\varrho} \nu^{\bar{t}} \, dx \right)^{1/\bar{t}} \varrho^{m(\bar{t}-1)/\bar{t}}.$$

Estimate of $I_{3,\varrho}$. We use (4.1), the Young's inequality, (4.6) and (4.7) to obtain

$$(4.15) \quad I_{3,\varrho} \leq \frac{1}{4\lambda(M)} \int_{B_\varrho} \nu |\nabla u|^p U^{-p} \zeta^p \, dx$$

$$+ d_{14} \varrho^{m(\tau-1)/\tau} + d_{15} \max_{B_\varrho} |\nabla \zeta|^p \left(\int_{B_\varrho} \nu^{\bar{t}} \, dx \right)^{1/\bar{t}} \varrho^{m(\bar{t}-1)/\bar{t}}.$$

Collecting (4.13), (4.14) and (4.15), we get

$$(4.16) \quad \frac{1}{2\lambda(M)} \int_{B_\varrho} \nu |\nabla u|^p U^{-p} \zeta^p \, dx \leq d_{14} \varrho^{m(\tau-1)/\tau} \\ + d_{16} \max_{B_\varrho} |\nabla \zeta|^p \left(\int_{B_\varrho} \nu^{\bar{\tau}} \, dx \right)^{1/\bar{\tau}} \varrho^{m(\bar{\tau}-1)/\bar{\tau}} + d_{17} \int_{B_\varrho} g U^{-p} \, dx$$

where $g = g^* + g_1$.

By Hölder's inequality, $U \geq R^r$ and $\varrho/2 < R < 1$, we have

$$\int_{B_\varrho} g U^{-p} \, dx \leq \|g\|_\tau \left(\int_{B_\varrho} U^{-p\tau/(\tau-1)} \, dx \right)^{(\tau-1)/\tau} \leq d_{18} \left(\frac{2}{\varrho} \right)^{r\tau} \varrho^{m(\tau-1)/\tau}.$$

From last inequality, taking into account relation (4.2), we get

$$(4.17) \quad \int_{B_\varrho} g U^{-p} \, dx \leq d_{19} \varrho^{m-p+m/s}.$$

Inequalities (4.16) and (4.17) imply inequality (4.12). The lemma is proved. \square

Define in Ω the function $v_0 = \ln F_1$.

Let us prove that v_0 satisfies some integral inequalities.

Lemma 4.3. *Let $r_0 = sp/(s+1)$. Then, there exist positive constants $d_{20}, d_{21}, d_{22}, d_{23}$ such that*

$$(4.18) \quad \int_{B_{3R/2}(x_0)} v_0^{r_0} \, dx \leq d_{20} R^m + R^{r_0} \left\{ \left[d_{21} \left\| \frac{1}{\nu} \right\|_{L^s(\Omega)} + d_{22} \right] R^{m-p+m/s} \right. \\ \left. + d_{23} \left\| \frac{1}{\nu} \right\|_{L^s(\Omega)} R^{m(\tau-1)/\tau} \right\}^{s/(s+1)}.$$

PROOF: We choose a function $\zeta_1 \in C_0^\infty(\Omega)$ such that

$$0 \leq \zeta_1 \leq 1 \quad \text{in } \Omega, \quad \zeta_1 = 1 \quad \text{in } B_{3R/2}(x_0), \quad \zeta_1 = 0 \quad \text{in } \Omega \setminus B_{7R/4}(x_0), \\ \left| \frac{\partial \zeta_1}{\partial x_i} \right| \leq K_1 R^{-1} \quad \text{for } i = 1, 2, \dots, m,$$

where K_1 is an absolute constant, not depending on R . By definition of v_0 it results $1 \leq v_0 \leq 1 + \ln 4$ on $G(R)$; moreover from (4.5) it follows that $\text{meas } G(R) \geq d_{24} R^m$.

Hence from Lemma 3.1 we find that:

$$(4.19) \quad \int_{B_{3R/2}(x_0)} v_0^{r_0} \, dx \leq d_{25} R^m + d_{25} R \int_{B_{3R/2}(x_0)} \left\{ \sum_{i=1}^m \left| \frac{\partial u}{\partial x_i} \right| \right\} U^{-1} v_0^{r_0-1} \, dx.$$

By means of Young inequality we get

$$(4.20) \quad d_{25}R \int_{B_{3R/2}(x_0)} \left\{ \sum_{i=1}^m \left| \frac{\partial u}{\partial x_i} \right| \right\} U^{-1} v_0^{r_0-1} dx \\ \leq \frac{1}{r_0} (d_{25}R)^{r_0} \int_{B_{3R/2}(x_0)} \left\{ \sum_{i=1}^m \left| \frac{\partial u}{\partial x_i} \right| \right\}^{r_0} U^{-r_0} dx + \frac{r_0-1}{r_0} \int_{B_{3R/2}(x_0)} v_0^{r_0} dx.$$

From (4.19) and (4.20) we have:

$$(4.21) \quad \int_{B_{3R/2}(x_0)} v_0^{r_0} dx \leq r_0 d_{26} R^m + d_{27} R^{r_0} \int_{B_{3R/2}(x_0)} \sum_{i=1}^m \left| \frac{\partial u}{\partial x_i} \right|^{r_0} U^{-r_0} dx.$$

Using Hölder inequality with p/r_0 and $(1 - r_0/p)^{-1} = (s+1)$ and the definition of the function ζ_1 we obtain

$$\int_{B_{3R/2}(x_0)} \sum_{i=1}^m \left| \frac{\partial u}{\partial x_i} \right|^{r_0} U^{-r_0} dx \leq d_{28} \left(\int_{B_{7R/4}(x_0)} \left(\frac{1}{\nu} \right)^s dx \right)^{1/(s+1)} \\ \times \left(\int_{B_{7R/4}(x_0)} \nu |\nabla u|^p \zeta_1^p U^{-p} dx \right)^{s/(s+1)}.$$

Finally, we acquire inequality (4.18) from (4.21) estimating the last integral of previous inequality by Lemma 4.2 and taking into account that from the Hypothesis 2.10 we have

$$\left\{ \int_{B_{7R/4}(x_0)} \left(\frac{1}{\nu} \right)^s dx \right\}^{1/s} \left\{ \int_{B_{7R/4}(x_0)} \nu^{\bar{t}} dx \right\}^{1/\bar{t}} \leq d_{29} R^{m(1/s+1/\bar{t})}.$$

The lemma is proved. \square

Lemma 4.4. *For every $\kappa \geq 1$ there is a positive constant $c = c(\mathbf{data}, \kappa)$ such that $\lim_{\kappa \rightarrow \infty} c(\mathbf{data}, \kappa) = \infty$ and*

$$(4.22) \quad \int_{B_{3R/2}(x_0)} v_0^\kappa dx \leq c R^m.$$

PROOF: At first, we estimate from above the integral average

$$(v_0)_{B_{3R/2}(x_0)} = \frac{1}{\text{meas } B_{3R/2}(x_0)} \int_{B_{3R/2}(x_0)} v_0 dx$$

by a constant depending only on \mathbf{data} .

Using Hölder's inequality and Lemma 4.3 we get

$$(4.23) \quad \begin{aligned} (v_0)_{B_{3R/2}(x_0)} &\leq d_{30}R^{-m/r_0} \left(\int_{B_{3R/2}(x_0)} v_0^{r_0} dx \right)^{1/r_0} \\ &\leq d_{30}R^{-m/r_0} [d_{31}R^{m/r_0} + d_{32}R(d_{33}R^{m-p+m/s} \\ &\quad + d_{34}R^{m(\tau-1)/\tau})^{1/p}] \leq d_{35}. \end{aligned}$$

Next, let $B_{2\varrho} \subset B_{2R}(x_0)$, and let $\zeta_2 \in C_0^\infty(\Omega)$ be a function such that

$$\begin{aligned} 0 \leq \zeta_2 \leq 1 \quad \text{in } \Omega, \quad \zeta_2 = 1 \quad \text{in } B_\varrho, \quad \zeta_2 = 0 \quad \text{in } \Omega \setminus B_{2\varrho}, \\ \left| \frac{\partial \zeta_2}{\partial x_i} \right| \leq K_2 \varrho^{-1} \quad \text{for } i = 1, 2, \dots, m, \end{aligned}$$

where K_2 is an absolute constant, not depending on ϱ . Using Hölder's inequality, Lemma 4.2, Hypothesis 2.10, the properties of the function ζ_2 and that $\tau > ms/(ps - m)$, $s > 1/(p - 1)$, we derive that

$$\begin{aligned} \sum_{i=1}^m \int_{B_\varrho} \left| \frac{\partial v_0}{\partial x_i} \right| dx &\leq d_{36} \left(\int_{B_\varrho} \nu^{-1/(p-1)} dx \right)^{(p-1)/p} \\ &\quad \times \left(\int_{B_{2\varrho}} \nu |\nabla u|^p \zeta_2^p U^{-p} dx \right)^{1/p} \leq d_{37} \varrho^{m-1}. \end{aligned}$$

Hence, by Lemma 3.2, we have

$$(4.24) \quad \int_{B_{3R/2}(x_0)} \exp(d_{38} |v_0 - (v_0)_{B_{3R/2}}|) dx \leq d_{39} R^m.$$

Now let $\kappa \geq 1$. Then inequalities (4.23) and (4.24) imply (4.22).

The lemma is proved. □

Lemma 4.5. *There is a positive constant $c_3 = c_3(\mathbf{data})$ such that*

$$(4.25) \quad \|v_0\|_{L^\infty(B_R(x_0))} \leq c_3.$$

PROOF: We proceed the proof in four steps.

Step 1. We fix a function $\psi_0 \in C_0^\infty(\mathbb{R})$ such that

$$0 \leq \psi_0 \leq 1 \quad \text{on } \mathbb{R}, \quad \psi = 1 \quad \text{in } [-1, 1], \quad \psi = 0 \quad \text{in } \mathbb{R} \setminus (-3/2, 3/2).$$

For any $x \in \Omega$ we set $\psi(x) = \psi_0(|x - x_0|/R)$,

$$\tilde{v}(x) = \begin{cases} [v_0(x)]^k \psi^t(x) [U(x)]^{1-p} & \text{if } x \in B_{2R}(x_0), \\ 0 & \text{if } x \in \Omega \setminus B_{2R}(x_0), \end{cases}$$

where $U = \omega_2(2R) - u + R^r$,

$$(4.26) \quad k \geq \bar{k} := \max\{p, 2(6M+1)\lambda^2(M)\},$$

$$(4.27) \quad t > p.$$

Simple calculations show that

$$\tilde{v} \in \mathring{W}^{1,p}(\nu, \Omega) \cap L^\infty(\Omega)$$

and the following assertion holds:

(c) for every $i = 1, 2, \dots, m$,

$$\left| \frac{\partial \tilde{v}}{\partial x_i} - z \frac{\partial u}{\partial x_i} \right| \leq \frac{d_{40} t v_0^k \psi^{t-1}}{R U^{p-1}} \quad \text{a.e. in } \Omega,$$

where $z = [(p-1)v_0^k + k v_0^{k-1}] \psi^t U^{-p}$ and $d_{40} > 0$ depends only on $\max_{\mathbb{R}} |\psi'_i|$.

Putting the function \tilde{v} into (2.4) instead of w and using (1.2), (2.1), and assertion (c), we obtain

$$(4.28) \quad \begin{aligned} & \frac{(p-1)}{\lambda(M)} \int_{B_{2R}(x_0)} \nu |\nabla u|^p U^{-p} v_0^k \psi^t \, dx \\ & \quad + \frac{k}{\lambda(M)} \int_{B_{2R}(x_0)} \nu |\nabla u|^p U^{-p} v_0^{k-1} \psi^t \, dx \\ & \leq 2\lambda(M) \int_{B_{2R}(x_0)} \nu |\nabla u|^p U^{1-p} v_0^k \psi^t \, dx \\ & \quad + \int_{B_{2R}(x_0)} g^* v_0^k \psi^t U^{1-p} \, dx + \mathcal{I}, \end{aligned}$$

where $g^*(x)$ was defined in Lemma 4.1 and

$$(4.29) \quad \mathcal{I} = \frac{d_{40} t}{R} \sum_{i=1}^m \int_{B_{2R}(x_0)} |a_i(x, u, \nabla u)| U^{1-p} v_0^k \psi^{t-1} \, dx.$$

Step 2. We show that the first term in the right-hand side of inequality (4.28) is absorbed by the second term in its left-hand side. For this we need the inequality

$$(4.30) \quad U v_0 \leq 6M + 1 \quad \text{a.e. in } B_{2R}(x_0).$$

To prove it, we consider the function

$$\chi(s) = (s + R^r) \ln \frac{2\omega(2R)}{s + R^r}, \quad s \in [0, \omega(2R)].$$

According to (4.4) and to the elementary inequality $\ln b < b$, $b > 0$, we obtain that for every $s \in [0, \omega(2R)]$

$$0 \leq \chi(s) \leq 2\omega(2R) \leq 4M.$$

Now inequality (4.30) follows from the relations $R \leq 1$ and

$$Uv_0 = \omega_2(2R) - u + R^r + \chi(\omega_2(2R) - u) \quad \text{a.e. in } B_{2R}(x_0).$$

Using (4.30), the first term on the right-hand side of inequality (4.28) is estimated in the following way

$$(4.31) \quad 2\lambda(M) \int_{B_{2R}(x_0)} \nu |\nabla u|^p U^{1-p} v_0^k \psi^t \, dx \\ \leq 2(6M + 1)\lambda(M) \int_{B_{2R}(x_0)} \nu |\nabla u|^p U^{-p} v_0^{k-1} \psi^t \, dx.$$

Now (4.26), (4.28) and (4.31) imply the inequality

$$(4.32) \quad \frac{(p-1)}{\lambda(M)} \int_{B_{2R}(x_0)} \nu |\nabla u|^p U^{-p} v_0^k \psi^t \, dx \\ \leq (2M + 1) \int_{B_{2R}(x_0)} g^* v_0^k \psi^t U^{-p} \, dx + \mathcal{I}.$$

Step 3. Let us estimate from above the quantity \mathcal{I} , which is defined by (4.29). We use (2.3) and Young's inequality

$$|yz| \leq \varepsilon |y|^{p/(p-1)} + \varepsilon^{1-p} |z|^p,$$

where

$$y = |a_i(x, u, \nabla u)| U^{1-p} \psi^{(p-1)t/p} (\nu(x))^{-1/p}, \quad i = 1, 2, \dots, m, \\ z = t \psi^{(t-p)/p} \nu(x)^{1/p} / R,$$

and ε is an appropriate positive number, to obtain

$$(4.33) \quad \mathcal{I} \leq \frac{(p-1)}{2\lambda(M)} \int_{B_{2R}(x_0)} \nu |\nabla u|^p U^{-p} v_0^k \psi^t \, dx \\ + d_{41} \int_{B_{2R}(x_0)} g_1 v_0^k \psi^t U^{-p} \, dx + \frac{d_{42} t^p}{R^p} \int_{B_{2R}(x_0)} \nu v_0^k \psi^{t-p} \, dx,$$

where $g_1(x)$ was defined in Lemma 4.1.

From (4.32) and (4.33), for every $k \geq \bar{k}$ and $t > p$, it follows that

$$(4.34) \quad \int_{B_{2R}(x_0)} \nu |\nabla u|^p U^{-p} v_0^k \psi^t \, dx \leq d_{43} t^p \int_{B_{2R}(x_0)} \psi_1 v_0^k \psi^{t-p} \, dx,$$

where $\psi_1(x) = R^{-rp}[g_1(x) + g^*(x)] + \nu(x)R^{-p}$.

It results

$$(4.35) \quad \left(\int_{B_{2R}(x_0)} \psi_1^{t_*} \, dx \right)^{1/t_*} \leq R^{-p} \left(\int_{B_{2R}(x_0)} \nu^{t_*} \, dx \right)^{1/t_*} + R^{-rp} \|g_1 + g^*\|_{L^{t_*}(\Omega)}.$$

Now, we fix arbitrary $k \geq \bar{k}\tilde{p}/p$ and $t > \tilde{p}$ and let

$$z_1 = v_0^{k/\tilde{p}} \psi^{t/\tilde{p}}.$$

We have $z_1 \in \mathring{W}^{1,p}(\nu, \Omega)$ and for every $i = 1, 2, \dots, m$,

$$\begin{aligned} \int_{\Omega} \nu \left| \frac{\partial z_1}{\partial x_i} \right|^p \, dx &\leq d_{44} k^p \int_{B_{2R}(x_0)} \nu \left| \frac{\partial u}{\partial x_i} \right|^p v_0^{kp/\tilde{p}} U^{-p} \psi^{t\tilde{p}/\tilde{p}} \, dx \\ &\quad + d_{45} t^p \int_{B_{2R}(x_0)} \psi_1 v_0^{kp/\tilde{p}} \psi^{t\tilde{p}/\tilde{p}-p} \, dx. \end{aligned}$$

From last inequality and (4.34) we obtain

$$\int_{\Omega} \nu \left| \frac{\partial z_1}{\partial x_i} \right|^p \, dx \leq d_{46} k^p t^p \int_{B_{2R}(x_0)} \psi_1 v_0^{kp/\tilde{p}} \psi^{(t/\tilde{p}-1)p} \, dx.$$

Estimating integral to second term of last inequality by Hölder's inequality with the exponents t_* and $t_*/(t_* - 1)$, we obtain that for every $k \geq \bar{k}\tilde{p}/p$ and $t > \tilde{p}$ the following inequality holds:

$$(4.36) \quad \begin{aligned} \int_{\Omega} \nu \left| \frac{\partial z_1}{\partial x_i} \right|^p \, dx &\leq d_{46} k^p t^p \left(\int_{B_{2R}(x_0)} \psi_1^{t_*} \, dx \right)^{1/t_*} \\ &\quad \times \left(\int_{B_{2R}(x_0)} v_0^{kpt_*/(\tilde{p}(t_*-1))} \psi^{tpt_*/(\tilde{p}(t_*-1))-pt_*/(t_*-1)} \, dx \right)^{(t_*-1)/t_*}. \end{aligned}$$

Step 4. We set

$$\begin{aligned} H(k, t) &= \int_{B_{2R}(x_0)} v_0^k \psi^t \, dx, \quad k \in \mathbb{R}, \, t > 0, \\ \theta &= \frac{\tilde{p}(t_* - 1)}{pt_*}, \quad \tilde{m} = \frac{pt_*}{t_* - 1}. \end{aligned}$$

Due to Hypothesis 2.2. we have

$$(4.37) \quad H(k, t) \leq \tilde{c}^{\tilde{p}} \left[\int_{B_{2R}(x_0)} \left(\frac{1}{\nu} \right)^s dx \right]^{\tilde{p}/(ps)} \left[\int_{\Omega} \sum_{i=1}^m \nu \left| \frac{\partial z_1}{\partial x_i} \right|^p dx \right]^{\tilde{p}/p}.$$

From (4.36) and (4.37) it follows that

$$(4.38) \quad H(k, t) \leq d_{47} (k+t)^{2\tilde{p}} \left\{ \left[\int_{B_{2R}(x_0)} \left(\frac{1}{\nu} \right)^s dx \right]^{1/s} \left[\int_{B_{2R}(x_0)} \psi_1^{t_*} dx \right]^{1/t_*} \right\}^{\tilde{p}/p} \\ \times \left[H\left(\frac{k}{\theta}, \frac{t}{\theta} - \tilde{m} \right) \right]^{\theta}.$$

Using (4.35), (4.2) and Hypothesis 2.10, we obtain

$$(4.39) \quad \left[\int_{B_{2R}(x_0)} \left(\frac{1}{\nu} \right)^s dx \right]^{1/s} \left[\int_{B_{2R}(x_0)} \psi_1^{t_*} dx \right]^{1/t_*} \leq d_{48} R^{-p+m/s+m/t_*}.$$

Note that due to the definition of \tilde{p} and θ we have

$$(4.40) \quad \left(p - \frac{m}{s} - \frac{m}{t_*} \right) \frac{\tilde{p}}{p} = m(\theta - 1).$$

From (4.38), (4.39) and (4.40) we get

$$(4.41) \quad H(k, t) \leq d_{49} (k+t)^{2\tilde{p}} R^{-m(\theta-1)} \left[H\left(\frac{k}{\theta}, \frac{t}{\theta} - \tilde{m} \right) \right]^{\theta}$$

for every $k \geq \bar{k}\tilde{p}/p$ and $t > \tilde{p}$.

We choose a number $i_0 \in \mathbb{N}$ such that $\theta^{i_0} > \bar{k}\tilde{p}/p$ and set

$$k_i = \theta^{i_0+i}, \quad t_i = \frac{\tilde{m}\theta}{\theta-1} (\theta^{i_0+i} - 1), \quad i = 0, 1, 2, \dots$$

Then (4.41) and the inequality $\theta > 1$ imply that for every $i = 1, 2, \dots$,

$$[H(k_i, t_i)]^{1/k_i} \leq [d_{50} R^{-m} H(k_0, t_0)]^{1/\theta^{i_0}}.$$

By Lemma 4.4 we have

$$H(k_0, t_0) \leq \int_{B_{3R/2}(x_0)} v_0^{\theta^{i_0}} dx \leq d_{51} R^m.$$

From the last two inequalities it follows that

$$\|v_0\|_{L^\infty(B_R(x_0))} = \lim_{i \rightarrow \infty} \left(\int_{B_R(x_0)} v_0^{k_i} dx \right)^{1/k_i} \leq \limsup_{i \rightarrow \infty} [H(k_i, t_i)]^{1/k_i} \leq c_3.$$

The lemma is proved. \square

Inequality (4.25) implies (4.3). Recall that we proved (4.3) under assumption (4.5). If (4.5) is not true, we take instead of F_1 the function $F_2 = 2e\omega(2R) \times (u - \omega_1(2R) + R^r)^{-1}$ in $B_{2R}(x_0)$, and $F_2 = e$ in $\Omega \setminus B_{2R}(x_0)$, and arguing as above, we establish (4.3) again. Hence according to Lemma 3.3, the assertions of Theorem 2.11 are true.

Remark. The technique used to prove the local Hölder continuity for bounded generalized solutions of the Dirichlet problem associated with equation (1.1), assuming the degenerate ellipticity condition (1.2), can be repeated for bounded generalized solutions of equation (1.1) with the following boundary Neumann condition:

$$\sum_{i=1}^m a_i(x, u, \nabla u) \cos(\vec{n}, x_i) + c_2 |u|^{p-2} u + F(x, u) = 0, \quad c_2 > 0, \quad x \in \partial\Omega,$$

where $\partial\Omega$ is locally Lipschitz boundary and $\vec{n} = \vec{n}(x)$ is the outwardly directed (relative to Ω) unit vector normal to $\partial\Omega$ at every point $x \in \partial\Omega$ (see [6] for the Existence theorem in such case). So, we can prove that every function $u \in W^{1,p}(\nu, \Omega) \cap L^\infty(\Omega)$ satisfying

$$\int_{\Omega} \left\{ \sum_{i=1}^m a_i(x, u, \nabla u) \frac{\partial w}{\partial x_i} + c_0 |u|^{p-2} u w + f(x, u, \nabla u) w \right\} dx + \int_{\partial\Omega} \{c_2 |u|^{p-2} u w + F(x, u) w\} ds = 0$$

for any $w \in W^{1,p}(\nu, \Omega) \cap L^\infty(\Omega)$, it is locally Hölder continuous in Ω .

5. Examples

Example 5.1. Let $\Omega \subset \mathbb{R}^m$ be an open bounded set. Suppose for simplicity that $0 \in \partial\Omega$ and, additionally, we assume that

$$p > \frac{m}{2}, \quad m \geq 4.$$

Let $0 < \gamma < (m/2)(p - m/2)(3m/2 - p)^{-1}$, and let $\nu: \Omega \rightarrow (0, \infty)$ be defined by

$$\nu(x) = |x|^\gamma.$$

Let s be such that

$$\frac{m}{p - m/2} < s < 1 + \frac{m}{2\gamma}.$$

It results $m/p < s < m/\gamma$, then the function ν satisfies Hypotheses 2.1 and 2.2. Moreover, it is easy to verify that

$$|x|^{2\gamma} \in A_{1+1/s-1} \quad (\text{Muckenhoupt's class})$$

then, Hypotheses 2.9 and 2.10 hold with $\bar{t} = 2$.

Consider the following boundary value problem

$$(5.1) \quad -\operatorname{div} \left(\frac{|x|^\gamma}{1+|u|} |\nabla u|^{p-2} \nabla u \right) + u\{|u|^{p-2} + e^{|u|}|x|^\gamma |\nabla u|^p\} = g(x) \quad \text{in } \Omega,$$

$$(5.2) \quad u = 0 \quad \text{on } \partial\Omega,$$

where $g(x) \in L^\infty(\Omega)$.

In this case we have:

$$a_i(x, u, \nabla u) = \frac{|x|^\gamma}{1+|u|} |\nabla u|^{p-2} \frac{\partial u}{\partial x_i}, \quad i = 1, 2, \dots, m,$$

$$f(x, u, \nabla u) = u e^{|u|} |x|^\gamma |\nabla u|^p - \frac{1}{2} u |u|^{p-2} - g(x), \quad c_0 = \frac{3}{2}.$$

If we put $\lambda(|u|) = (1+|u|)e^{|u|}$, it is possible to verify all the Hypotheses 2.3–2.8. To verify (2.1), for example, it will be sufficient to note that the function $|u|^{p-2}/e^{-|u|}$ is bounded from above by $((p-2)/e)^{p-2}$ in $]-\infty, \infty[$.

Hence, boundary value problem (5.1), (5.2) has at least one weak solution in the sense (2.4), i.e., there exists at least one $u \in W$ such that

$$\int_{\Omega} \frac{|x|^\gamma}{1+|u|} |\nabla u|^{p-2} \nabla u \nabla w \, dx + \int_{\Omega} u\{|u|^{p-2} + e^{|u|}|x|^\gamma |\nabla u|^p\} w \, dx = \int_{\Omega} g(x) w \, dx$$

holds for every $w \in W$.

Moreover, from Theorem 2.11, u is locally Hölder continuous in Ω .

Example 5.2. Let $\Omega \subset \mathbb{R}^m$ be an open bounded set and let $g \in L^\infty(\Omega)$. Put $\nu(x) = 1$ in Ω and consider boundary value problem

$$-\operatorname{div} \left(\frac{1}{1+|u|^p} |\nabla u|^{p-2} \nabla u \right) + e^u - |u|^p + |\nabla u|^p = g(x) \quad \text{in } \Omega,$$

$$u = 0 \quad \text{on } \partial\Omega.$$

In this case we have:

$$a_i(x, u, \nabla u) = \frac{1}{1+|u|^p} |\nabla u|^{p-2} \frac{\partial u}{\partial x_i}, \quad i = 1, 2, \dots, m,$$

$$f(x, u, \nabla u) = e^u - |u|^p - u|u|^{p-2} + |\nabla u|^p - g(x), \quad c_0 = 1,$$

and $\lambda(|u|) = e^{|u|^p}$. All Hypotheses 2.1–2.8 are satisfied. It may be worth noting that the function $u(e^u - |u|^p - u|u|^{p-2})$ has minimum (negative) in \mathbb{R} . Hence

every functions $u \in W$ satisfying

$$\int_{\Omega} \frac{1}{1 + |u|^p} |\nabla u|^{p-2} \nabla u \nabla w \, dx + \int_{\Omega} \{e^u - |u|^p + |\nabla u|^p\} w \, dx = \int_{\Omega} g(x) w \, dx$$

for every $w \in W$ is locally Hölder continuous in Ω .

Acknowledgment. I would like to thank the referee for carefully reading my manuscript and for giving such constructive comments which substantially helped improving the quality of the paper.

REFERENCES

- [1] Bensoussan A., Boccardo L., Murat F., *On a nonlinear partial differential equation having natural growth terms and unbounded solution*, Ann. Inst. Henri Poincaré **5** (1988), no. 4, 347–364.
- [2] Boccardo L., Murat F., Puel J. P., *Existence de solutions faibles pour des équations elliptiques quasi-linéaires à croissance quadratique*, Nonlinear Partial Differential Equations and Their Applications, College de France Seminar, Vol. IV, Res. Notes in Math., 84, Pitman, London, 1983, 19–73 (French. English summary).
- [3] Boccardo L., Murat F., Puel J. P., *Résultat d'existence pour certains problèmes elliptiques quasiliéaires*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **11** (1984), no. 2, 213–235 (French).
- [4] Bonafede S., *Quasilinear degenerate elliptic variational inequalities with discontinuous coefficients*, Comment. Math. Univ. Carolin. **34** (1993), no. 1, 55–61.
- [5] Bonafede S., *Existence and regularity of solutions to a system of degenerate nonlinear elliptic equations*, Br. J. Math. Comput. Sci. **18** (2016), no. 5, 1–18.
- [6] Bonafede S., *Existence of bounded solutions of Neumann problem for a nonlinear degenerate elliptic equation*, Electron. J. Differential Equations **2017** (2017), no. 270, 1–21.
- [7] Cirmi G. R., D'Asero S., Leonardi S., *Fourth-order nonlinear elliptic equations with lower order term and natural growth conditions*, Nonlinear Anal. **108** (2014), 66–86.
- [8] Del Vecchio T., *Strongly nonlinear problems with hamiltonian having natural growth*, Houston J. Math. **16** (1990), no. 1, 7–24.
- [9] Drábek P., Nicolosi F., *Existence of bounded solutions for some degenerated quasilinear elliptic equations*, Ann. Mat. Pura Appl. **165** (1993), 217–238.
- [10] Fabes E. B., Kenig C. E., Serapioni R. P., *The local regularity of solutions of degenerate elliptic equations*, Comm. Partial Differential Equations **7** (1982), 77–116.
- [11] Gilbarg D., Trudinger N. S., *Elliptic Partial Differential Equations of Second Order*, Springer, Berlin, 1983.
- [12] Guglielmino F., Nicolosi F., *W-solutions of boundary value problems for degenerate elliptic operators*, Ricerche Mat. **36** (1987), suppl., 59–72.
- [13] Guglielmino F., Nicolosi F., *Existence theorems for boundary value problems associated with quasilinear elliptic equations*, Ricerche Mat. **37** (1988), 157–176.
- [14] John F., Nirenberg L., *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. **14** (1961), 415–426.
- [15] Kovalevsky A., Nicolosi F., *Boundedness of solutions of variational inequalities with nonlinear degenerated elliptic operators of high order*, Appl. Anal. **65** (1997), 225–249.
- [16] Kovalevsky A., Nicolosi F., *On Hölder continuity of solutions of equations and variational inequalities with degenerate nonlinear elliptic high order operators*, Current Problems of Analysis and Mathematical Physics, Taormina 1998, Aracne, Rome, 2000, 205–220
- [17] Kovalevsky A., Nicolosi F., *Boundedness of solutions of degenerate nonlinear elliptic variational inequalities*, Nonlinear Anal. **35** (1999), 987–999.
- [18] Kovalevsky A., Nicolosi F., *On regularity up to the boundary of solutions to degenerate nonlinear elliptic high order equations*, Nonlinear Anal. **40** (2000), 365–379.

- [19] Ladyzhenskaya O., Ural'tseva N., *Linear and Quasilinear Elliptic Equations*, translated from the Russian, Academic Press, New York-London, 1968.
- [20] Landes R., *Solvability of perturbed elliptic equations with critical growth exponent for the gradient*, J. Math. Anal. Appl. **139** (1989), 63–77.
- [21] Moser J., *A new proof of De Giorgi's theorem concerning the regularity problem for elliptic differential equations*, Comm. Pure Appl. Math. **13** (1960), pp. 457–468.
- [22] Murthy M. K. V., Stampacchia G., *Boundary value problems for some degenerate elliptic operators*, Ann. Mat. Pura Appl. (4) **80** (1968), 1–122.
- [23] Serrin J. B., *Local behavior of solutions of quasi-linear equations*, Acta Math. **111** (1964), 247–302.
- [24] Skrypnik I. V., *Nonlinear Higher Order Elliptic Equations*, Naukova dumka, Kiev, 1973 (Russian).
- [25] Skrypnik I. V., *Higher order quasilinear elliptic equations with continuous generalized solutions*, Differ. Equ. **14** (1978), no. 6, 786–795.
- [26] Trudinger N. S., *On Harnack type inequalities and their application to quasilinear elliptic equations*, Comm. Pure Appl. Math. **20** (1967), 721–747.
- [27] Trudinger N. S., *On the regularity of generalized solutions of linear non-uniformly elliptic equations*, Arch. Ration. Mech. Anal. **42** (1971), 51–62.
- [28] Trudinger N. S., *Linear elliptic operators with measurable coefficients*, Ann. Scuola Norm. Sup. Pisa **27** (1973), 265–308.
- [29] Voitovich M. V., *Existence of bounded solutions for a class of nonlinear fourth-order equations*, Differ. Equ. Appl. **3** (2011), no. 2, 247–266.
- [30] Voitovich M. V., *Existence of bounded solutions for nonlinear fourth-order elliptic equations with strengthened coercivity and lower-terms with natural growth*, Electron. J. Differential Equations **2013** (2013), no. 102, 25 pages.
- [31] Voitovich M. V., *On the existence of bounded generalized solutions of the Dirichlet problem for a class of nonlinear high-order elliptic equations*, J. Math. Sci. (N.Y.) **210** (2015), no. 1, 86–113.
- [32] Voitovych M. V., *Hölder continuity of bounded generalized solutions for nonlinear fourth-order elliptic equations with strengthened coercivity and natural growth terms*, Electron. J. Differential Equations **2017** (2017), no. 63, 18 pages.
- [33] Zamboni P., *Hölder continuity for solutions of linear degenerate elliptic equations under minimal assumptions*, J. Differential Equations **182** (2002), 121–140.

S. Bonafede:

DIPARTIMENTO DI AGRARIA, UNIVERSITÀ DEGLI STUDI MEDITERRANEA DI REGGIO CALABRIA, LOCALITÀ FEO DI VITO, 89122 REGGIO CALABRIA, ITALY

E-mail: salvatore.bonafede@unirc.it

(Received October 16, 2017, revised February 23, 2018)