

Order-enriched solid functors

LURDES SOUSA, WALTER THOLEN

In memory of Věra Trnková

Abstract. Order-enriched solid functors, as presented in this paper in two versions, enjoy many of the strong properties of their ordinary counterparts, including the transfer of the existence of weighted (co)limits from their codomains to their domains. The ordinary version of the notion first appeared in Trnková's work on automata theory of the 1970s and was subsequently studied by others under various names, before being put into a general enriched context by C. Anghel. Our focus in this paper is on differentiating the order-enriched notion from the ordinary one, mostly in terms of the functor's behaviour with respect to specific weighted (co)limits, and on the presentation of examples, which include functors of general varieties of ordered algebras and special ones, such as ordered vector spaces.

Keywords: ordered category; (strongly) order-solid functor; weighted (co)limit; ordered algebra

Classification: 18A22, 18A30, 18B35, 06F99

1. Introduction

Inspired by E. Čech's book [14] and M. Hušek's article [23], in her work [44] on *Automata and Categories* Věra Trnková defined a concrete category \mathcal{A} (which therefore comes with a faithful functor $|-|: \mathcal{A} \rightarrow \mathit{Set}$) to admit weak inductive generation if for every possibly large (!) family $(D_i)_{i \in I}$ of \mathcal{A} -objects, equipped with maps $\xi_i: |D_i| \rightarrow X$, $i \in I$, into a given set X , there exist an \mathcal{A} -object A and a map $q: X \rightarrow |A|$ such that

1. all maps $q \cdot \xi_i: |D_i| \rightarrow |A|$ underlie \mathcal{A} -morphisms $D_i \rightarrow A$, and
2. the pair (q, A) is universal with this property, i.e. for every map $f: X \rightarrow |B|$ with B in \mathcal{A} , such that all maps $f \cdot \xi_i: |D_i| \rightarrow |B|$ underlie \mathcal{A} -morphisms $D_i \rightarrow B$, there is a unique \mathcal{A} -morphism $t: A \rightarrow B$ with $|t| \cdot q = f$.

When q may always be chosen to be the identity map (so that A has underlying set X), this gives precisely the notion of concrete *topological category* or, when one

DOI 10.14712/1213-7243.2020.002

This work was partially supported by the Centre for Mathematics of the University of Coimbra – UID/MAT/00324/2019, funded by the Portuguese Government through FCT/MEC and co-funded by the European Regional Development Fund through the Partnership Agreement PT2020. The second author acknowledges partial financial support under the Discovery Grants Program by the Natural Sciences and Engineering Research Council of Canada.

trades *Set* for any “base category” \mathcal{X} , of (faithful) *topological functor* $P: \mathcal{A} \rightarrow \mathcal{X}$, see, for example, [1], [20]. This, as it turns out, self-dual notion was first introduced (in dual form and under a different name) in G. C. L. Brümmer’s thesis in [11]; V. Trnková, not aware of [11], calls \mathcal{A} to *admit inductive generation* in this case.

Other precursors to the notion of topological category or functor (first just over *Set*, but then over any category \mathcal{X}), such as [38], [8], [32], [47], [48], [30], [34], [18], [45], [22], limited the concept of inductive generation and its dualization to the consideration of small families, or even singleton-families (thus essentially considering Grothendieck’s *bifibrations*), and then imposed smallness conditions to effectively enable inductive generation for large families of data, such as asking each fibre of the given functor (i.e. each category of \mathcal{A} -objects with fixed underlying \mathcal{X} -object) to form a small complete lattice, as it was done by O. Wyler in his milestone papers [47], [48]. While credit for having elegantly introduced topologicity of a concrete category using large families is due to [11], this approach resonated with a wider audience only after the appearance of H. Herrlich’s important article [17], with its notions quickly expanded upon in other papers, such as [39], [46], [40].

For small families and, more generally, for small cocones, the concept of *weak inductive generation* was, without Trnková’s knowledge at the time, considered earlier by R.-E. Hoffmann in [18], under a different name and in rather cumbersome notation, and it appeared in published form only later, in [19]. Afterwards, unaware of Trnková’s notion, the authors of [41], [43], [37], [42] undertook a systematic and coherent study of the \mathcal{X} -based categories \mathcal{A} admitting weak inductive generation, showing their usefulness in the categorical investigation of a wide range of mathematical structures. These papers reconcile many themes studied earlier in the more restrictive context of topological categories and therefore call the functors involved (presenting the categories \mathcal{A} as concrete over \mathcal{X}) *semi-topological*, a term that had been used somewhat hiddenly in [19]. Even though every such functor may be presented as the composite of a full reflective embedding followed by a topological functor, the occurrence of these functors is by no means restricted to the realm of topology. Therefore, on Herrlich’s suggestion, they were renamed as *solid* later on, a term adopted in [10], [1] and used henceforth by others, for example in [35].

In his thesis [5], published and extended in [6], [7], C. Anghel takes the study of the (then) semi-topological functors comprehensively to the level of enriched category theory [25]. However, in order to do so, he needed to utilize the full range of the theory and often to impose additional conditions on the categories at issue, making it somewhat hard for the non-expert to apply his results. The purpose of this article is therefore to present a largely self-contained theory of solid functors in the easily presentable context of *order-enriched* categories and functors and their applications, that is: in an environment that has gained considerable attention in

recent years; see, for example, [3], [4], [12], [13], [21], [36].¹ Explicitly then, the hom-sets of our categories come equipped with a partial order that is preserved by the composition of the category and by the hom-maps of any functors departing from them, thus providing also an elementary 2-categorical context in which all 2-cells are given by order. However, even in this very simplified context one quickly arrives at subtleties that hinder a seamless transition of notions and results from the ordinary to the enriched context.

Therefore, in Section 2 we first present a notion of solidity for ordered functors, called *strongly order-solid*², which on first sight seems to add only a minor order-related condition to the ordinary notion. Nevertheless, it captures an extensive list of relevant examples, some of which appear in Section 3. It then turns out that the seemingly mild additional condition which makes ordinarily solid functors strongly order-solid already guarantees that they become solid as order-enriched functors in Anghel’s sense in [6], called *order-solid* here. We present these functors in Section 4 without assuming the reader’s familiarity with [25]. While strongly order-solid functors are easily seen to be order-solid, the converse question, whether every order-solid functor is strongly order-solid, is still open.

A central goal of the paper is the characterization of strongly order-solid and of order-solid functors in terms of their behaviour vis-a-vis weighted limits and colimits. In Theorem 2.6 we characterize strongly order-solid functors using inserters, and in Theorem 4.5 we state that they “lift” the existence of weighted (co)limits for diagrams of any given shape. We study the behaviour of order-solid functors on weighted colimits in Section 5 and characterize order-solid functors when the “base” category is tensored (Theorem 5.11). The list of examples in Section 3 culminates in a theorem on categories of general ordered algebras; Theorem 3.7 asserts that algebraic functors between them are always strongly order-solid as soon as they admit free algebras over every ordered set. The category of ordered vector spaces, considered as an ordered category via the positive cones of its objects, falls outside the scope of this theorem, but its positive-cone functor to the category of partially ordered sets is still strongly order-solid. When considered as a discretely ordered category, it serves as a resource to demonstrate that certain conditions of our characterization theorems are essential.

¹For the purpose of consistency with these papers, but at the price of divergence from other works (such as [20], [16]), in this paper we understand “order” to mean what is generally referred to as “partial order”. But we stress the fact that the theory presented here carries through smoothly when “order” means just “preorder” in more common parlance, perhaps even more so than in the partially ordered context. In fact, many general constructions lead from partially ordered sets just to preordered sets, which at the end have to be subjected to the reflector to enforce separation (= anti-symmetry), as demonstrated also by some of the examples presented in Section 3.

²In this paper the easily defined strongly order-solid functors appear before the more natural, but also slightly more complex, notion of order-solid functor, since we are not aware of examples of the latter type of functors not already covered by the former.

2. Strongly order-solid functors

We generally assume our categories and functors to be enriched in the Cartesian closed category \mathcal{Pos} of (partially) ordered sets and their monotone (= order-preserving) maps and simply call them *ordered*³. Hence, the hom-sets of an ordered category \mathcal{A} carry an order which is preserved by composition with morphisms from either side, and the hom-maps of an ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ preserve the order as well. In accordance with [25], whenever necessary for clarity, we write \mathcal{A}_o for the underlying ordinary category of an ordered category \mathcal{A} , and likewise for ordered functors.

Recall that the ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ has a left adjoint $F: \mathcal{X} \rightarrow \mathcal{A}$ in the order-enriched sense if there are order-isomorphisms

$$\mathcal{A}(FX, B) \cong \mathcal{X}(X, PB)$$

that are natural in $X \in \text{ob}\mathcal{X}$ and $B \in \text{ob}\mathcal{A}$; we call P *order-right adjoint* in this case. For that to happen it suffices that for every \mathcal{X} -object X , one finds a (tacitly chosen) P -universal arrow $e: X \rightarrow PA$ with $A \in \text{ob}\mathcal{A}$, see [29], which has the additional property of being *order- P -epi(morphic)*, that is: whenever $Pr \cdot e \leq Ps \cdot e$ for any morphisms $r, s: A \rightarrow B$ in \mathcal{A} , then $r \leq s$; equivalently, the ordered functor P is right adjoint in the ordinary sense such that all adjunction units are order- P -epic.

Given a (potentially large⁴) family $D = (D_i)_{i \in I}$ of objects in \mathcal{A} , we consider the (potentially very large⁵) category $D \downarrow \mathcal{A}$ whose objects are pairs (α, A) with an \mathcal{A} -object A and a family $\alpha = (\alpha_i)_{i \in I}$ of \mathcal{A} -morphisms $\alpha_i: D_i \rightarrow A$, shortly written as $\alpha: D \rightarrow A$; a morphism $t: (\alpha, A) \rightarrow (\beta, B)$ is given by an \mathcal{A} -morphism $t: A \rightarrow B$ satisfying $t \cdot \alpha_i = \beta_i$ for all $i \in I$, shortly written as $t \cdot \alpha = \beta$. Of course, $D \downarrow \mathcal{A}$ and, likewise, $PD \downarrow \mathcal{X}$ inherit the order from \mathcal{A} and \mathcal{X} , respectively, making both categories ordered, as well as the P -induced functor

$$P_D: (D \downarrow \mathcal{A}) \rightarrow (PD \downarrow \mathcal{X}), \quad (\alpha, A) \mapsto (P\alpha, PA).$$

Definition 2.1. An ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ is *strongly order-solid* if the functor P_D is order-right adjoint for every family D of \mathcal{A} -objects. Equivalently, given any D , for every family $\xi: PD \rightarrow X$ in \mathcal{X} there is a (tacitly chosen) family $\alpha: D \rightarrow A$ in \mathcal{A} and an \mathcal{X} -arrow $q: X \rightarrow PA$ such that

³As mentioned in the Introduction, and as will become apparent in Section 3, for many purposes we may alternatively work with the Cartesian closed category Ord of preordered sets and their monotone maps.

⁴That is: the size of the indexing system I may be as large as the size of the class of all morphisms of \mathcal{A} .

⁵We use the term “very large” informally, to refer to collections of (potentially proper) classes, called conglomerates in [1]. A formalization of the term does not seem to be justified in this paper, since one may, of course, avoid the formation of $D \downarrow \mathcal{A}$ and $PD \downarrow \mathcal{X}$ (the individual objects of which may already be large), but one will then have to accept universal quantification over these entities: see Definition 2.1.

1. (α, A, q) is a P -extension of ξ , that is: $P\alpha = q \cdot \xi$;
2. (α, A, q) is *universal* with respect to property 1, that is: for every family $\beta: D \rightarrow B$ in \mathcal{A} and every \mathcal{X} -arrow $f: X \rightarrow PB$ with $P\beta = f \cdot \xi$ one has a unique⁶ \mathcal{A} -morphism $t: A \rightarrow B$ with $t \cdot \alpha = \beta$ and $Pt \cdot q = f$;
3. $q: X \rightarrow PA$ is order- P -epimorphic.

The three properties together make (α, A, q) a *strongly order-universal P -extension* of ξ .

Remarks 2.2. (1) Just as order-right adjoint functors are in particular right-adjoint ordinary functors, every strongly order-solid functor is in particular solid in the ordinary sense and therefore faithful; see Lemma 3.2 of [41], the proof of which uses a Cantor-type diagonal argument, as presented more generally in [9]. But if P is faithful, given a family $\xi: PD \rightarrow X$, any family $\beta: D \rightarrow B$ with $P\beta = f \cdot \xi$ is already determined by B and $f: X \rightarrow PB$. Hence, the existence requirement of universal P -extensions for all D and ξ amounts precisely to Trnková’s admittance of weak inductive generation, as recorded at the beginning of the Introduction, to which we have only added the condition that all universal P -extensions be order- P -epic to make the ordered functor P strongly order-solid.

To see that a strongly order-solid functor as defined in Definition 2.1 is faithful, one in fact does not need to resort to the above argument, as a stronger property may be shown easily: see Proposition 2.4 below.

(2) Being in particular solid in the ordinary sense, a strongly order-solid functor $P: \mathcal{A} \rightarrow \mathcal{X}$ certainly enjoys all the “lifting properties” of solid functors, such as: if \mathcal{X} has all ordinary (co)limits (of diagrams of a specified shape), so does \mathcal{A} , see [41], [1]; if \mathcal{X} is totally cocomplete (so that its Yoneda embedding has a left adjoint in the ordinary sense), so is \mathcal{A} , see [42].

(3) All fully faithful order-right adjoint functors are strongly order-solid, and so are composites of strongly order-solid functors.

While we postpone the discussion of the behaviour of strongly order-solid functors with respect to *weighted* (co)limits until Sections 4 and 5, here we consider one (easy, but important) type of weighted limit since it helps clarifying the relationship of the notions of strongly order-solid functor and ordinarily solid functor.

Definition 2.3. (1) Recall that an *inserter* of a pair of morphisms $r, s: A \rightarrow B$ in an ordered category \mathcal{A} is a morphism $u: U \rightarrow A$ with $r \cdot u \leq s \cdot u$ that is universal with this property: any $v: V \rightarrow A$ with $r \cdot v \leq s \cdot v$ factors (uniquely) as $v = u \cdot j$; moreover, u is required to be *order mon(omorph)ic*, so that $u \cdot h \leq u \cdot k$ always implies $h \leq k$.

(2) An ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ is *order-faithful* if 1_{PA} is order- P -epic for all objects A in \mathcal{A} , that is: $Pr \leq Ps$ for morphisms $r, s: A \rightarrow B$ in \mathcal{A} always implies $r \leq s$.

⁶Uniqueness comes for free in the presence of condition 3, but only so because here we understand “ordered” to entail anti-symmetry. This observation applies analogously to many subsequent notions in this paper.

Proposition 2.4. *A strongly order-solid functor is solid in the ordinary sense, as well as order-right adjoint and order-faithful, and it preserves (any existing) inserters.*

PROOF: The first claim is obvious; see Remark 2.2 (1). Since, in Definition 2.1, families are allowed to be empty, order-right adjointness follows. It is standard to confirm the preservation of inserters (or any weighted limits) by order-right adjoint functors. So, only order-faithfulness of a strongly order-solid functor P needs to be shown here. But given $r, s: A \rightarrow B$ in \mathcal{A} with $Pr \leq Ps$, let $(a: A \rightarrow C, C, q: PA \rightarrow PC)$ be a universal P -extension of the singleton family $(1_{PA}: PA \rightarrow PA)$. Then Pr and Ps must both factor through q , so that for some $r', s': C \rightarrow B$ one has $Pr = Pr' \cdot q$ and $Ps = Ps' \cdot q$, as well as $r = r' \cdot a$ and $s = s' \cdot a$. Since q is order- P -epic, $r' \leq s'$ follows, which implies $r \leq s$. \square

Proposition 2.5. *Let \mathcal{A} have inserters. Then an ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ is strongly order-solid if and only if P is solid in the ordinary sense and order-faithful and preserves inserters.*

PROOF: After Proposition 2.4, only the “if”-part needs proof. To this end, it suffices to show that for the given families D and $\xi: PD \rightarrow X$ as in Definition 2.1, the universal P_o -extension (α, A, q) with respect to the ordinary functor $P_o: \mathcal{A}_o \rightarrow \mathcal{X}_o$ serves also as a strongly order-universal P -extension, that is: $q: X \rightarrow PA$ is necessarily order- P -epic. Hence, assuming $Pr \cdot q \leq Ps \cdot q$ for $r, s: A \rightarrow B$ in \mathcal{A} we form the inserter $u: U \rightarrow A$ of the pair r, s in \mathcal{A} which, by hypothesis, is preserved by P . So q factors as $q = Pu \cdot f$, with $f: X \rightarrow PU$. Since $P(r \cdot \alpha) = Pr \cdot q \cdot \xi \leq Ps \cdot q \cdot \xi = P(s \cdot \alpha)$ and P is order-faithful, $r \cdot \alpha \leq s \cdot \alpha$ follows. Consequently, the inserter u makes the family α factor as $\alpha = u \cdot \beta$. Since $Pu \cdot f \cdot \xi = q \cdot \xi = Pu \cdot P\beta$ and Pu (as an inserter) is monic in \mathcal{X} , one obtains $f \cdot \xi = P\beta$ and therefore an \mathcal{A} -morphism $t: A \rightarrow U$ with $Pt \cdot q = f$. From $P(u \cdot t) \cdot q = Pu \cdot f = q$ and q being (ordinarily) P -epic, one derives $u \cdot t = 1$. Since $r \cdot u \leq s \cdot u$, this finally implies $r \leq s$. \square

Theorem 2.6. *Let \mathcal{X} have inserters. An ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ is strongly order-solid if and only if*

- (a) P is solid as an ordinary functor;
- (b) \mathcal{A} has inserters and P preserves them;
- (c) P is order-faithful.

PROOF: That the conditions (a–c) are sufficient for P to be strongly order-solid has been confirmed in Proposition 2.5. Conversely, only the existence of inserters in \mathcal{A} still needs to be shown when \mathcal{X} has them and P is strongly order-solid. To this end, for any morphisms $r, s: A \rightarrow B$ in \mathcal{A} we form the inserter $k: X \rightarrow PA$ of Pr, Ps in \mathcal{X} and then consider the family ξ of all pairs (D, x) with $D \in \text{ob}\mathcal{A}$ and $x: PD \rightarrow X$ an \mathcal{X} -morphism such that there is a (necessarily unique) \mathcal{A} -morphism $a: D \rightarrow A$ with $Pa = k \cdot x$. (Note that, as an ordinarily solid functor, P is faithful.) With $q: X \rightarrow PU$ forming a universal P -extension of ξ we then see

that k must factor as $k = Pu \cdot q$ with $u: U \rightarrow A$ in \mathcal{A} . Since $P(r \cdot u) \cdot q = Pr \cdot k \leq Ps \cdot k = P(s \cdot u) \cdot q$ and q is order- P -epic, $r \cdot u \leq s \cdot u$ follows.

Furthermore, by the inserter property of k , any $v: V \rightarrow A$ in \mathcal{A} with $r \cdot v \leq s \cdot v$ produces a morphism $y: PV \rightarrow X$ with $k \cdot y = Pv$. This makes (V, y) a member of the family ξ , which implies that there is an \mathcal{A} -morphism $j: V \rightarrow U$ with $Pj = q \cdot y$. From $P(u \cdot j) = Pu \cdot q \cdot y = k \cdot y = Pv$ one obtains $u \cdot j = v$, as required. We note that the same argumentation may also be applied to u in place of v ; it produces morphisms $z: PU \rightarrow X$ and $t: U \rightarrow U$ with $k \cdot z = Pu$ and $Pt = q \cdot z$. Since k is monic, from $k \cdot z \cdot q = k$ one first obtains $z \cdot q = 1_X$, and then $Pt \cdot q = q \cdot z \cdot q = q$ forces $t = 1_U$ since q is P -epic. Consequently, $q \cdot z = 1_{PU}$, so that q and z must be isomorphisms in \mathcal{X} .

It remains to be shown that u is order-monic. If $u \cdot c \leq u \cdot d$ with $c, d: C \rightarrow U$ in \mathcal{A} , applying P to the inequality we first obtain $k \cdot z \cdot Pc \leq k \cdot z \cdot Pd$ and then $Pc \leq Pd$, since k is order-monic and z an isomorphism. As P is order-faithful, $c \leq d$ follows. □

We suppose that the existence assumptions regarding inserters are essential in Proposition 2.5 and Theorem 2.6 but have not been able yet to confirm this conjecture. However, preservation of inserters is: in Example 3.9 we exhibit a solid and order-faithful functor $P: \mathcal{A} \rightarrow \mathcal{X}$ (thus satisfying conditions (a) and (c) of the above theorem), with both \mathcal{X} and \mathcal{A} having inserters, but with P failing to preserve them; in particular, P fails to be order-right adjoint and, a fortiori, strongly order-solid. This still leaves open the following question:

Open Problem 2.7. Is a (ordinarily) solid, order-right adjoint and order-faithful functor $P: \mathcal{A} \rightarrow \mathcal{X}$ strongly order-solid? Equivalently, when \mathcal{X} has inserters, do these conditions on P imply the existence of inserters in \mathcal{A} ?

Remark 2.8. We recall from [41] (see Theorem 1.2 of [10] for a “direct” proof) that an ordinary functor $P: \mathcal{A} \rightarrow \mathcal{X}$ is solid if and only if P is right adjoint and there is a class \mathcal{E} of morphisms in \mathcal{A} such that

- (A) all adjunction co-units lie in \mathcal{E} ;
- (P) the pushout of a morphism in \mathcal{E} along any morphism exists in \mathcal{A} , and any such lies in \mathcal{E} ;
- (W) the wide pushout (= co-intersection) of a (possibly large) family of morphisms in \mathcal{E} with common domain exists in \mathcal{A} , and any such lies in \mathcal{E} .

For any morphism class \mathcal{E} , the category \mathcal{A} is said to be \mathcal{E} -cocomplete if conditions (P) and (W) hold. Note that (W) forces every morphism in \mathcal{E} to be an epimorphism in \mathcal{A} , see [9], [41]. Hence, the class \mathcal{E} may be assumed to be a class of epimorphisms a priori. Furthermore then, if \mathcal{A} is \mathcal{E} -cowellpowered, the consideration of small (= set-indexed) families in (W) suffices.

Following the proof for the ordinary characterization theorem of Remark 2.8 as given in Theorem 1.2 of [10], we easily arrive at the following characterization for strongly order-solid functors, which entails the ordinary version as the discretely ordered case.

Theorem 2.9. *An ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ is strongly order-solid if and only if P is order-right adjoint, and there exists a class \mathcal{E} of order-epimorphisms in \mathcal{A} such that the (ordinary) conditions (A), (P), (W) hold.*

PROOF: If P is strongly order-solid, P is order-right adjoint. Like in the proof for the ordinary case (see Theorem 2.1 of [10]) one considers the class \mathcal{E} of all those morphisms $e: A \rightarrow B$ in \mathcal{A} for which $Pe: PA \rightarrow PB$ is part of a universal P -extension of some family $\xi: PD \rightarrow PA$. But here, being order- P -epimorphic, such extension will make e order-epic, i.e., $r \cdot e \leq s \cdot e$ always implies $r \leq s$. Hence, \mathcal{E} is a class of order-epimorphisms which, being chosen as in the ordinary case, satisfies conditions (P) and (W). Furthermore, for the adjunction $F \dashv P$ with unit η and co-unit ε , as in the ordinary case one has that for every object A in \mathcal{A} , $P\varepsilon_A: PFPA \rightarrow PA$ serves as a universal P -extension (of the pair $(\eta_{PA}: PA \rightarrow PFPA, 1_{PFPA})$); but here we have to confirm that $P\varepsilon_A$ is order- P -epic. Indeed, since $P\varepsilon_A \cdot \eta_{PA} = 1_{PA}$ and P is order-faithful by Proposition 2.4, for all $r, s: A \rightarrow B$ with $Pr \cdot P\varepsilon_A \leq Ps \cdot P\varepsilon_A$ one obtains $Pr \leq Ps$ and then $r \leq s$. Consequently, $\varepsilon_A \in \mathcal{E}$, which shows (A).

Conversely, we know that conditions (A), (P), (W) make P solid as an ordinary functor, with universal P -extensions $(\alpha, A, q = Pe \cdot \eta_X: X \rightarrow PA)$ constructed in such a way that $e: FX \rightarrow A$ lies in the class \mathcal{E} (see Theorem 2.1 of [10]). As η_X is order- P -epic and e is order-epic, q must be order- P -epic, making it part of a strongly order-universal P -extension. □

3. Examples of strongly order-solid functors

For many of our examples it is convenient to first consider them in a preorder-enriched context, so that \mathcal{Pos} gets replaced by the larger Cartesian closed category \mathcal{Ord} of preordered sets. We will freely use the terms introduced in Section 2 in this context and thus talk about *preordered categories* and *functors*, *strongly preorder-universal P -extensions* and *strongly preorder-solid* functors, as well as about *preorder- P -epic* morphisms and *preorder-faithful* functors, keeping in mind that the latter two notions will no longer automatically imply that the morphisms will be P -epic or the functors be faithful in the ordinary sense.

The following proposition turns out to be useful in many concrete situations.

Proposition 3.1. *In the commutative diagram*

$$\begin{array}{ccc}
 \mathcal{A} & \xrightarrow{H} & \mathcal{A}' \\
 P \downarrow & & \downarrow P' \\
 \mathcal{X} & \xrightarrow{J} & \mathcal{X}'
 \end{array}$$

of preordered functors, let H and J be full embeddings, with H preorder-right adjoint. If P' is strongly preorder-solid, then P is also strongly preorder-solid, and trivially even strongly order-solid when it is an ordered functor.

PROOF: Being preorder-right adjoint, H is strongly preorder-solid, and so is its composite with the strongly preorder-solid functor P' (see Remark 2.2). Quite trivially now, as JP is strongly preorder-solid, with J being fully faithful, also P is strongly preorder-solid. Explicitly then, one constructs a strongly preorder-universal P -extension (α, A, q) of a P -cocone $\xi: PD \rightarrow X$ by composing a strongly preorder-universal P' -extension (α', A', q') of $J\xi: P'HD \rightarrow JX$ with (the P' -image of) a reflection $r: A' \rightarrow HA$ into A :

$$\begin{array}{ccccc}
 JPD & \xrightarrow{J\xi} & JX & \xrightarrow{q'} & P'A' & \xrightarrow{P'r} & P'HA = JPA \\
 & & \searrow^{Jf} & & \downarrow P't' & & \swarrow JPt \\
 & & & & P'HB = JPB & &
 \end{array}$$

(Note: A curved arrow labeled Jq also connects JX to $P'HA$)

□

Example 3.2. The functor

$$S: \mathcal{Top}_0 \rightarrow \mathcal{Pos}$$

provides the underlying set of a T_0 -topological space A with the (dual of the) *specialization order*, so that $x \leq y$ in SA means that the neighbourhood filter of x is finer than that of y (or that the ultrafilter fixed at x converges to y). With S , the category \mathcal{Top}_0 becomes order-enriched, that is: $f \leq g: A \rightarrow B$ in \mathcal{Top}_0 means $f \leq g: SA \rightarrow SB$ in \mathcal{Pos} , or $f(x) \leq g(x)$ in SB for all $x \in A$. We show that S is strongly order-solid.

In fact, since the specialization *preorder* may be defined for all topological spaces, so that S is the restriction of a preordered functor S' as in the diagram

$$\begin{array}{ccc}
 \mathcal{Top}_0 & \hookrightarrow & \mathcal{Top} \\
 s \downarrow & & \downarrow s' \\
 \mathcal{Pos} & \hookrightarrow & \mathcal{Ord}
 \end{array}$$

and since \mathcal{Top}_0 is epireflective in \mathcal{Top} , so that the surjective reflection morphisms make the embedding order-right adjoint, by Proposition 3.1 it suffices to show that S' is strongly preorder-solid.

Indeed, given a preordered set (X, \leq) and any family of monotone maps $\xi_i: SD_i \rightarrow (X, \leq)$ defined on topological spaces $D_i, i \in I$, we obtain a topology τ on the set X by declaring open all those down-closed sets $U \subseteq X$ for which the set $\xi_i^{-1}(U)$ is open in D_i for every $i \in I$. Then, obviously, $\text{id}_X: (X, \leq) \rightarrow S(X, \tau)$ is monotone and preorder- S -epic, and all maps $\xi_i: D_i \rightarrow (X, \tau)$ are continuous. When we are given any monotone map $f: (X, \leq) \rightarrow SB$ with a topological space B , such that all maps $f \cdot \xi_i: D_i \rightarrow B$ are continuous, then $f^{-1}(V)$ is down-closed for every open set V of B and indeed open in (X, τ) , thus making $f: (X, \tau) \rightarrow B$ continuous.

As a particular consequence of S being strongly order-solid, with Theorem 4.5 and Proposition 5.1 below one concludes that \mathcal{Top}_0 has all (small-indexed) weighted limits and colimits (as described in Section 4) since \mathcal{Pos} has them (see Examples 5.7 (1)), as previously observed in [12] and [4]. Likewise for \mathcal{Top} .

Example 3.3. Every *frame* (= complete lattice in which the binary meet distributes over arbitrary joins) A has an underlying *meet-semilattice* UA which just forgets the existence of arbitrary joins; likewise, one may forget the information that a homomorphism of frames preserves arbitrary joins and just keep the information of preservation of finite meets, to obtain a functor


$$U: \mathcal{Frm} \rightarrow \mathcal{SLat}.$$

With the order in both categories inherited from \mathcal{Pos} , this functor is order-enriched and right adjoint as such: for a meet-semilattice X , the adjunction unit $\downarrow: X \rightarrow UDX$ into the lattice DX of down-closed subsets of X (ordered by “ \subseteq ”) assigns to $x \in X$ the principal down-set $\downarrow x = \{z \in X : z \leq x\}$ in X ; it is easily seen to be order- U -epic since every down-closed subset of X is a join of principal down-sets.

In order to show that U is strongly order-solid, we consider a meet-semilattice X and a family of homomorphisms $\xi_i: UC_i \rightarrow X$, with frames $C_i, i \in I$. On the frame DX , one lets “ \sim ” be the least congruence relation such that

$$\forall i \in I, K \subseteq C_i \left(\downarrow \xi_i \left(\bigvee K \right) \sim \bigcup_{a \in K} \downarrow \xi_i(a) \right).$$

It is clear that, with the projection $p: DX \rightarrow A := DX/\sim$, all maps $p \cdot \downarrow \cdot \xi_i: UC_i \rightarrow A$ become frame homomorphisms. Furthermore, any meet-semilattice homomorphism $f: X \rightarrow UB$ to a frame B , for which all maps $f \cdot \xi_i: UC_i \rightarrow B$ are frame homomorphisms, gives us a frame homomorphism $f^\sharp: DX \rightarrow B$ whose induced congruence relation must contain “ \sim ”. Consequently, f^\sharp factors as $f^\sharp = t \cdot p$ with a frame homomorphism $t: A \rightarrow B$. Since $q := p \cdot \downarrow: X \rightarrow UA$ is clearly order- U -epic, this shows that q belongs to a strongly order-universal U -extension of the family ξ , as desired.



The above construction raises the question of how to “compute” the least congruence relation C on a frame A containing a given relation R on A —even though an answer is actually not needed in the proof above. In any case, the reader may consult [31] to see that the underlying set of A/C may be taken to contain all elements of A that are saturated with respect to R , that is: every

$s \in A$ such that for all $a, b, c \in A$, $a R b$ implies $(a \wedge c \leq s \iff b \wedge c \leq s)$. In this way, A/C becomes a frame, with the map $\pi: A \rightarrow A/C$ that assigns to $x \in A$ the infimum of all saturated elements s with $x \leq s$, acting as the quotient map; π satisfies the condition $(a R b \implies \pi(a) = \pi(b))$ and is universal with respect to it, that is: any frame homomorphism $g: A \rightarrow B$ with $(a R b \implies g(a) = g(b))$ factors as $g = h \cdot \pi$ with a frame homomorphism $h: A/R \rightarrow B$.

Example 3.4. That also the forgetful functor

$$V: \mathcal{SLat} \rightarrow \mathcal{Pos}$$

is strongly order-solid may be shown analogously to the previous example. Its left adjoint E is described as follows: for an ordered set X , one takes EX to contain the up-closures $\uparrow F$ of all finite subsets $F \subseteq X$, ordered by reverse inclusion “ \supseteq ”. Since $(\uparrow F) \cup (\uparrow G) = \uparrow(F \cup G)$, this makes EX a meet-semilattice and the map $\uparrow: X \rightarrow VEX$, $x \mapsto \uparrow x$, monotone and, in fact, as one easily sees, the unit of an adjunction, since $\uparrow F = \bigcup_{x \in F} \uparrow x$, i.e., every element in EX is a finite meet of “generic” elements.

Given a family of monotone maps $\xi_i: VC_i \rightarrow X$ with meet-semilattices C_i , $i \in I$, one considers the least congruence relation “ \sim ” on EX satisfying the condition

$$\forall i \in I, \quad a, b \in C_i \quad (\uparrow(\xi_i(a \wedge b)) \sim \uparrow \xi_i(a) \cup \uparrow \xi_i(b) \text{ and } \uparrow \xi_i(\top_i) \sim \emptyset),$$

where \top_i denotes the top element in C_i . By definition of “ \sim ”, with the projection $p: EX \rightarrow A = EX/\sim$, one obtains meet-semilattice homomorphisms $p \cdot \uparrow \cdot \xi_i: C_i \rightarrow A$ for all $i \in I$. Since “ \sim ” is contained in the congruence relation induced by the canonical extension $f^\#: EX \rightarrow B$ of any monotone map $f: X \rightarrow VB$ to a meet-semilattice B making all $f \cdot \xi_i$ homomorphisms, f factors uniquely through $p \cdot \uparrow: X \rightarrow VA$. That this map is order- V -epic follows again from the presentation $\uparrow F = \bigcup_{x \in F} \uparrow x$ of elements in EX .

As in Example 3.2, from Examples 3.3 and 3.4 we can draw the conclusion that \mathcal{SLat} and \mathcal{Frm} have all (small-indexed) weighted (co)limits.

Example 3.5. By an *ordered Abelian monoid* A we understand a commutative monoid object in the category \mathcal{Pos} , that is: A is a commutative monoid equipped with a partial order that makes its binary operation $+: A \times A \rightarrow A$ monotone. The morphisms of the resulting category $\mathbf{AbMon}(\mathcal{Pos})$ are monotone monoid homomorphisms. With the order of the hom-sets of $\mathbf{AbMon}(\mathcal{Pos})$ inherited from \mathcal{Pos} , we want to show that the forgetful functor $W: \mathbf{AbMon}(\mathcal{Pos}) \rightarrow \mathcal{Pos}$ is strongly order-solid. For that, in consideration of the commutative diagram

$$\begin{array}{ccc} \mathbf{AbMon}(\mathcal{Pos}) & \hookrightarrow & \mathbf{AbMon}(\mathcal{Ord}) \\ \downarrow W & & \downarrow W' \\ \mathcal{Pos} & \hookrightarrow & \mathcal{Ord} \end{array},$$

by Proposition 3.1 it suffices to show that the forgetful functor $W': \mathbf{AbMon}(\mathit{Ord}) \rightarrow \mathit{Ord}$ of *preordered Abelian monoids* (which, in comparison to ordered Abelian monoids, are missing only the anti-symmetry) is strongly preorder-solid, and that the top-row full inclusion functor is preorder-right adjoint. But the latter fact is easily guaranteed by General-Adjoint-Function-Theorem-type arguments (see, for example, [1], [20]), since $\mathbf{AbMon}(\mathit{Pos})$ is closed under point-separating families in $\mathbf{AbMon}(\mathit{Ord})$, so that we can focus on the former and first show that W' is preorder-right adjoint.

To this end, since we are not aware of a proof presented in the specific situation considered here (see [16] and the literature cited in there), we rely on general principles to confirm that W' , as an ordinary functor, is right adjoint, and apply the construction provided by O. Wyler’s *Taut Lift theorem* in [47]. Hence, for a preordered set X , we consider all monotone maps $f: X \rightarrow A_f$ whose codomain is any preordered Abelian monoid, and denote by $f^\sharp: FX \rightarrow A_f$ the homomorphism that extends f to the free Abelian monoid FX over the set X ; it consists of all formal sums $\sum_{x \in X} n_x x$ (with nonnegative integers n_x , all but finitely many being 0), and f^\sharp sends them to $\sum_{x \in X} n_x f(x)$. With

$$a \leq b \iff \forall f: X \rightarrow A_f (f^\sharp(a) \leq f^\sharp(b)),$$

it is easy to see that FX becomes a preordered Abelian monoid, making the insertion $\delta_X: X \rightarrow W'FX$ a W' -universal arrow, which turns out to be also order- W' -epic.

To finally see that W' is strongly preorder-solid, given a family $\xi = (\xi_i: W'C_i \rightarrow X)_{i \in I}$ of monotone maps from preordered Abelian monoids C_i , $i \in I$, to a preordered set X , we consider the least monoid congruence relation “ \sim ” on FX which, with the projection $p: FX \rightarrow FX/\sim$, makes all maps $p \cdot \delta_X \cdot \xi_i$ monoid homomorphisms. We must now define a preorder on FX/\sim , in such a way that FX/\sim becomes a preordered Abelian monoid with monotone projection p . To this end, let us call a monotone map $f: X \rightarrow A_f$ ξ -admissible if $f \cdot \xi_i: C_i \rightarrow A_f$ is a monotone homomorphism for all $i \in I$ and then define for all $a, b \in FX$,

$$p(a) \leq p(b) \iff \forall f: X \rightarrow A_f \xi\text{-admissible} (f^\sharp(a) \leq f^\sharp(b)).$$

Since “ \sim ” is generated by the pairs $(\delta_X(\xi_i(c+d)), \delta_X(\xi_i(c)) + \delta_X(\xi_i(d)))$, $c, d \in C_i$, $i \in I$, one sees that this preorder is well defined and has the desired properties.

Categories of ordered algebras, of which $\mathbf{AbMon}(\mathit{Pos})$ is an example, have gained the attention of several authors; see, for instance, [26] and the references given there. Hence, in what follows, we extend the previous example and consider any variety of any (possibly infinitary) type of general algebras instead of Abelian monoids. These are sets that come equipped with a class of (possibly infinitary) operations (instead of one binary and one nullary operation for monoids), which are required to satisfy certain equations (instead of the associativity, neutrality and commutativity requirements). Moreover, we must assume that one can form the free (pre)ordered general algebra of that type over a (pre)ordered set, with the

insertion of generators being order- $(-)^1$ -epic; here, as we explain next, $(-)^1$ denotes the forgetful functor from the category of (pre)ordered general algebras of the given type and their monotone homomorphisms to the category $\mathcal{P}os$ (or $\mathcal{O}rd$).

In the following theorem we formulate these facts in terms of Lawvere–Linton (infinitary) algebraic theories (as originally introduced in [27], [28]; for a modern treatment in the finitary case, see [2]). Explicitly then, paraphrasing [28] in the spirit of [2], by an (infinitary) algebraic theory \mathcal{T} we mean a category whose class of objects is the class of cardinal numbers, such that every cardinal n is the n -fold power of 1 in \mathcal{T} . An ordered \mathcal{T} -algebra A is a product-preserving functor $A: \mathcal{T} \rightarrow \mathcal{P}os$; its underlying ordered set is the value of A at 1. When we denote the value of A at n more suggestively by A^n , then A assigns to every n -ary term t of \mathcal{T} , i.e., to every morphism $t: n \rightarrow 1$ in \mathcal{T} , an n -ary monotone operation $At: A^n \rightarrow A^1$, written more conveniently as t_A .⁷ A monotone \mathcal{T} -homomorphism $f: A \rightarrow B$ of ordered \mathcal{T} -algebras is simply a natural transformation; its underlying monotone map is the component of the transformation at 1, which must commute with the n -ary operations t ; that is, when we write the underlying map of f as f again, $f \cdot t_A = t_B \cdot f^n$. With the order on its hom-sets inherited from $\mathcal{P}os$, this defines the ordered category $\mathbf{Alg}(\mathcal{T}, \mathcal{P}os)$, as a full subcategory of the ordered functor category $\mathcal{P}os^{\mathcal{T}}$. By replacing $\mathcal{P}os$ by $\mathcal{O}rd$ one obtains the category of preordered \mathcal{T} -algebras and the commutative diagram

$$\begin{array}{ccc}
 \mathbf{Alg}(\mathcal{T}, \mathcal{P}os) & \hookrightarrow & \mathbf{Alg}(\mathcal{T}, \mathcal{O}rd) \\
 (-)^1 \downarrow & & \downarrow (-)^1 \\
 \mathcal{P}os & \hookrightarrow & \mathcal{O}rd
 \end{array}$$

of preordered functors. We call the algebraic theory \mathcal{T} preorder-variatal if the functor $(-)^1: \mathbf{Alg}(\mathcal{T}, \mathcal{O}rd) \rightarrow \mathcal{O}rd$ is preorder-right adjoint and obtain, as in Example 3.5, the following quite general result:

Theorem 3.6. *For every preorder variatal algebraic theory \mathcal{T} , the forgetful functor*

$$U^{\mathcal{T}}: \mathbf{Alg}(\mathcal{T}, \mathcal{P}os) \rightarrow \mathcal{P}os, \quad A \mapsto A^1,$$

is strongly order-solid, and likewise when $\mathcal{P}os$ is traded for $\mathcal{O}rd$.

We do not give the proof of the theorem, not only since it follows the same argumentation as that of Example 3.5, but also since the theorem is a special case of Theorem 3.7, the proof of which we sketch in sufficient detail, albeit with a variation which avoids the use of Wyler’s theorem.

⁷For example, the morphisms $t: n \rightarrow m$ of the theory \mathcal{T} of Abelian monoids may be taken to be the homomorphisms $t: Fm \rightarrow Fn$ of the free Abelian monoids on m and n generators. Hence, for $m = 1$, t just picks an element in Fn , that is: an n -ary formal term, to which an algebra A as defined here assigns the actual n -ary operation t_A on its underlying set. Likewise for any other general algebraic structures admitting free algebras.

While Theorem 3.6 covers Examples 3.4 and 3.5, a generalization of Example 3.3 requires the consideration of *algebraic functors*, induced by morphisms of algebraic theories. Recall that a *morphism* $K : \mathcal{S} \rightarrow \mathcal{T}$ of algebraic theories \mathcal{S}, \mathcal{T} is simply a functor that maps objects identically and preserves their status as direct products. For example, the embedding of the theory of meet-semilattices into the theory of frames is a morphism of algebraic theories. Any morphism K of algebraic theories gives rise to the *ordered algebraic functor*

$$\text{Alg}(\mathcal{T}, \mathcal{Pos}) \rightarrow \text{Alg}(\mathcal{S}, \mathcal{Pos}), \quad A \mapsto AK,$$

which, for convenience, we denote by K again. In the example just mentioned, this then is the forgetful functor $\text{Frm} \rightarrow \text{SLat}$ as considered in Example 3.3.

Theorem 3.7. *The (pre)ordered algebraic functor induced by any morphism of preorder-variatal algebraic theories is strongly (pre)order-solid.*

PROOF: (Sketch) As in Example 3.5, by Proposition 3.1 it suffices that the (analogously defined) preordered functor K' of the commutative diagram

$$\begin{array}{ccc} \text{Alg}(\mathcal{T}, \mathcal{Pos}) & \hookrightarrow & \text{Alg}(\mathcal{T}, \mathcal{Ord}) \\ K \downarrow & & \downarrow K' \\ \text{Alg}(\mathcal{S}, \mathcal{Pos}) & \hookrightarrow & \text{Alg}(\mathcal{S}, \mathcal{Ord}) \end{array}$$

is strongly preorder-solid. With the notation for ordered algebras used also in the preordered case, the algebraic functor K' commutes with the forgetful functors of the algebraic categories, that is: $U^{\mathcal{S}}K' = U^{\mathcal{T}}$. Assuming that both $U^{\mathcal{T}}$ and $U^{\mathcal{S}}$ are preorder-right adjoint, we first show that K' is also preorder-right adjoint. To this end, we note that, according to Dubuc’s *adjoint triangle theorem*, see [15], the left adjoint L of the ordinary functor K' may be constructed with the help of the left adjoints $F^{\mathcal{T}} \dashv U^{\mathcal{T}}$, $F^{\mathcal{S}} \dashv U^{\mathcal{S}}$ and their adjunction units η and co-units ε . An inspection of the proof of Dubuc’s theorem reveals that the unit $\kappa : 1 \rightarrow K'L$ of $L \dashv K'$ makes the diagram

$$\begin{array}{ccc} F^{\mathcal{S}}U^{\mathcal{S}} & \xrightarrow{\varepsilon^{\mathcal{S}}} & 1 \\ \mu^{U^{\mathcal{S}}} \downarrow & & \downarrow \kappa \\ K'F^{\mathcal{T}}U^{\mathcal{S}} & \xrightarrow{K'\pi} & K'L \end{array}$$

commute; here $\mu : F^{\mathcal{S}} \rightarrow K'F^{\mathcal{T}}$ is the mate of $\eta^{\mathcal{T}} : 1 \rightarrow U^{\mathcal{S}}(K'F^{\mathcal{T}}) = U^{\mathcal{T}}F^{\mathcal{T}}$, and $\pi : F^{\mathcal{T}}U^{\mathcal{S}} \rightarrow L$ is a (pointwise) regular epimorphism in $\text{Alg}(\mathcal{T}, \mathcal{Ord})$. Now we can easily see that for every preordered \mathcal{S} -algebra B , the unit κ_B is preorder- K' -epic. Indeed, for $r, s : LB \rightarrow A$ in $\text{Alg}(\mathcal{T}, \mathcal{Pos})$ with $K'r \cdot \kappa_B \leq K's \cdot \kappa_B$, the commutativity of the diagram gives $K'(r \cdot \pi_{B^1}) \cdot \mu_{B^1} \leq K'(s \cdot \pi_{B^1}) \cdot \mu_{B^1}$, where $B^1 = U^{\mathcal{S}}B$. But the mate μ_{B^1} of $\eta_{B^1}^{\mathcal{T}}$ satisfies $U^{\mathcal{S}}\mu_{B^1} \cdot \eta_{B^1}^{\mathcal{S}} = \eta_{B^1}^{\mathcal{T}}$, so that with

$U^{\mathcal{S}}K' = U^{\mathcal{T}}$ we obtain

$$U^{\mathcal{T}}(r \cdot \pi_{B^1}) \cdot \eta_{B^1}^{\mathcal{T}} \leq U^{\mathcal{T}}(s \cdot \pi_{B^1}) \cdot \eta_{B^1}^{\mathcal{T}}.$$

Since $\eta_{B^1}^{\mathcal{T}}$ is preorder- $U^{\mathcal{T}}$ -epic and π_{B^1} surjective, $r \leq s$ follows.

Let $\xi = (\xi_i : K'C_i \rightarrow B)_{i \in I}$ be a family of monotone \mathcal{S} -homomorphisms from preordered \mathcal{T} -algebras $C_i, i \in I$, to a preordered \mathcal{S} -algebra B . Largely neglecting to write down forgetful functors now, on the \mathcal{T} -algebra LB we consider the least congruence relation “ \sim ” which makes the \mathcal{S} -homomorphisms $p \cdot \kappa_B \cdot \xi_i$ \mathcal{T} -homomorphisms, where $p : LB \rightarrow LB/\sim$ is the projection map. As in Example 3.5, we equip LB/\sim with the preorder defined by

$$p(a) \leq p(b) \iff \forall f : B \rightarrow A_f \text{ } \xi\text{-admissible } (f^\sharp(a) \leq f^\sharp(b)),$$

for all $a, b \in LB$; here f runs through all monotone \mathcal{S} -homomorphisms into some preordered \mathcal{T} -algebra, and the ξ -admissibility of f means that all maps $f \cdot \xi_i : C_i \rightarrow A$ need to be monotone \mathcal{T} -homomorphisms; $f^\sharp : LB \rightarrow A$ denotes the \mathcal{T} -homomorphism with $f^\sharp \cdot \kappa_B = f$. This makes LB/\sim an object of $\text{Alg}(\mathcal{T}, \text{Ord})$ and $p \cdot \kappa_B$ belong to a preorder-universal K -extension of ξ . □

Theorem 3.6 appears as a special case of Theorem 3.7 when one chooses for \mathcal{S} the initial algebraic theory, given by the dual of the full subcategory of Set with object class of all cardinal numbers, i.e. a skeleton of Set .

We continue with an important example of a strongly order-solid functor of a category of a generalized type of ordered algebras which, however, is not covered by Theorem 3.7, since only some of the algebraic operations are assumed to be monotone and, more importantly, since the order of the homomorphisms is not taken to be given pointwise by universal quantification over *all* elements of their common domain, but only over a part of it.

Example 3.8. By an *ordered vector space* V we understand a real vector space that comes equipped with a partial order for which the (binary) addition and all unary operations given by multiplication with any nonnegative scalar λ are monotone. Such V defines the *positive cone* $PV = \{v \in V : v \geq 0\}$, and a linear map $f : V \rightarrow W$ is said to be *positive* if it maps PV into PW ; equivalently, if $f : V \rightarrow W$ is monotone. Given another positive linear map $g : V \rightarrow W$, one writes

$$f \leq g \iff \forall v \geq 0 (f(v) \leq g(v)).$$

But to make sure that this preorder is anti-symmetric, we must assume that the positive cone PV is *generating*, that is, $V = PV + (-PV)$. Hence, we denote by OVec the category of all ordered vector spaces V whose positive cone is generating, and their positive linear maps. We obtain the ordered functor $P : \text{OVec} \rightarrow \text{Pos}$, and claim that P is *strongly order-solid*.

As in Example 3.5, we use Proposition 3.1. We note that it suffices to show that the analogously defined functor P' of *preordered vector spaces* with generating

positive cones, which extends P as in the diagram

$$\begin{array}{ccc}
 \mathcal{OVec} & \hookrightarrow & \mathcal{POVec} \\
 P \downarrow & & \downarrow P' \\
 \mathcal{Pos} & \hookrightarrow & \mathcal{Ord} ,
 \end{array}$$

is strongly preorder-solid, since it is easy to see that any such preordered vector space V admits a surjective reflection into \mathcal{OVec} : just consider V/U , where $U = \{v \in V : v \leq 0 \leq v\}$.

Proving first that P' is preorder-right adjoint, given a preordered set X , one extends its preorder “ \leq ” and considers the least preorder “ \leq ” of the free real vector space FX with basis X satisfying

1. if $x \leq y$ in X , then $0 \leq x \leq y$ in FX ;
2. if $u \leq v$ in FX and $w \in FX$, $\lambda \geq 0$, then $\lambda u + w \leq \lambda v + w$ in FX .

In this way the positive cone of FX becomes generating and the insertion $\eta_X : X \hookrightarrow FX$ a P' -universal arrow, which is also preorder- P' -epimorphic. Now, given a family of monotone maps $\xi_i : P'V_i \rightarrow X$ with preordered vector spaces V_i , $i \in I$, we consider all the vector space quotients $q : FX \rightarrow FX/K_q$, where the preorder of FX/K_q is such that it makes the quotient a preordered vector space, q a positive map and all $g_i = q \cdot \eta_X \cdot \xi_i$ positively linear, so that $g_i(\lambda u + \mu w) = \lambda g_i(u) + \mu g_i(w)$ for all $v, w \in PV_i$ and $\lambda, \mu \geq 0$. For K the intersection of all the subspaces K_q , the vector space FX/K comes then equipped with the preorder given by

$$v + K \leq w + K \iff \forall q (v + K_q \leq w + K_q).$$

This way we obtain a quotient $p : FX \rightarrow FX/K$ and monotone maps $p \cdot \eta_X \cdot \xi_i$. Since for each i , PV_i is generating in V_i , every $p \cdot \eta_X \cdot \xi_i$ has a unique linear extension $\alpha_i : V_i \rightarrow FX/K$ (because we may obtain a base contained in the positive cone). The monotone map $p \cdot \eta_X$ together with the family α forms the desired strongly preorder-universal P' -extension of ξ . We leave all details to the reader and refer to the literature, such as [24] or [33].

In the next example, we consider ordered vector spaces again, but now take the order of the hom-sets to be given by the pointwise order over the entire vector space, not just over the positive cone. Then the order becomes necessarily discrete and, although we still obtain an order-faithful forgetful functor to \mathcal{Pos} that is solid in the ordinary sense, it fails to be strongly order-solid. This shows in particular that, in Theorem 2.6, we cannot omit condition (b).

Example 3.9. Let $\mathcal{OVec}_=$ be the category of ordered real vector spaces and linear maps which preserve the order. Given a pair of morphisms $f, g : V \rightarrow U$, since the inequality $f(u) \leq g(u)$ implies $g(-u) \leq f(-u)$, imposing the inequality $f(v) \leq g(v)$ for all $v \in V$ forces $f = g$. Hence, $\mathcal{OVec}_=$ is trivially order-enriched

via the discrete order, and the forgetful ordered functor

$$R: \mathcal{OVec}_= \rightarrow \mathcal{Pos}$$

is order-faithful. We show that R is also solid in the ordinary sense but fails to be strongly order-solid.

In order to show that R is solid, we can follow a path completely analogous to the one used in the previous example when we showed that P is strongly order-solid, including the use of Proposition 3.1 in its non-enriched version (that is, with the discrete order between morphisms). Here, to show that the forgetful functor into \mathcal{Ord} is a right-adjoint, we change condition 1 of the description of the preorder on the freely generated vector space FX of Example 3.8, by replacing $0 \leq x \leq y$ with $x \leq y$.

As a solid ordinary functor, R has a left adjoint H , but the adjunction units $X \hookrightarrow RHX$ will generally fail to be order- R -epic. Indeed, for X the 2-chain $\{a < b\}$, let the real valued maps $f^\sharp, g^\sharp: HX \rightarrow \mathbb{R}$ be determined by the monotone maps $f, g: X \rightarrow \mathbb{R}$ with $0 = f(a) < g(a) = f(b) = 2 < g(b) = 3$. Then, for $u = b - a$, $f^\sharp(u) \not\leq g^\sharp(u)$. Consequently R is not strongly order-solid.

In conclusion, the functor R fulfils conditions (a) and (c) of Theorem 2.6, that is, R is ordinarily solid and order-faithful, but does not fulfil (b), since R fails to preserve inserters. Indeed, in $\mathcal{OVec}_=$ inserters are just equalizers, since the order between morphisms is discrete, but not so in \mathcal{Pos} .

The above arguments also show that, analogously, we have a preorder-faithful functor which is solid but not strongly preorder-solid, and which does not preserve inserters.

4. Order-enriched solid functors

Following Anghel’s lead [5], [6] we now look at notions of universal P -extension and solidity for order-enriched functors from the general enriched categorical perspective. An *ordered diagram (of shape \mathcal{D})* in an ordered category \mathcal{A} is an ordered functor $D: \mathcal{D} \rightarrow \mathcal{A}$; we do not restrict the size of the ordered category \mathcal{D} . For a given \mathcal{D} and an object A in \mathcal{A} , when there is no risk of confusion we denote the constant functor $\mathcal{D} \rightarrow \mathcal{A}$ with value A again by A ; a morphism $t: A \rightarrow B$ is then treated as a natural transformation of constant functors. A *weight* for an ordered diagram of shape \mathcal{D} is an ordered functor $W: \mathcal{D}^{\text{op}} \rightarrow \mathcal{Ord}$. (Note that, in forming \mathcal{D}^{op} , one turns around the arrows of \mathcal{D} while maintaining their order.) Every object B in \mathcal{A} gives the weight

$$\mathcal{A}(D-, B) = (\mathcal{D}^{\text{op}} \xrightarrow{D^{\text{op}}} \mathcal{A}^{\text{op}} \xrightarrow{\mathcal{A}(-, B)} \mathcal{Pos}), \quad i \mapsto \mathcal{A}(Di, B),$$

and an \mathcal{A} -morphism $t: A \rightarrow B$ then becomes a natural transformation $\mathcal{A}(D-, t): \mathcal{A}(D-, A) \rightarrow \mathcal{A}(D-, B)$, i.e., a morphism in the (potentially very large) ordered

category $\mathcal{Pos}^{\mathcal{D}^{\text{op}}}$ of weights for \mathcal{D} , the morphisms of which are ordered component-wise. Pushing things even further, we note that, of course, $\mathcal{A}(D-, B)$ is functorial in B , i.e., one has the hom-functor

$$H_D^W : \mathcal{A} \rightarrow \mathcal{POS}, \quad B \mapsto \mathcal{Pos}^{\mathcal{D}^{\text{op}}}(W, \mathcal{A}(D-, B)),$$

whose codomain may be very large⁸.

To fix our notation and terminology, we recall the notion of weighted colimit in both, elementary “pointwise” form and standard terms of enriched category theory, before proceeding similarly for the enriched notion of order-solidity.

Remarks 4.1. (1) A *weighted cocone* $\alpha: D \rightarrow A$ over an ordered diagram $D: \mathcal{D} \rightarrow \mathcal{A}$ of weight $W: \mathcal{D}^{\text{op}} \rightarrow \mathcal{Pos}$ (briefly referred to as a W -weighted cocone over D) is given by its *vertex* $A \in \text{ob}\mathcal{A}$ and a natural transformation $\alpha: W \rightarrow \mathcal{A}(D-, A)$, that is, a family of \mathcal{A} -morphisms $\alpha_i^u: Di \rightarrow A$, $i \in \text{ob}\mathcal{D}$, $u \in Wi$, satisfying the conditions

- $\forall i \in \text{ob}\mathcal{D}, u, v \in Wi (u \leq \implies \alpha_i^u \leq \alpha_i^v)$;
- $\forall d: i \rightarrow j \text{ in } \mathcal{D}, v \in Wj (\alpha_j^v \cdot Dd = \alpha_i^{Wd(v)})$.

\mathcal{D} is the *shape* of the cocone.

(2) A W -*weighted colimit* of D is given by a W -weighted cocone $\alpha: D \rightarrow A$ such that

- $\alpha: D \rightarrow A$ is universal amongst all W -weighted cocones $\beta: D \rightarrow B$, i.e., any such β factors through α , so that there is a unique⁹ \mathcal{A} -morphism $t: A \rightarrow B$ with $\beta = t \cdot \alpha$; that is,
 - $\forall i \in \text{ob}\mathcal{D}, u \in Wi (\beta_i^u = t \cdot \alpha_i^u)$;
- $\alpha: D \rightarrow A$ is *order-epic*, so that for all $r, s: A \rightarrow B$ in \mathcal{A} one has the implication
 - $\forall i \in \text{ob}\mathcal{D}, u \in Wi (r \cdot \alpha_i^u \leq s \cdot \alpha_i^u) \implies r \leq s$;
 we write more economically $(r \cdot \alpha \leq s \cdot \alpha \implies r \leq s)$ for this implication.

It is easy to check that this equivalently means that H_D^W is representable, i.e., $H_D^W \cong \mathcal{A}(A, -)$ as \mathcal{POS} -valued functors, making H_D^W in effect \mathcal{Pos} -valued.

(3) A weighted cocone $\alpha: D \rightarrow A$ whose weight $W = 1: \mathcal{D}^{\text{op}} \rightarrow \mathcal{Pos}$ maps the \mathcal{D} -objects constantly to the terminal ordered set 1 is simply an ordinary cocone over the ordinary diagram D_o . Consequently, a 1-weighted cocone $\alpha: D \rightarrow A$ is a weighted colimit precisely when it is an ordinary colimit of D_o and order-epic. Such weighted colimits are usually called *conical*.

(4) A weighted cocone in \mathcal{A} over an empty diagram is just an object of \mathcal{A} . By (3), a weighted colimit over the empty diagram is just an ordinary colimit, i.e., an initial object of the category \mathcal{A} .

⁸The objects of \mathcal{POS} are the partially ordered classes. Concerning the informal term “very large” and the formation of \mathcal{POS} , the same comment as the one made before Definition 2.1 (as footnote 5) applies here.

⁹Uniqueness is automatically guaranteed by the subsequent condition of α being order-epic.

(5) A diagram D in \mathcal{A} over the terminal (ordered) category $\mathcal{D} = 1$ can be viewed as an object D of \mathcal{A} ; likewise, a weight W with domain 1 is to be considered as an ordered set W . A W -weighted cocone with vertex $A \in \text{ob}\mathcal{A}$ is then a monotonely W -indexed family of morphisms $\alpha^u : D \rightarrow A$ in the ordered category \mathcal{A} , so that $\alpha^u \leq \alpha^v$ whenever $u \leq v$ in W . If it is even a weighted colimit, A is usually written as a *tensor product* $W \otimes D$, so that then the colimit property is described by the existence of order-isomorphisms

$$\mathcal{A}(W \otimes D, B) \cong \text{Pos}(W, \mathcal{A}(D, B)),$$

naturally in $B \in \text{ob}\mathcal{A}$. The category \mathcal{A} is *tensored* if $W \otimes D$ exists for all $D \in \text{ob}\mathcal{A}$ and $W \in \text{ob}\text{Pos}$.

(6) Of particular interest is also the discretely ordered category $\mathcal{D} = \{a, b : 0 \rightarrow 1\}$ with exactly two objects and exactly two non-identical arrows, together with the weight $W : \mathcal{D}^{\text{op}} \rightarrow \text{Pos}$ defined by

$$W1 = 1 = \{*\}, \quad W0 = 2 = \{u < v\}, \quad Wa(*) = u, \quad Wb(*) = v,$$

sometimes referred to as the *Walking Two*. A diagram $D : \mathcal{D} \rightarrow \mathcal{A}$ is simply a pair $f, g : C \rightarrow B$ of morphisms in \mathcal{A} , and a W -weighted colimit of that diagram is called a *co-inserter* for f, g , i.e., it is an order-epic arrow $e : B \rightarrow A$, universal with respect to the property $e \cdot f \leq e \cdot g$, so that any $h : B \rightarrow E$ with $h \cdot f \leq h \cdot g$ factors as $h = t \cdot e$.

(7) *Weighted limits* in \mathcal{A} are, by definition, weighted colimits in \mathcal{A}^{op} .

Definition 4.2.

(1) For an ordered functor $P : \mathcal{A} \rightarrow \mathcal{X}$ and a W -weighted cocone $\xi : PD \rightarrow X$ in \mathcal{X} , we call the triple (α, A, q) , consisting of a W -weighted cocone $\alpha : D \rightarrow A$ in \mathcal{A} and an \mathcal{X} -morphism $q : X \rightarrow PA$, a *P-extension* of ξ if $q \cdot \xi = P\alpha$; that is, if $q \cdot \xi_i^u = P\alpha_i^u$ for all $i \in \text{ob}\mathcal{D}$, $u \in Wi$.

(2) (α, A, q) is an *order-universal P-extension* of the W -weighted cocone $\xi : PD \rightarrow X$ if

1. (α, A, q) is a *P-extension* of ξ ;
2. (α, A, q) is *universal* with respect to property 1, that is, for every *P-extension* (β, B, f) of ξ there is a (unique) \mathcal{A} -morphism $t : A \rightarrow B$ with $t \cdot \alpha = \beta$ and $Pt \cdot q = f$;
3. (α, A, q) is *order-P-epi(morphi)c*, that is, for all $r, s : A \rightarrow B$ in \mathcal{A} one has the implication

$$Pr \cdot e \leq Ps \cdot e, r \cdot \alpha \leq s \cdot \alpha \implies r \leq s.$$

(3) An ordered functor $P : \mathcal{A} \rightarrow \mathcal{X}$ is *order-solid* if every weighted cocone $\xi : PD \rightarrow X$ with any ordered diagram $D : \mathcal{D} \rightarrow \mathcal{A}$ of any weight $W : \mathcal{D}^{\text{op}} \rightarrow \text{Pos}$ has an order-universal *P-extension*.

Remarks 4.3. (1) It is important to observe the difference of condition 3 in Definitions 2.1 and 4.2: if the \mathcal{X} -morphism $q : X \rightarrow PA$ with $A \in \text{ob}\mathcal{A}$ is order-*P-epic*, so is the *P-extension* (α, A, q) , but not necessarily conversely. Reconciliation

Theorem 4.5. *An ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ is strongly order-solid if and only if it is order-solid and order-faithful. In this case, if \mathcal{X} has all weighted limits of a given shape \mathcal{D} , so does \mathcal{A} .*

PROOF: For the first statement, if $P: \mathcal{A} \rightarrow \mathcal{X}$ is order-solid and order-faithful, the morphism q of any order-universal P -extension of a weighted cocone ξ is order-P-epic, by Lemma 4.4. This holds particularly when ξ is a 1-weighted cocone over a discrete diagram, as needed to satisfy Definition 2.1. Conversely, let P be strongly order-solid. By Proposition 2.4, P is order-faithful. Furthermore, in order to construct an order-universal P -extension of any W -weighted cocone $\xi: PD \rightarrow X$ with $D: \mathcal{D} \rightarrow \mathcal{A}$, one considers $\xi = (\xi_i^u: Di \rightarrow X)_{i \in \text{ob } \mathcal{D}, u \in Wi}$ as a discretely indexed family of morphisms for which we have a strongly universal P -extension $(\alpha = (\alpha_i^u: Di \rightarrow A)_{i,u}, A, q)$, by hypothesis. Since P is order-faithful, from $u \leq v$ in Wi and, hence, $P\alpha_i^u = \xi_i^u \cdot q \leq \xi_i^v \cdot q = P\alpha_i^v$, one concludes $\alpha_i^u \leq \alpha_i^v$; and, analogously, for $d: i \rightarrow j$ in \mathcal{D} , the faithfulness of P implies $\alpha_j^v \cdot Dd = \alpha_i^{Wd(v)}$. This makes (α, A, q) an order-universal P -extension of the W -weighted cocone ξ .

For the second statement (on the existence of weighted limits), one proceeds analogously to the proof of Theorem 2.6 which deals with the special case of inserters. □

In the ordinary case, that is, when the categories are ordered discretely, the notions of solid and strongly solid are equivalent, since, as proved in [41], every solid functor is faithful. But we have not been able to decide whether order-faithfulness is an essential condition in Theorem 4.5:

Open Problem 4.6. Is every order-solid functor order-faithful?

5. Order-solid functors and weighted colimits

For the study of the behaviour of order-solid functors with respect to weighted colimits, we first consider order-universal P -extensions of individual cocones, without the universal quantification over all such data. An easy, but nevertheless fundamental, observation in this regard is that, in generalization of a well-known property of the ordinary notions, the order-universal P -extension of a weighted colimit of PD gives a weighted colimit of D , as stated in the next proposition. In Remark 5.5 we recall some important types of weighted colimits and some of their properties.

Proposition 5.1. *For an ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$, an ordered diagram $D: \mathcal{D} \rightarrow \mathcal{A}$ and a weight $W: \mathcal{D}^{\text{op}} \rightarrow \text{Ord}$, let $\xi: PD \rightarrow X$ be a W -weighted colimit of PD in \mathcal{X} . Then:*

- (1) *If $(\alpha: D \rightarrow A, A, q: X \rightarrow PA)$ is an order-universal P -extension of ξ , then α is a W -weighted colimit of D in \mathcal{A} .*
- (2) *If $\alpha: D \rightarrow A$ is a W -weighted colimit of D and $q: X \rightarrow PA$ the comparison morphism induced by ξ , then (α, A, q) is an order-universal P -extension of ξ .*

PROOF: (1) The colimit property of ξ makes $P\beta$ for any cocone $\beta: D \rightarrow B$, factor as $P\beta = f \cdot \xi$, with $f: X \rightarrow PB$ in \mathcal{X} . Order-universality of (α, A, q) gives $t: A \rightarrow B$ in \mathcal{A} with $Pt \cdot q = f$ and $t \cdot \alpha = \beta$. For any morphisms $r, s: A \rightarrow B$ with $r \cdot \alpha \leq s \cdot \alpha$, order preservation by P gives $(Pr \cdot q) \cdot \xi \leq (Ps \cdot q) \cdot \xi$, whence $Pr \cdot q \leq Ps \cdot q$ follows since ξ is order-epic. With (α, A, q) being order- P -epic, we conclude $r \leq s$.

(2) The proof proceeds similarly to the proof of (1). □

Corollary 5.2. *Let the ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ admit order-universal P -extensions for all weighted cocones of shape \mathcal{D} . Then, if \mathcal{X} has weighted colimits for diagrams of shape \mathcal{D} , so does \mathcal{A} ; likewise for conical colimits instead of weighted colimits.*

Corollary 5.3. *If $P: \mathcal{A} \rightarrow \mathcal{X}$ is order-solid and \mathcal{X} has weighted colimits of shape \mathcal{D} , so does \mathcal{A} .*

Corollary 5.4. *If the ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ admits order-universal P -extensions for all weighted cocones of shape $\mathbf{1}$ and \mathcal{X} is tensored, so is \mathcal{A} .*

Remarks 5.5. (1) Recall that for morphisms $f: A \rightarrow B, g: A \rightarrow C$ in an ordered category \mathcal{A} , a *cocomma object* for (f, g) is given by an object D and morphisms $p: B \rightarrow D, q: C \rightarrow D$ in \mathcal{A} with $p \cdot f \leq q \cdot g$

$$\begin{array}{ccc}
 A & \xrightarrow{g} & C \\
 f \downarrow & \lrcorner & \downarrow q \\
 B & \xrightarrow{p} & D
 \end{array}$$

and (p, q) is universal with that property (so that any pair (k, l) with common codomain E and $k \cdot f \leq l \cdot g$ must factor through (p, q) by a morphism $t: D \rightarrow E$; moreover, the pair (p, q) is required to be jointly order-epic.

(2) Similarly to co-inserters, also cocomma objects are easily recognized as weighted colimits: instead of the discretely ordered diagram shape with a parallel pair, consider a span and define the weight of its domain and codomains as for the Walking Two (see Remark 4.1 (7)).

(3) It is easy to see that one may construct the cocomma object of (f, g) as in (1) by forming the conical coproduct $B + C$ with injections i, j and then the co-inserter $c: B + C \rightarrow D$ of $(f \cdot i, g \cdot j)$.

(4) Conversely to (3), having cocomma objects at one's disposal, one may construct the co-inserter $c: B \rightarrow C$ of a pair $(f, g: A \rightarrow B)$ by forming their cocomma object $(p, q: B \rightarrow D)$ and then the conical coequalizer $e: D \rightarrow C$ of the pair (p, q) .

(5) As a consequence of (3) and (4), in the presence of finite conical colimits, the existence of cocomma objects is equivalent to the existence of co-inserters.

(6) It is well known (see Lemma 3.13 of [4] in the dual situation) that the tensor product $W \otimes A$ may be constructed with conical copowers and co-inserters, as

follows: presenting the order of W as a subset W_1 of $W_0 \times W_0$, with W_0 the underlying set of W , which comes with projections $d_1, d_2: W_1 \rightarrow W_0$, one forms the conical copowers $W_1 \otimes A, W_0 \otimes A$ (which are, in fact tensor products with discretely ordered sets) and then the co-inserters of the induced morphisms $d_1 \otimes A, d_2 \otimes A: W_1 \otimes A \rightarrow W_0 \otimes A$.

(7) A standard result of enriched category theory (see Theorem 3.73 of [25]) guarantees the existence of all weighted colimits of small shape in \mathcal{A} when \mathcal{A} has tensor products, conical coproducts and conical coequalizers. Taking the preceding remarks into account, one obtains: *The ordered category \mathcal{A} has all weighted colimits of small shape if and only if it has small-indexed conical coproducts, conical coequalizers and at least one—and then all—of the following types of weighted colimits: tensor products, co-inserters, or cocomma objects.*

With Remark 5.5 (2), Proposition 5.1 gives:

Corollary 5.6. *If the ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ admits order-universal P -extensions of cocones of finite shape and \mathcal{X} has cocomma objects (respectively, co-inserters), so does \mathcal{A} .*

We return to the examples presented in Section 3.

Examples 5.7. (1) Conical colimits in $\mathcal{P}os$ are given by ordinary colimits. The tensor product $W \otimes A$ may be given as $W \times A$, ordered like the direct product. The cocomma object of $(f: A \rightarrow B, g: A \rightarrow C)$ has as its underlying set the union $B \cup C$, which may be assumed to be disjoint; one then maintains the orders of its subsets B and C and adds to that the condition that $y \leq z$ holds for $y \in B$ and $z \in C$ if $y \leq f(x)$ and $g(x) \leq z$ for some $x \in A$.

(2) In $\mathcal{S}Lat$, the (conical) copower of A indexed by a set W_0 , denoted $A^{(W_0)}$, is the sub-semilattice of the power A^{W_0} whose elements have all but finitely many coordinates equal to the top element \top ; each injection p_w maps every a to $(a_w) \in A^{W_0}$ with $a_w = a$ and $a_u = \top$ for $u \neq w$. The tensor product $A \otimes W$ is the quotient $A^{(W_0)}/\sim$ where W_0 is the underlying set of W and “ \sim ” is the least congruence containing the pairs $(p_u(a) \wedge p_v(a), p_u(a))$ for all $a \in A$ and $u \leq v$ in W . Given $f: A \rightarrow B$ and $g: A \rightarrow C$ in $\mathcal{S}Lat$, let $B \times C$ be the product in $\mathcal{P}os$ (then also the conical product and conical coproduct in $\mathcal{S}Lat$); the cocomma object of (f, g) is the quotient $B \times C/\sim$, where “ \sim ” is the least congruence relation with $(f(a), r) \sim (f(a), s)$ for all $a \in A$ and $r, s \geq g(a)$.

(3) The characterization of weighted colimits in $\mathcal{F}rm$ is more involved. Concerning the conical coproduct, if we first take it in $\mathcal{S}Lat$ and then form the order-universal U -extension of the corresponding U -sink using the construction of Example 3.3, we obtain precisely the description of the coproduct given in [31]. We can proceed in an analogous way for coequalizers (see also [31]), tensor products, co-inserters and cocomma objects.

(4) In $\mathcal{O}Vec$, given morphisms $f, g: V \rightarrow W$, we describe the co-inserters of (f, g) . Let C be the cone given by the sum of PW with the cone $S = \{g(v) - f(v): v \in PV\}$, so $C = PW + S = \{u + w: u \in PW, w \in S\}$. The intersection

$U = C \cap (-C)$ is a subspace of W . Let W/U be the quotient space of W whose order has positive cone $P(W/U) = \{w + U : w \in C\}$. Then the co-inserter of the pair (f, g) is precisely the projection $W \rightarrow W/U$. With this characterization of the co-inserters, it is easy to obtain similar descriptions for tensor products and cocomma objects, using the fact that conical coproducts in \mathcal{OVec} are just the usual direct sums of spaces with the positive cone given by the sum of the positive cones of the components of the sum; see Remark 5.5 (3) and (6).

Guided by Anghel’s Theorem 2.2.8 in [6], we now give a step-by-step analysis of what may be needed to construct an order-universal P -extension $(\alpha : D \rightarrow A, A, q : X \rightarrow PA)$ of a given W -weighted cocone $\xi : PD \rightarrow X$ with $D : \mathcal{D} \rightarrow \mathcal{A}$, assuming that we have some particular weighted colimits and a certain order-universal P -extension over an $\text{ob}\mathcal{D}$ -indexed discrete diagram at our disposal.

Step 1: For every $i \in \text{ob}\mathcal{D}$, we assume that the tensor products $(\lambda_i^u : Di \rightarrow Wi \otimes Di)_{u \in Wi}$ and $(\kappa_i^u : PDi \rightarrow Wi \otimes PDi)_{u \in Wi}$ with comparison morphisms $c_i : Wi \otimes PDi \rightarrow P(Wi \otimes Di)$ exist in \mathcal{A} and \mathcal{X} , respectively. (Of course, by Corollary 5.4, the former tensor product may be obtained from the latter by an order-universal P -extension.) For every i , we let $\xi_i : Wi \otimes PDi \rightarrow X$ be the induced \mathcal{X} -morphism satisfying $\xi_i \cdot \kappa_i^u = \xi_i^u$ for all $u \in Wi$.

Step 2: We assume that in \mathcal{X} there exists the conical *generalized pushout* diagram

$$\begin{array}{ccc}
 Wi \otimes PDi & \xrightarrow{c_i} & P(Wi \otimes Di) \\
 \xi_i \downarrow & & \downarrow \overline{\xi}_i \\
 X & \xrightarrow{p} & \overline{X}
 \end{array}$$

(which, of course, one may construct by first forming the conical pushout (ξ'_i, c'_i) of each pair (ξ_i, c_i) and then the conical wide pushout (= co-intersection) of $(c'_i)_{i \in \text{ob}\mathcal{D}}$).

Step 3: We assume that the (discretely) $\text{ob}\mathcal{D}$ -indexed and 1-weighted cocone $(\overline{\xi}_i : P(Wi \otimes Di) \rightarrow \overline{X})_{i \in \text{ob}\mathcal{D}}$ has an order-universal P -extension $(\alpha_i : Wi \otimes Di \rightarrow A)_{i \in \text{ob}\mathcal{D}}, A, q_o : \overline{X} \rightarrow PA$.

We set $q := q_o \cdot p : X \rightarrow PA$ and $\alpha_i^u := \alpha_i \cdot \lambda_i^u : Di \rightarrow A$ for all $i \in \text{ob}\mathcal{D}, u \in Wi$, and prove:

Proposition 5.8. *Under the assumptions of Steps 1–3, and when (the ordinary functor) P is faithful, one obtains a W -weighted cocone $\alpha = (\alpha_i^u)_{i,u}$ which, together with q , gives an order-universal P -extension of the given W -weighted cocone ξ .*

PROOF: Clearly, from $\lambda_i^u \leq \lambda_i^{u'}$ one obtains $\alpha_i^u \leq \alpha_i^{u'}$ for all $u \leq u' \in Wi, i \in \text{ob}\mathcal{D}$. Also, from $\xi_j^v \cdot PDd = \xi_i^{Wd(v)}$ one obtains $P(\alpha_j^v \cdot Dd) = P(\alpha_i^{Wd(v)})$ and then, when (the ordinary functor) P is faithful, $\alpha_j^v \cdot Dd = \alpha_i^{Wd(v)}$ for all $d : i \rightarrow j$ in \mathcal{D} and $v \in Wj$. Hence, α is a W -weighted cocone, obviously satisfying $P\alpha = q \cdot \xi$.

Given any W -weighted cocone $\beta: D \rightarrow B$ in \mathcal{A} and an \mathcal{X} -morphism $f: X \rightarrow PB$ with $P\beta = f \cdot \xi$, we consider the $\text{ob}\mathcal{D}$ -indexed cocone $(\beta_i)_i$ with $\beta_i \cdot \lambda_i^u = \beta_i^u$ for all $u \in Wi, i \in \text{ob}\mathcal{D}$. Then, from $P\beta_i \cdot c_i \cdot \kappa_i^u = f \cdot \xi_i \cdot \kappa_i^u$ for all u one derives $P\beta_i \cdot c_i = f \cdot \xi_i$ for all i . The generalized pushout now gives an \mathcal{X} -morphism $g: \overline{X} \rightarrow PB$ with $g \cdot p = f$ and $g \cdot \xi_i = P\beta_i$ for all i . Order-universality of the discrete cocone $(\alpha_i)_i$ together with q_o finally produces an \mathcal{A} -morphism $t: A \rightarrow B$ with $Pt \cdot q_o = g$ and $t \cdot \alpha_i = \beta_i$ for all $i \in \text{ob}\mathcal{D}$, from which one easily deduces $Pt \cdot q = f$ and $t \cdot \alpha = \beta$.

To show that (α, A, q) is order- P -epic, we consider \mathcal{A} -morphisms $r, s: A \rightarrow B$ with $Pr \cdot q \leq Ps \cdot q, r \cdot \alpha \leq s \cdot \alpha$. The latter inequality gives $r \cdot \alpha_i \leq s \cdot \alpha_i$ for every $i \in \text{ob}\mathcal{D}$ since the cocone $(\lambda_i^u)_u$ of the tensor product $Wi \otimes Di$ is order-epic, while the first inequality and the cocone $(\kappa_i^u)_u$ of the tensor product $Wi \otimes PDi$ being order-epic give $Pr \cdot q_o \cdot \overline{\xi_i} \leq Ps \cdot q_o \cdot \overline{\xi_i}$. Since also the conical generalized pushout is order-epic, with $Pr \cdot q_o \cdot p \leq Ps \cdot q_o \cdot p$ one obtains $Pr \cdot q_o \leq Ps \cdot q_o$. In conjunction with $r \cdot \alpha_i \leq s \cdot \alpha_i$ for every $i \in \text{ob}\mathcal{D}$ one can finally conclude $r \leq s$ since the order-universal P -extension $((\alpha_i)_{i \in \text{ob}\mathcal{D}}, A, q_o)$ is order- P -epic. \square

Remarks 5.9. (1) For any $A \in \text{ob}\mathcal{A}, W \in \text{ob}\mathcal{P}os$, such that the respective tensor products in \mathcal{A} and \mathcal{X} exist, we call the canonical morphism $c: W \otimes PA \rightarrow P(W \otimes A)$ a *tensor comparison morphism*. In order to perform Step 2 it suffices that \mathcal{X} has *conical generalized pushouts of tensor comparison morphisms*; more precisely, the conical pushout of a tensor comparison morphism along any morphism exists in \mathcal{X} , and the conical wide pushout of any family of such pushouts exists as well.

(2) If \mathcal{A} has tensor products preserved by P , then the needed tensor products and pushouts in \mathcal{X} as described in (1) trivially exist and are conical since then, by definition of preservation, all tensor comparison morphisms are isomorphisms, so that by putting $\overline{X} = X, \overline{\xi_i} = \xi_i \cdot c_i^{-1}, p = 1_X$ one obtains the needed generalized pushout diagram of Step 2.

(3) While, when tensor products exist in \mathcal{X} , Corollary 5.4 guarantees their existence also in \mathcal{A} if P admits order-universal P -extensions of weighted cocones over diagrams of shape 1, these will generally not be preserved by P , even when P is strongly order-solid. For instance, let A be the 2-chain $2 = \{0 < 1\}$, and let W be the discrete 2-element poset, thus $W \otimes A$ is just a conical copower. In $\mathcal{S}\mathcal{L}at$, it is the diamond poset, but in $\mathcal{P}os$ it is just the disjoint union of two copies of 2. Hence, the strongly order-solid functor V of Example 3.4 does not preserve tensor products.

(4) If one tightens the condition of Proposition 5.8 that P be faithful to P being order-faithful, then the construction leads us more generally from an (op)lax cocone ξ to an (op)lax cocone α , as the beginning of the proof of the proposition shows. (Recall that an (op)lax W -weighted cocone $\alpha: D \rightarrow A$ is given by an (op)lax natural transformation $\alpha: W \rightarrow \mathcal{A}(D-, A)$, so that the identities $\alpha_j^v \cdot Dd = \alpha_i^{Wd(v)}$ of Remarks 4.1 (1) get traded for the inequalities $\alpha_j^v \cdot Dd \leq \alpha_i^{Wd(v)}$ (“ \geq ” in the op-lax case).) Consequently, with Propositions 5.1, 5.8, the construction

leads from *(op)lax colimits* (= universal (op)lax cocones) in \mathcal{X} to (op)lax colimits in \mathcal{A} when P is order-faithful.

With the Remarks 5.9 we obtain from Proposition 5.8 the following corollary:

Corollary 5.10. *For an ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$ and any ordered category \mathcal{D} , all weighted cocones $\xi: PD \rightarrow X$ with $D: \mathcal{D} \rightarrow \mathcal{A}$ have order-universal P -extensions if*

- (a) *all 1-weighted and discrete $\text{ob}\mathcal{D}$ -indexed cocones $(\xi_i: PD_i \rightarrow X)_i$ have order-universal P -extensions;*
- (b) *\mathcal{A} has tensor products and P preserves them;*
- (c) *as an ordinary functor, P is faithful.*

We can now combine some of the previous statements and formulate a characterization of order-solid functors:

Theorem 5.11. *For the ordered functor $P: \mathcal{A} \rightarrow \mathcal{X}$, assume that \mathcal{X} has all tensor products $W \otimes PD$ (with W in $\mathcal{P}\text{os}$ and D in \mathcal{A}), as well as conical generalized pushouts of arbitrary families of tensor comparison morphisms. Then P is order-solid if and only if \mathcal{A} is tensored and P admits order-universal P -extensions for all 1-weighted cocones of discrete shape. The assumption on \mathcal{X} is particularly satisfied when \mathcal{A} is tensored and P preserves tensor products.*

PROOF: When P is order-solid, by Proposition 5.1 (1), the existence of tensor products of the form $W \otimes PD$ in \mathcal{X} is sufficient to make \mathcal{A} tensored; also, trivially, the specified weighted cocones have order-universal P -extensions. Conversely, the existence of the specified order-universal P -extensions suffices to make the ordinary functor P solid and, hence, faithful. With our assumptions on \mathcal{X} , Proposition 5.8 now guarantees that P is order-solid.

The additional claim follows from Remark 5.9 (2). □

REFERENCES

- [1] Adámek J., Herrlich H., Strecker G. E., *Abstract and Concrete Categories*, Pure and Applied Mathematics (New York), A Wiley-Interscience Publication, John Wiley & Sons, New York, 1990.
- [2] Adámek J., Rosický J., Vitale E. M., *Algebraic Theories*, Cambridge Tracts in Mathematics, 184, Cambridge University Press, Cambridge, 2011.
- [3] Adámek J., Sousa L., *KZ-monadic categories and their logic*, Theory Appl. Categ. **32** (2017), paper no. 10, 338–379.
- [4] Adámek J., Sousa L., Velebil J., *Kan injectivity in order-enriched categories*, Math. Structures Comput. Sci. **25** (2015), no. 1, 6–45.
- [5] Anghel C., *Factorizations and Initiality in Enriched Categories*, Doctoral Dissertation, Fernuniversität, Hagen, 1987.
- [6] Anghel C., *Semi-initial and semi-final V-functors*, Comm. Algebra **18** (1990), no. 1, 135–181.
- [7] Anghel C., *Lifting properties of V-functors*, Comm. Algebra **18** (1990), no. 1, 183–192.
- [8] Antoine P., *Étude élémentaire des catégories d'ensembles structurés*, Bull. Soc. Math. Belg. **18** (1966), 142–164, 387–414 (French).

- [9] Börger R., Tholen W., *Cantors Diagonalprinzip für Kategorien*, Math. Z. **160** (1978), no. 2, 135–138 (German).
- [10] Börger R., Tholen W., *Total categories and solid functors*, Canad. J. Math. **42** (1990), no. 2, 213–229.
- [11] Brümmer G. C. L., *A Categorical Study of Initiality in Uniform Topology*, Ph.D. Thesis, University of Cape Town, Cape Town, 1971.
- [12] Carvalho M., Sousa L., *Order-preserving reflectors and injectivity*, Topology Appl. **158** (2011), no. 17, 2408–2422.
- [13] Carvalho M., Sousa L., *On Kan-injectivity of locales and spaces*, Appl. Categorical Structures **25** (2017), no. 1, 83–104.
- [14] Čech E., *Topological Spaces*, Academia, Prague, 1966.
- [15] Dubuc E., *Adjoint triangles*, Reports of the Midwest Category Seminar, Springer, Berlin, 1968, pages 69–91.
- [16] Fritz T., *Resource convertibility and ordered commutative monoids*, Math. Structures Comput. Sci. **27** (2017), no. 6, 850–938.
- [17] Herrlich H., *Topological functors*, General Topology and Appl. **4** (1974), 125–142.
- [18] Hoffmann R.-E., *Die kategorielle Auffassung der Initial- und Finaltopologie*, Doctoral Dissertation, Ruhr-Universität, Bochum, 1972 (German).
- [19] Hoffmann R.-E., *Semi-identifying lifts and a generalization of the dual theorem for topological functors*, Math. Nachr. **74** (1976), 295–307.
- [20] Hofmann D. (ed.), Seal G. J. (ed.), Tholen W. (ed.), *Monoidal Topology*, Encyclopedia of Mathematics and Its Applications, 153, Cambridge University Press, Cambridge, 2014.
- [21] Hofmann D., Sousa L., *Aspects of algebraic algebras*, Log. Methods Comput. Sci. **13** (2017), no. 3, paper no. 4, 25 pages.
- [22] Hong Y. H., *Studies on Categories of Universal Topological Algebras*, Ph.D. Thesis, McMaster University, Hamilton, 1974.
- [23] Hušek M., *S-categories*, Comment. Math. Univ. Carolinae **5** (1964), 37–46.
- [24] Jameson G., *Ordered Linear Spaces*, Lecture Notes in Mathematics, 141, Springer, Berlin, 1970.
- [25] Kelly G. M., *Basic Concepts of Enriched Category Theory*, London Mathematical Society Lecture Note Series, 64, Cambridge University Press, Cambridge, 1982.
- [26] Kurz A., Velebil J., *Quasivarieties and varieties of ordered algebras: regularity and exactness*, Math. Structures Comput. Sci. **27** (2017), no. 7, 1153–1194.
- [27] Lawvere F. W., *Functorial Semantics of Algebraic Theories*, Ph.D. Thesis, Columbia University, New York, 1963.
- [28] Linton F. E. J., *Some aspects of equational categories*, Proc. of the Conf. Categorical Algebra, La Jolla, 1965, Springer, New York, 1966, pages 84–94.
- [29] MacLane S., *Categories for the Working Mathematician*, Graduate Texts in Mathematics, 5, Springer, Berlin, 1971.
- [30] Manes E. G., *A pullback theorem for triples in a lattice fibering with applications to algebra and analysis*, Algebra Universalis **2** (1972), 7–17.
- [31] Picado J., Pultr A., *Frames and Locales: Topology without Points*, Frontiers in Mathematics, Birkhäuser/Springer Basel AG, Basel, 2012.
- [32] Roberts J. E., *A characterization of initial functors*, J. Algebra **8** (1968), 181–193.
- [33] Schaefer H. H., *Topological Vector Spaces*, Graduate Texts in Mathematics, 3, Springer, New York, 1971.
- [34] Shukla W., *On Top Categories*, Ph.D. Thesis, Indian Institute of Technology, Kanpur, 1971.
- [35] Sousa L., *Solid hulls of concrete categories*, Appl. Categ. Structures **3** (1995), no. 2, 105–118.
- [36] Sousa L., *A calculus of lax fractions*, J. Pure Appl. Algebra **221** (2017), no. 2, 422–448.

- [37] Street R., Tholen W., Wischnewsky M., Wolff H., *Semitopological functors. III. Lifting of monads and adjoint functors*, J. Pure Appl. Algebra **16** (1980), no. 3, 299–314.
- [38] Taylor J. C., *Weak families of maps*, Canad. Math. Bull. **8** (1965), 771–781.
- [39] Tholen W., *M-functors*, Nordwestdeutsches Kategorienseminar, Tagung, Bremen, 1976, Math.-Arbeitspapiere, No. 7, Teil A: Math. Forschungspapiere, Universität Bremen, Bremen, 1976, pages 178–185.
- [40] Tholen W., *On Wyler's taut lift theorem*, General Topology and Appl. **8** (1978), no. 2, 197–206.
- [41] Tholen W., *Semitopological functors I*, J. Pure Appl. Algebra **15** (1979), no. 1, 53–73.
- [42] Tholen W., *Note on total categories*, Bull. Austral. Math. Soc. **21** (1980), no. 2, 169–173.
- [43] Tholen W., Wischnewsky M. B., *Semitopological functors. II. External characterizations*, J. Pure Appl. Algebra **15** (1979), no. 1, 75–92.
- [44] Trnková V., *Automata and categories*, Mathematical Foundations of Computer Science 1975, Fourth Sympos., Mariánské Lázně, 1975, Lecture Notes in Computer Science, 32, Springer, Berlin, 1975, 138–152.
- [45] Wischnewsky M. B., *Partielle Algebren in Initialkategorien*, Math. Z. **127** (1972), 83–91 (German).
- [46] Wischnewsky M. B., *A lifting theorem for right adjoints*, Cahiers Topologie Géom. Différentielle **19** (1978), no. 2, 155–168.
- [47] Wyler O., *On the categories of general topology and topological algebra*, Arch. Math. (Basel) **22** (1971), 7–17.
- [48] Wyler O., *Top categories and categorical topology*, General Topology and Appl. **1** (1971), no. 1, 17–28.

L. Sousa:

CMUC, UNIVERSITY OF COIMBRA, DEPARTMENT OF MATHEMATICS, APARTADO 3008,
EC SANTA CRUZ 3001-501 COIMBRA, PORTUGAL

and

POLYTECHNIC INSTITUTE OF VISEU, AV. COR. JOSÉ MARIA VALE DE ANDRADE S/N,
CAMPUS POLITÉCNICO SANTA MARIA, 3504-510 VISEU, PORTUGAL

E-mail: sousa@estv.ipv.pt

W. Tholen:

DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, 4700 KEELE ST,
TORONTO, ONTARIO, M3J 1P3, CANADA

E-mail: tholen@mathstat.yorku.ca

(Received December 28, 2018, revised March 12, 2019)