

## Normal structure and weakly normal structure of Orlicz spaces

SHUTAO CHEN, YANZHENG DUAN

*Abstract.* Every Orlicz space equipped with Orlicz norm has weak sum property, therefore, it has weakly normal structure and fixed point property. A criterion of sum property also of normal structure for such spaces is given as well, which shows that every Orlicz space has isonormal structure.

*Keywords:* Orlicz space, normal structure

*Classification:* 46B20

### Introduction.

T. Landes [4] shows that, under certain conditions, an Orlicz sequence space with Luxemburg norm has normal or weakly normal structure iff it is separable. For Orlicz spaces with Orlicz norm, we will discover that the results are much different.

We begin with some notations. A sequence  $(x_n)$  of a Banach space  $X$  is called limit affine, if the limit  $\lambda(x) := \lim_n \|x_n - x\| > 0$  exists for every  $x \in \text{conv}(x_n)$ , and  $\lambda$  is an affine function on  $\text{conv}(x_n)$ .  $(x_n)$  is called growing, if  $\lambda(x_n) \leq \lambda(x_{n+1})$  for all  $n \in N$ .  $X$  is said to have sum property, if it contains no growing limit affine sequence.  $X$  is said to have weak sum property, if it contains no growing weakly converging limit affine sequence.  $X$  is said to have normal structure, if it contains no limit affine sequence  $(x_n)$  with  $\lambda(x_n) = \lambda(x_{n+1}) > 0$  for all  $n \in N$ .  $X$  is said to have weakly normal structure, if it contains no weakly converging limit affine sequence  $(x_n)$  with  $\lambda(x_n) = \lambda(x_{n+1}) > 0$  for all  $n \in N$ .  $X$  is said to have isonormal structure, if it is isomorphic to a Banach space with normal structure.  $X$  is said to have fixed point property if every nonexpansive selfmapping on a weakly compact convex subset of  $X$  has a fixed point.

It is well known that sum property  $\Rightarrow$  normal structure and that weak sum property  $\Rightarrow$  weakly normal structure  $\Rightarrow$  fixed point property.

Throughout this paper, we always denote by  $(G, \Sigma, \mu)$  a complete, nonatomic, finite measure space. We say  $M : R \rightarrow R^+$  to be an  $N$ -function, if it is a continuous, convex, even function satisfying  $M(u) = 0$ , iff  $u = 0$  and  $M(u)/u \rightarrow 0$  (resp.  $\infty$ ) as  $u \rightarrow 0$  (resp.  $\infty$ ). If  $M(u)$  is an  $N$ -function, then we denote by  $p(u)$  its right-hand derivative and by  $N(v)$  the conjugate  $N$ -function of  $M(u)$ , i.e.,  $N(v) := \max_u \{uv - M(u)\}$ .

---

\*Supported by the Special Scientific Research Fund of China for University Doctoral Instruction.

Let  $M$  be an  $N$ -function. For every  $\mu$ -measurable function  $x : G \rightarrow R$  we define  $\varrho_M(x) = \int_G M(x(t)) d\mu$ , and

$$\begin{aligned}
 (1) \quad & L_M = \{x : \varrho_M(\beta x) < \infty \text{ for some } \beta > 0\}, \\
 & E_M = \{x : \varrho_M(\beta x) < \infty \text{ for all } \beta > 0\}, \\
 & \|x\| = \|x\|_M = \inf_{k>0} [1 + \varrho_M(kx)]/k, \quad x \in L_M.
 \end{aligned}$$

Then the Orlicz space  $(L_M, \|\cdot\|)$  and its subspace  $(E_M, \|\cdot\|)$  are Banach spaces.

**Main results.**

**Lemma 1.** *Suppose  $x_n \in L_M, \|x_n\| \leq K, n \in N$  and  $x_n(t) \rightarrow x(t)$  in measure as  $n \rightarrow \infty$ , then  $x \in L_M$ .*

PROOF: Since  $\|x_n/K\| \leq 1$ , by [1],  $\varrho_M(x_n/K) \leq \|x_n/K\| \leq 1, n \in N$ . Without loss of generality, we may assume  $x_n(t)$   $\mu$ -a.e. on  $G$  (pass a subsequence, if necessary), then, by Fatou’s lemma,  $\varrho_M(x/K) \leq \liminf_{n \rightarrow \infty} \varrho_M(x_n/K) \leq 1$ , i.e.,  $x \in L_M$ .  $\square$

**Lemma 2.** *If  $x_n \rightarrow 0$  weakly in  $L_M$  and  $x_n(t) \rightarrow y(t)$  in measure, then  $y = 0$ .*

PROOF: Again, we may assume  $x_n(t) \rightarrow y(t)$   $\mu$ -a.e. on  $G$ . Let  $F = \{t \in G; y(t) \neq 0\}$ . If  $\mu^F > 0$ , then there exists  $E \in \Sigma$  with  $\mu E < \mu F$  and  $x_n(t) \rightarrow y(t)$  uniformly on  $G \setminus E$ . Define  $v(t) = \text{signy}(t)\chi_{F \setminus E}(t)$ , then  $v \in L_M^*$  and  $\langle v, x_n \rangle = \int_{F \setminus E} v(t)x_n(t) d\mu \rightarrow \int_{F \setminus E} |y(t)| d\mu > 0$  contradicting the hypothesis  $x_n \rightarrow 0$  weakly.  $\square$

We say that an interval  $[a, b]$  is a structural affine interval of the  $N$ -function  $M$ , if  $M$  is affine on  $[a, b]$  and it is neither affine on  $[a - \varepsilon, b]$  nor on  $[a, b + \varepsilon]$  for any  $\varepsilon > 0$ .

**Theorem 1.** *For any  $N$ -function  $M, L_M$  has weak sum property, therefore, it has weakly normal structure.*

**Theorem 2.** *The following are equivalent,*

- (i)  $L_M$  has sum property,
- (ii)  $L_M$  has normal structure,
- (iii) *there exist  $a > 0, C > 1$  such that for any structural affine interval  $[u, v]$  of  $M$  with  $u \geq a$ , we have  $v/u \leq C$ .*

PROOF OF THEOREMS 1 AND 2: For any limit affine sequence  $(x_n)$  in  $L_M$  with  $x_i \neq x_j$  whenever  $i \neq j$ , by [1], the “inf” in (1) is attainable for all  $x \neq 0$ . Therefore, for all  $i \neq j$ , we may find  $k_{ij} > 0$  such that

$$(2) \quad \|x_i - x_j\| = [1 + \varrho_M(k_{ij}(x_i - x_j))]/k_{ij}.$$

First we show that there exists a subsequence  $N_1$  of  $N$  such that for any  $j \in N_1, \{k_{ij}\}_{i \in N_1}$  is bounded. Indeed, if  $\{k_{ij}\}_i$  is bounded for all  $j \in N$ , then we let  $N_1 = N$ . Otherwise, there exist some  $m \in N$  and a subsequence  $I$  of  $N$  such that

$k_{im} \rightarrow \infty$  as  $i(\in I) \rightarrow \infty$ . Hence, for any  $\sigma > 0$ , if we define  $G_i = \{t \in G : |x_i(t) - x_m(t)| > \sigma\}$ , then by (2), we have

$$\|x_i - x_m\| > \varrho_M(|x_i - x_m|^{k_{im}})/k_{im} \geq [M(\sigma k_{im})/k_{im}]\mu G_i.$$

Since  $\lambda(x_m) < \infty$  and  $M(u)/u \rightarrow \infty$  as  $u \rightarrow \infty$ , we must have  $\mu G_i \rightarrow 0$  as  $i(\in I) \rightarrow \infty$ . This shows that  $\{x_i\}_{i \in I}$  converges to  $x_m$  in measure. We may assume that  $I$  does not contain  $m$ . We claim that  $N_1 = I$  satisfies our requirement. In fact, if  $k_{ij} \rightarrow \infty$  as  $i(\in I_1) \rightarrow \infty$  for some  $j \in I$  and some subsequence  $I$ , of  $I$ , then in the same way we can show that  $x_i \rightarrow x_j$  in measure as  $i(\in I_1) \rightarrow \infty$ . This is impossible since  $x_j \neq x_m$ .

By the diagonal method, we can find a subsequence  $N_2$  of  $N_1$  such that  $k_{ij} \rightarrow k_j < \infty$  as  $i(\in N_2) \rightarrow \infty$  for each  $j \in N_1$ . We claim that  $k_j \rightarrow \infty$  as  $j(\in N_2) \rightarrow \infty$ . In fact, if this is not true, then  $N_2$  contains a subsequence  $N_3$  such that  $k_j \rightarrow k < \infty$  as  $j(\in N_3) \rightarrow \infty$ . By (1) and (2), for all  $n, i, j \in N_3$ ,  $n \neq i, j$ ,

$$\begin{aligned} & \|x_n - x_i\| + \|x_n - x_j\| - \|2x_n - x_i - x_j\| \geq \\ & \geq [1 + \varrho_M(k_{ni}(x_n - x_i))]/k_{ni} + [1 + \varrho_M(k_{nj}(x_n - x_j))]/k_{nj} - \\ (3) \quad & - [1 + \varrho_M((2x_n - x_i - x_j)k_{ni}k_{nj}/(k_{ni} + k_{nj}))](k_{ni} + k_{nj})/k_{ni}k_{nj} = \\ & = \int_G [M((x_n(t) - x_i(t))k_{ni})/k_{ni} + M((x_n(t) - x_j(t))k_{nj})/k_{nj} - \\ & - M((2x_n(t) - x_i(t) - x_j(t))k_{ni}k_{nj}/(k_{ni} + k_{nj}))](k_{ni} + k_{nj})/k_{ni}k_{nj} d\mu. \end{aligned}$$

Denote the last integrand in (3) by  $f_{nij}(t)$ , then by the convexity of  $M$ ,  $f_{nij}(t) \geq 0$  for all  $t \in G$ . Since  $\lambda$  is affine on  $\text{conv}(x_k)$ , let  $n \rightarrow \infty$ , by (3),  $\int_G f_{nij}(t) d\mu \rightarrow 0$ , therefore,  $f_{nij}(t) \rightarrow 0$  in measure. By the diagonal method, we can choose a subsequence  $N_4$  of  $N_3$  such that  $f_{nij}(t) \rightarrow 0$   $\mu$ -a.e. on  $G$  as  $n(\in N_4) \rightarrow \infty$  for all  $i, j \in N_3$ .

For each  $t \in G$ , choose a subsequence  $\{n_\tau = n_\tau(t)\}$  of  $N_4$  such that

$$(*) \quad |v(t)| = \liminf_{n \in N_4} |x_n(t)|, \quad \lim_{\tau} x_{n_\tau}(t) = v(t),$$

then by Fatou's lemma,  $|v(t)| < \infty$   $\mu$ -a.e. on  $G$  (one may prove this analogously as in Lemma 1). Let  $\tau \rightarrow \infty$ , by the continuity of  $M$ ,

$$\begin{aligned} (4) \quad 0 &= \lim_{\tau} f_{n_\tau i j}(t) = \\ &= M((v(t) - x_i(t))k_i)/k_i + M((v(t) - x_j(t))k_j)/k_j - \\ & - M((2v(t) - x_i(t) - x_j(t))k_i k_j / (k_i + k_j))(k_i + k_j)/k_i k_j \end{aligned}$$

$\mu$ -a.e. on  $G$ . Since for  $\mu$ -a.e.  $t \in G$ , (4) holds for all  $i, j \in N_3$ , by replacing  $j$  by  $n_\tau$  in (4) and taking  $\tau \rightarrow \infty$ , for each  $t \in G$ , we have

$$\begin{aligned} (5) \quad & M((v(t) - x_i(t))k_i)/k_i = \\ & = M((v(t) - x_i(t))k_i k / (k_i + k))(k_i + k)/k_i k \end{aligned}$$

$\mu$ -a.e. on  $G$ . Since  $0 < k/(k_i + k) < 1$ , and  $u \neq 0, 0 < \alpha < 1$  implies  $M(\alpha u) < \alpha M(u)$ , for all  $t \in G$  satisfying  $v(t) \neq x_i(t)$ , we have

$$M((v(t) - x_i(t))k_i k / (k_i + k))(k_i + k) / k_i k < M((v(t) - x_i(t))k_i) / k_i.$$

Since this inequality contradicts (5), we have  $x_i(t) = v(t)$   $\mu$ -a.e. on  $G$  for all  $i \in N_3$ , which contradicts the assumption  $x_i \neq x_j$  whenever  $i \neq j$ .

Now, we prove (iii)  $\Rightarrow$  (i) in Theorem 2. If  $L_M$  does not have sum property, then there exists a growing limit affine sequence. Without loss of generality, we may assume that every two points in the sequence are different. By the above discussion, it contains a subsequence  $(x_n)$  satisfying  $k_{ij} \rightarrow k_j < \infty$  as  $i \rightarrow \infty$  and  $k_j \rightarrow \infty$  as  $j \rightarrow \infty$ , where  $k_{ij}$  satisfies (2),  $i, j \in N$ . Since  $M(u)/u \rightarrow \infty$  as  $u \rightarrow \infty$ , for the constant  $a > 0$  in (iii), we can find  $b > a$  such that  $M(\frac{1}{2}(a + b)) < \frac{1}{2}[M(a) + M(b)]$ . Since  $M$  is convex, by (iii)

$$(6) \quad M(\alpha u + (1 - \alpha)v) < \alpha M(u) + (1 - \alpha)M(v)$$

for all  $0 < \alpha < 1$  and all  $u \leq a, v \geq b$  or  $u \geq a, v \geq Cu$ . If we define  $v(t)$  as in (\*), then by (4) and (6), for  $\mu$ -a.e.  $t \in G$ , if  $k_i|v(t) - x_i(t)| \leq a$ , then  $k_j|v(t) - x_j(t)| \leq b$ ; if  $k_i|v(t) - x_i(t)| > a$ , then  $k_j|v(t) - x_j(t)| \leq Ck_i|v(t) - x_i(t)|$ . Therefore, for  $\mu$ -a.e.  $t \in G$ ,

$$(7) \quad k_j|v(t) - x_j(t)| \leq \max\{b, Ck_i|v(t) - x_i(t)|\} := u_i(t).$$

By (2) and Fatou's lemma, we have

$$(8) \quad \lambda(x_j) \geq [1 + \varrho_M(k_j(v - x_j))]/k_j \geq \|v - x_j\|.$$

Thus,  $v - x_j \in L_M$ , therefore,  $u_i \in L_M$ . Since  $\lambda > 0$ , we have  $\liminf \|v - x_j\| := \tau > 0$ . It follows from (7) that

$$k_j = \|k_j(v - x_j)\|/\|v - x_j\| \leq \|u_i\|/\|v - x_j\|.$$

Let  $j \rightarrow \infty$ , we get a contradiction  $\infty \leq \|u_i\|/\tau < \infty$ .

Next, we turn to Theorem 1. If  $L_M$  does not have weak sum property, then by (3), there exists a weakly converging (to zero) limit affine sequence  $(x_n)$  with  $\|x_n\| \rightarrow 1$  and  $\lambda(x_n) \rightarrow 1$ . By the first part of the proof, passing a subsequence if necessary, we may assume  $k_{ij} \rightarrow k_j < \infty$  as  $i \rightarrow \infty$  and  $k_j \rightarrow \infty$  as  $j \rightarrow \infty$ , where  $k_{ij}$  satisfies (2). It follows from (8) that  $x_j \rightarrow v$  in measure (similarly verified as in the first part of the proof). Therefore, by Lemma 2,  $v = 0$ . We may also assume  $x_j \rightarrow 0$   $\mu$ -a.e. on  $G$ . We prove the theorem by showing  $\lim \lambda(x_j) \geq 4/3$  contradicting the assumption  $\lambda(x_j) \rightarrow 1$ .

For each  $j \in N$ , we choose a set  $G_j \in \Sigma$  such that  $x_j$  is bounded on  $G_j$  and

$$[1 + \varrho_M(k_j x_j \chi_{G_j})]/k_j > [1 + \varrho_M(k_j x_j)]/k_j - 1/k_j,$$

then by (1) and (8),

$$\begin{aligned} \lambda(x_j) &= [1 + \varrho_M(k_j x_j)]/k_j = \\ &= [1 + \varrho_M(k_j x_j \chi_{G_j})]/k_j + [1 + \varrho_M(k_j x_j \chi_{G \setminus G_j})]/k_j - 1/k_j > \\ &> [1 + \varrho_M(k_j x_j)]/k_j - 1/k_j + \|x_j \chi_{G \setminus G_j}\| - 1/k_j \geq \\ &\geq \|x_j\| + \|x_j \chi_{G \setminus G_j}\| - 2/k_j, \end{aligned}$$

i.e.

$$(9) \quad \|x_j \chi_{G \setminus G_j}\| < \lambda(x_j) - \|x_j\| + 2/k_j.$$

It follows that

$$(10) \quad \|x_j \chi_{G_j}\| \geq \|x_j\| - \|x_j \chi_{G \setminus G_j}\| > 2\|x_j\| - \lambda(x_j) - 2/k_j.$$

Since  $x_j$  is bounded on  $G_j$ , there exists  $\delta = \delta(j) > 0$  such that

$$(11) \quad \|x_j \chi_E\| < 1/k_j \quad \text{whenever } E \subset G_j \text{ and } \mu E < \delta.$$

Since  $x_i \rightarrow 0$   $\mu$ -a.e. on  $G$ , there exists  $F \in \Sigma$  with  $\mu F < \delta$  such that  $x_i \rightarrow 0$  uniformly on  $G \setminus F$ . Hence, there exists  $I = I(j) \in N$  such that for all  $i > I$ , we have

$$(12) \quad \|x_i \chi_{G \setminus F}\| < 1/k_j.$$

It follows that

$$(13) \quad \|x_i \chi_F\| \geq \|x_i\| - \|x_i \chi_{G \setminus F}\| > \|x_i\| - 1/k_j.$$

Hence, by (1), (2), (9)–(13),

$$\begin{aligned} \|x_i - x_j\| &= [1 + \varrho_M(k_{ij}(x_i - x_j) \chi_{G \setminus (G_j \setminus F)})]/k_{ij} + \\ &+ [1 + \varrho_M(k_{ij}(x_i - x_j) \chi_{G_j \setminus F})]/k_{ij} - 1/k_{ij} \geq \\ &\geq \|(x_i - x_j) \chi_{G \setminus (G_j \setminus F)}\| + \|(x_i - x_j) \chi_{G_j \setminus F}\| - 1/k_{ij} \geq \\ &\geq \|x_i \chi_{G \setminus (G_j \setminus F)}\| - \|x_j \chi_{G \setminus (G_j \setminus F)}\| + \\ &+ \|x_j \chi_{G_j \setminus F}\| - \|x_i \chi_{G_j \setminus F}\| - 1/k_{ij} = \\ &= \|x_i \chi_{G \setminus (G_j \setminus F)}\| - \|x_j \chi_{G \setminus G_j} + x_j \chi_{G_j \setminus F}\| + \\ &+ \|x_j \chi_{G_j} - x_j \chi_{G_j \setminus F}\| - \|x_i \chi_{G_j \setminus F}\| - 1/k_{ij} > \\ &> (\|x_i\| - 1/k_j) - (\lambda(x_j) - \|x_j\| + 2/k_j + 1/k_j) + \\ &+ (2\|x_j\| - \lambda(x_j) - 2/k_j - 1/k_j) - 1/k_j - 1/k_{ij} = \\ &= \|x_i\| + 3\|x_j\| - 2\lambda(x_j) - 8/k_j - 1/k_{ij}. \end{aligned}$$

Let  $i \rightarrow \infty$ , we have

$$\lambda(x_j) \geq 1 + 3\|x_j\| - 2\lambda(x_j) - 9/k_j.$$

Let  $j \rightarrow \infty$ , then  $\lim \lambda(x_j) \geq 4/3$ .

Finally, we prove (ii)  $\Rightarrow$  (iii) in Theorem 2. If (iii) does not hold, then there exist the sequences  $\{u_j\}, \{v_j\}$  such that  $M(u_1)\mu G > 1, u_{j+1} > 2^j u_j, v_j > 2^j u_j$  and  $p(u)$  is constant on  $[u_j, v_j], j \in N$ . By the first two assumptions, we can choose disjoint sets  $g_j \in \Sigma$  such that  $\mu G \setminus U_{j \in N} G_j > 0$  and

$$(14) \quad 2^{-j} = u_j p(u_j) \mu G_j = [M(u_j) + N(p(u_j))] \mu G_j$$

(the last equality holds by the special case of Young's inequality). Hence, we can find  $u_0$  large enough so that there is  $G_0$  satisfying

$$(15) \quad \sum_{j \in N} N(p(u_j)) \mu G_j + N(p(u_0)) \mu G_0 = 1.$$

Define

$$\begin{aligned} v &= \sum_{j \geq 0} p(u_j) \chi_{G_j}, \\ x_n &= u_0 \chi_{G_0} + \sum_{j \in N} v_j \chi_{G_j} + \sum_{j > n} u_j \chi_{G_j}, \end{aligned}$$

then by (15),  $\varrho_N(v) = 1$ , therefore,  $v \in L_M^*$  and  $\|v\| = 1$  (cf. [1]).

First we show that  $x_n \in E_M$  for any  $n \in N$ . Given arbitrary  $K > 1$ , choose  $J > n$  such that  $2^j > K$ , then  $v_j > 2^j u_j > K u_j > u_j$  for all  $j > J$ . Therefore

$$\begin{aligned} \sum_{j > J} M(K u_j) \mu G_j &= \sum_{j > J} [K u_j p(K u_j) - N(p(K u_j))] \mu G_j < \\ < \sum_{j > J} K u_j p(K u_j) \mu G_j &= \sum_{j > J} K u_j p(u_j) \mu G_j = K \sum_{j > J} 2^{-j} < \infty. \end{aligned}$$

This implies  $\varrho_M(K x_n) < \infty$ . Since  $K > 1$  is arbitrary, we have  $x_n \in E_M$ .

Let  $k_n = \|x_n\|$  and  $y_n = x_n/k_n$ , then  $y_n \in E_M$  and  $\|y_n\| = 1$ . By (1) and (15),

$$\begin{aligned} \|y_n\| \geq \langle v, y_n \rangle &= [u_0 p(u_0) \mu G_0 + \sum_{j \leq n} v_j p(u_j) \mu G_j + \sum_{j > n} u_j p(u_j) \mu G_j] / k_n = \\ &= [\varrho_N(v) + \varrho_M(k_n y_n)] / k_n \geq \|y_n\| = 1. \end{aligned}$$

Moreover, since

$$k_n = \|x_n\| \geq \langle v, x_n \rangle > \sum_{j \leq n} v_j p(u_j) \mu G_j \geq \sum_{j \leq n} 2^j u_j p(u_j) \mu G_j = n,$$

we have  $k_n \rightarrow \infty$  as  $n \rightarrow \infty$ .

We complete the proof by showing  $\lambda = 2$  on  $\text{conv}(y_n)$ . Indeed, for any  $y \in \text{conv}(y_n)$ , there exist  $\lambda_i \geq 0, \sum_{i \leq m} \lambda_i = 1$  such that  $y = \sum_{i \leq m} \lambda_i y_i$ . Since  $\langle v, y_n \rangle = 1$ , we have  $\langle v, y \rangle = \sum_{i \leq m} \lambda_i \langle v, y_i \rangle = 1$ . For any  $\varepsilon > 0$ , since  $y \in E_M$ , there exists  $I > m$  such that  $\|y \chi_F\| < \varepsilon$ , where  $F = U_{i > I} G_i$ . In view of  $x_n(t) \leq \max\{v_I, u_0\}$

on  $G \setminus F$  and  $k_n \rightarrow \infty$  as  $n \rightarrow \infty$ , we can find  $n_0 \in N$  such that  $\|y_n \chi_{G \setminus F}\| < \varepsilon$  for all  $n > n_0$ . Define  $v_0 = v \chi_{G \setminus F} - v \chi_F$ , then  $\|v_0\| = \|v\| = 1$  and for all  $n > n_0$ ,

$$\begin{aligned} 2 &\geq \|y\| + \|y_n\| \geq \|y - y_n\| \geq \langle v_0, y - y_n \rangle = \\ &= \langle v_0, y \chi_{G \setminus F} \rangle + \langle v_0, y \chi_F \rangle - \langle v_0, y_n \chi_{G \setminus F} \rangle - \langle v_0, y_n \chi_F \rangle = \\ &= \langle v, y \chi_{G \setminus F} \rangle - \langle v, y \chi_F \rangle - \langle v, y_n \chi_{G \setminus F} \rangle + \langle v, y_n \chi_F \rangle = \\ &= \langle v, y \rangle - 2\langle v, y \chi_F \rangle - 2\langle v, y_n \chi_{G \setminus F} \rangle + \langle v, y_n \rangle > \\ &> 1 - 2\|y \chi_F\| - 2\|y_n \chi_{G \setminus F}\| + 1 > 2 - 4\varepsilon, \end{aligned}$$

which shows that  $\lambda(y) = 2$ . □

**Theorem 3.**  $L_M$  has isonormal structure.

PROOF: If  $M$  is strictly convex, then the condition (iii) in Theorem 2 holds for all  $u \neq 0$  and all  $C \neq 1$ . Therefore,  $L_M$  has normal structure in this case. By [1], for any Orlicz function  $M$  and any  $\varepsilon > 0$ , we can construct a strictly convex Orlicz function  $H$  such that

$$\|x\|_M \leq \|x\|_H \leq (1 + \varepsilon)\|x\|_M$$

for all  $x \in L_M$ , which shows that  $L_M$  is isomorphic to  $L_H$ , i.e.  $L_M$  has isonormal structure. □

### REFERENCES

- [1] Wu Congxin, Wang Tingfu, Chen Shutao, Wang Yuwen, *Geometry of Orlicz spaces*, Harbin, 1986.
- [2] Wu Congxin, Wang Tingfu, *Orlicz spaces and their applications*, Harbin, 1983.
- [3] Landes T., *Permanence properties of normal structure*, Pacific J. Math. **110**, No. 1 (1984), 125–143.

HARBIN NORMAL UNIVERSITY, DEPARTMENT OF MATHEMATICS, HARBIN, CHINA

HARBIN INSTITUTE OF TECHNOLOGY, DEPARTMENT OF MATHEMATICS, HARBIN, CHINA

(Received September 13, 1990, revised February 19, 1991)