

## A note on splittable spaces

VLADIMIR V. TKACHUK

*Abstract.* A space  $X$  is splittable over a space  $Y$  (or splits over  $Y$ ) if for every  $A \subset X$  there exists a continuous map  $f : X \rightarrow Y$  with  $f^{-1}fA = A$ . We prove that any  $n$ -dimensional polyhedron splits over  $\mathbf{R}^{2n}$  but not necessarily over  $\mathbf{R}^{2n-2}$ . It is established that if a metrizable compact  $X$  splits over  $\mathbf{R}^n$ , then  $\dim X \leq n$ . An example of  $n$ -dimensional compact space which does not split over  $\mathbf{R}^{2n}$  is given.

*Keywords:* splittable, polyhedron, dimension

*Classification:* 54A25

The notion of splittability was introduced by A.V. Arhangel'skii [1]. A space  $X$  is splittable (or splits) over a space  $Y$  if for any  $A \subset X$  there exists a continuous map  $f : X \rightarrow Y$  such that  $f^{-1}fA = A$ . Many results were obtained by A.V. Arhangel'skii and D.B. Shakhmatov ([1]–[3]) on spaces splittable over  $\mathbf{R}^\omega$ . The author had also written a paper [4] on this topic.

Recently, A.V. Arhangel'skii had shown that a compact space  $X$  splits over  $\mathbf{R}$  iff it embeds in  $\mathbf{R}$ . He also proved that any 1-dimensional polyhedron splits over  $\mathbf{R}^2$  so that not every compact space splittable over  $\mathbf{R}^2$  embeds in  $\mathbf{R}^2$ . We prove here that any  $n$ -dimensional polyhedron splits over  $\mathbf{R}^{2n}$  but not necessarily over  $\mathbf{R}^{2n-2}$ . We establish also that there exists a compact space  $X_n \subset \mathbf{R}^{2n+1}$  with  $\dim X_n = n$  and not splittable over  $\mathbf{R}^{2n}$ . Another result is Corollary 8 answering Question 2 and 3 in [1].

**Notations and terminology.** All spaces under consideration are Tychonoff ones. Given two spaces  $X$  and  $Y$ , denote by  $C(X, Y)$  the set of all continuous functions from  $X$  to  $Y$ . The space  $\mathbf{R}$  is the real line with its usual topology and  $2 = \{0, 1\}$ . If  $x, y \in \mathbf{R}^n$  then  $[x, y] = \{tx + (1-t)y : 0 \leq t \leq 1\}$  is the segment connecting  $x$  and  $y$ ,  $|x - y|$  is its length. It is always clear from the context whether  $|\cdot|$  denotes cardinality of some set or length of a segment. Of course,  $[x, y] = \{tx + (1-t)y : 0 < t \leq 1\}$  and  $(x, y) = \{tx + (1-t)y : 0 < t < 1\}$ . The simplex in  $\mathbf{R}^n$  with vertices  $a_0, \dots, a_k$  will be denoted by  $[a_0, \dots, a_k]$ , then  $\langle T \rangle = \{t_0a_0 + \dots + t_ka_k : t_i > 0 \text{ for all } i \in k + 1 \text{ and } \sum_{i=0}^k t_i = 1\}$ . Let  $x \in \mathbf{R}^n$  and  $A \subset \mathbf{R}^n$ . Then  $\text{con}(x, A) = \bigcup\{[x, a] : a \in A\}$ .

Other notations are standard and can be found in [9].

**1. Lemma.** *Given any spaces  $X$  and  $Y$  and an infinite cardinal  $k$  with  $d(X) \leq k$ , ( $d$  is the density character),  $|Y| \leq 2^k$ , suppose that  $\bigcup\{X_s : s \in 2^k\} \subset X$ ,  $X_s \cap X_t = \emptyset$  if  $s \neq t$  and no  $X_s$  can be continuously injected into  $Y$ . Then  $X$  does not split over  $Y$ .*

PROOF: Clearly,  $|C(X, Y)| \leq 2^k$ , so let  $C(X, Y) = \{f_s : s < 2^k\}$ . For any  $s < 2^k$  there is an  $x_s \in X$  such that  $|f_s^{-1}f_s(x_s) \cap X_s| > 1$  because  $f \upharpoonright X$  is not an injection. Then the set  $A = \{x_s : s < 2^k\}$  witnesses non-splittability of  $X$  over  $Y$ .  $\square$

**2. Example.** For any natural  $n$  there exists an  $n$ -dimensional metrizable compact space  $X_n$  which does not split over  $\mathbf{R}^{2n}$ .

PROOF: Take any metrizable compact space  $Y_n$  with  $Y_n \not\hookrightarrow \mathbf{R}^{2n}$  and  $\dim Y_n = n$ . Then let  $X_n = 2^\omega \times Y$ . It is obvious that the family  $\{\{s\} \times Y_n : s \in 2^\omega\}$  satisfies the conditions of Lemma 1 for  $X = X_n$ ,  $k = \omega$  and  $Y = \mathbf{R}^{2n}$ . Hence  $X_n$  does not split over  $\mathbf{R}^{2n}$ .  $\square$

**3. Example.** There exists an  $n$ -dimensional compact polyhedron  $P_n$  which is not splittable over  $\mathbf{R}^{2n-2}$ .

PROOF: There exists an  $(n - 1)$ -dimensional compact polyhedron  $Y_n$  which does not embed in  $\mathbf{R}^{2n-2}$ . Then  $P = Y_n \times [0, 1]$  is what was required, because the family  $\{Y_n \times \{t\} : t \in [0, 1]\}$  satisfies the conditions of Lemma 1 for  $X = P_n$ ,  $k = \omega$  and  $Y = \mathbf{R}^{2n-2}$ .  $\square$

**4. Theorem.** Let  $P$  be an  $n$ -dimensional compact polyhedron. Then  $P$  splits over  $\mathbf{R}^{2n}$ .

PROOF: Denote by  $a_1, \dots, a_k$  the vertices of  $P$ . Let  $\{S_1, \dots, S_r\}$  be the set of all  $(n - 1)$ -dimensional simplexes from  $P$  and let  $\mu = \{T_1, \dots, T_m\}$  be some set of its  $n$ -dimensional ones. Take any hyperplane  $H \subset \mathbf{R}^{2n}$  and let  $b_1, \dots, b_k$  be some points generally positioned in  $H$ . Define a polyhedron  $Q_{n-1}$  in the following way: the vertices of  $Q_{n-1}$  are  $b_1, \dots, b_k$  and a simplex  $[b_{i_1}, \dots, b_{i_l}]$ ,  $l \leq n$  belongs to  $Q_{n-1}$  iff the simplex  $[a_{i_1}, \dots, a_{i_l}]$  belongs to  $P$ . If  $P_{n-1}$  is the union of all  $\leq (n - 1)$ -dimensional simplexes, then the simplicial map  $f : P_{n-1} \rightarrow Q_{n-1}$  defined by  $f(a_i) = b_i$  is a homeomorphism because  $H$  is isomorphic to  $\mathbf{R}^{2n-1}$ . Pick any  $D \in \mathbf{R}^{2n} \setminus H$ .  $\square$

**5. Lemma.** There exist  $m$  sets  $L_1, \dots, L_m$  and a continuous map

$$g = g(f, D) : P_{n-1} \cup T_1 \cup \dots \cup T_m \rightarrow \mathbf{R}^{2n}$$

with the following properties:

- (1)  $L_i$  is a subset of  $\mathbf{R}^{2n}_D$ , where the last set is the component of  $\mathbf{R}^{2n} \setminus H$  containing  $D$ ,  $i = 1, \dots, m$ ;
- (2)  $L_i$  is homeomorphic to  $\langle T_i \rangle$ ,  $i = 1, \dots, m$ ;
- (3)  $L_i \cap L_j = \{D\}$  if  $i \neq j$ ;
- (4)  $g \upharpoonright P_{n-1} = f$ ;
- (5)  $g \upharpoonright (P_{n-1} \cup T_i)$  is a homeomorphism onto  $Q_{n-1} \cup L_i$ .

PROOF OF THE LEMMA: Let  $R = \text{con}(D, Q_{n-1})$ ,  $R_i = \text{con}(D, f(S_i))$ ,  $i = 1, \dots, r$ . The set  $R$  being compact, there is a sphere (in  $\mathbf{R}^{2n}$ ) containing it. Pick any  $E \in \mathbf{R}^{2n}_D$  outside this sphere and not belonging to any of  $(2n - 1)$ -dimensional planes, spanned in  $\mathbf{R}^{2n}$  by some  $2n$  points from the set  $\{b_1, \dots, b_k, D\}$ .

We are going to construct a continuous function  $q : R \rightarrow [0, 1)$  such that

$$(6) \quad q^{-1}(0) = Q_{n-1} \cup \{D\};$$

$$(7) \quad \text{if } x \in R \setminus (Q_{n-1} \cup \{D\}) \text{ and } y \in (x, E) \cap R, \text{ then } q(x) < \frac{|x - y|}{|y - E|}.$$

To do that let  $W_i = \{x \in R : (x, E) \cap R_i \neq \emptyset\}$ . For each  $x \in R$  and each  $i$ , there is at most one point in  $(x, E) \cap R_i$ , for otherwise  $E$  would belong to the  $n$ -dimensional plane spanned by  $R_i$ . Put for  $x \in R_i$ :

$$r_i(x) = \frac{|x - y|}{|y - E|} \text{ where } y \in (x, E) \cap R_i, \text{ and } r_i(D) = 0.$$

Let us prove that  $\text{dom } r_i = W_i \cup \{D\}$  is a closed subset of  $R$ . It suffices to show that  $\text{dom } r_i \cap R_j$  is closed for each  $j$ . If  $R_i \cap R_j \cap H \neq \emptyset$  then, owing to the choice of  $E$ ,  $\text{dom } r_i \cap R_j = \{D\}$ . Suppose  $R_i \cap R_j \cap H = \emptyset$ , hence  $R_i \cap R_j = \{D\}$ . Let  $x \in R_j \setminus \text{dom } r_i$ . Then  $(x, E) \cap R_i = \emptyset$  and  $[x, E] \cap R_i = \emptyset$  as well. Since  $[x, E]$  is compact and  $R_i$  is closed, the distance between these sets is positive, say  $\varepsilon$ , and whenever  $|z - x| < \varepsilon$  then clearly  $[z, E] \cap R_i = \emptyset$ . Since  $r_i$  is continuous, there exists a continuous  $q_i : R \rightarrow [0, 1)$  with  $q^{-1}(0) = Q_{n-1} \cup \{D\}$  and  $q_i(x) < r_i(x)$  for all  $x \in W_i \setminus (Q_{n-1} \cup \{D\})$ . Now it suffices to put

$$q(x) = \min\{q_i(x) : i = 1, \dots, r\}.$$

Let  $M_i = \text{con}(D, f(T_i \setminus \langle T_i \rangle))$ ,  $i = 1, \dots, m$ . It is clear that  $M_i$  is homeomorphic to  $T_i$ . Define an injective continuous map  $s_i : M_i \rightarrow \mathbf{R}_D^{2n}$  in the following way: if  $x \in M_i$  then find the point  $y \in [x, E)$  with

$$\frac{|x - y|}{|y - E|} = \frac{q(x)}{i + 1}$$

and put  $s_i(x) = y$ .

Evidently,  $s_i$  is a homeomorphism. Let  $L_i = s_i(M_i \setminus f(T_i \setminus \langle T_i \rangle))$ . We are going to define the map  $g = g(f, D)$  and verify (1)–(5). Take any homeomorphism  $u_i : T_i \rightarrow M_i$  with  $u_i \upharpoonright (T_i \setminus \langle T_i \rangle) = f$ . Then let  $g(x)$  be equal to  $f(x)$  if  $x \in P_{n-1}$  and to  $s_i(u_i(x))$  for  $x \in T_i \setminus P_{n-1}$ ,  $i = 1, \dots, m$ .

Only (3) needs to be verified.

Let  $x \in M_i \setminus (\{D\} \cup Q_{n-1})$ ,  $y \in M_j \setminus (\{D\} \cup Q_{n-1})$ . If  $g(x) = g(y)$  then  $x, y$  and  $E$  are linearly dependent. We may assume without loss of generality that  $y \in [x, E]$ . There are two possibilities:  $x = y$ , and  $x \neq y$ .

If  $x = y$  then

$$\frac{|x - g(x)|}{|g(x) - E|} = \frac{q(x)}{i + 1} \quad \text{and} \quad \frac{|x - g(y)|}{|g(y) - E|} = \frac{q(y)}{j + 1} = \frac{q(x)}{j + 1} \neq \frac{q(x)}{i + 1},$$

so that  $g(x) \neq g(y)$ , which is a contradiction.

If  $x \neq y$  and  $x \in R_{t_1}, y \in R_{t_2}, R_{t_1} \cap R_{t_2} \cap Q_{n-1} \neq \emptyset$  then it is impossible that  $y \in [x, E]$  — a contradiction.

If  $R_{t_1} \cap R_{t_2} \cap Q_{n-1} = \emptyset$  then

$$\frac{|x - g(x)|}{|g(x) - E|} < \frac{|x - y|}{|y - E|}$$

and therefore  $g(x) \in [x, y]$  while  $g(y) \in [y, E]$  and  $g(x) \neq g(y)$  — a contradiction again and we established (3) together with our lemma.  $\square$

We have all we need to split  $P$  over  $\mathbf{R}^{2n}$ . Let  $A \subset P$ . Pick a point  $D_1 \in \mathbf{R}^{2n} \setminus (H \cup \mathbf{R}_D^{2n})$ . Let  $T_1, \dots, T_m, T_{m+1}, \dots, T_{m_1}$  be all  $n$ -dimensional simplexes of  $P$  numerated in such a way that  $A \cap \langle T_i \rangle \neq \emptyset, i = 1, \dots, m, (P \setminus A) \cap \langle T_i \rangle \neq \emptyset, i = m + 1, \dots, m_1$ . Using Lemma 5 find the sets  $L_1, \dots, L_{m_1}$  and maps  $g = g(f, D)$  and  $g_1 = g(f, D_1)$  such that

- (8)  $L_i \subset \mathbf{R}_D^{2n}, i = 1, \dots, m, L_i \subset \mathbf{R}_{D_1}^{2n}, i = m + 1, \dots, m_1;$
- (9)  $L_i$  is homeomorphic to  $\langle T_i \rangle, i = 1, \dots, m_1;$
- (10)  $L_i \cap L_j = \{D\}, i \neq j, j \in 1, \dots, m;$
- (11)  $L_i \cap L_j = \{D\}, i = j, i, j \in m + 1, \dots, m_1;$
- (12)  $g \upharpoonright P_{n-1} = g_1 \upharpoonright P_{n-1} = f;$
- (13)  $g \upharpoonright (P_{n-1} \cup T_i)$  is a homeomorphism onto  $Q_{n-1} \cup L_i, i = 1, \dots, m;$
- (14)  $g \upharpoonright (P_{n-1} \cup T_i)$  is a homeomorphism onto  $Q_{n-1} \cup L_i, i = m + 1, \dots, m_1;$

Pick some points  $c_1, \dots, c_{m_1}$  with  $c_i \in A \cap \langle T_i \rangle, i = 1, \dots, m, c_i \in (P \setminus A) \cap \langle T_i \rangle, i = m + 1, \dots, m_1$  and the points  $d_1, \dots, d_{m_1}$  with  $g(d_i) = D, i = 1, \dots, m, g_1(d_i) = D_1, i = m + 1, \dots, m_1$  (observe that automatically  $d_i \in \langle T_i \rangle$  for each  $i$ ). Let  $G = g \cup g_1$ . Then  $G$  is a continuous map,  $G : P \rightarrow \mathbf{R}^{2n}$ . There exists a homeomorphism  $h : P \rightarrow P$  with  $h \upharpoonright P_{n-1} = \text{id}_{P_{n-1}}$  and  $h(c_i) = d_i, i = 1, \dots, m_1$ . The map  $F = G \circ h$  separates  $A$  from  $P \setminus A$ , because  $|F^{-1}(x)| = 1$  if  $x \notin \{D, D_1\}$  and  $F^{-1}(D) = \{c_1, \dots, c_m\} \subset A, F^{-1}(D_1) = \{c_{m+1}, \dots, c_{m_1}\} \subset P \setminus A$ , and our theorem is proved.  $\square$

**6. Proposition.** *Let  $X$  be a compact space and  $X = X_1 \cup X_2$  where  $X_1 \cap X_2 = \emptyset$  and any compact  $K \subset X_i$  is scattered ( $i = 1, 2$ ). Assume that  $X$  splits over a space  $Y$  with  $\dim Y \leq n$ . Then  $\dim X \leq n$ .*

PROOF: Take a continuous  $f : X \rightarrow Y$  with  $f^{-1}f(X_1) = X_1$ . If  $y \in Y$  then  $f^{-1}(y)$  is a compact subset of some  $X_i$  ( $i = 1, 2$ ) and is thus scattered. Hence  $\dim f^{-1}(y) = 0$  for every  $y \in f(X)$ . But  $\dim X \leq \dim f(X) + \dim f \leq n$  [10] and the proof is over.  $\square$

**7. Corollary.** *If a compact space  $X$  is splittable over  $\mathbf{R}^n$ , then  $\dim X \leq n$ .*

PROOF: The space  $X$  must be metrizable [3]. It is widely known (see e.g. [5]) that metrizable compact spaces satisfy the assumptions of Proposition 6, so our proof is over.  $\square$

This corollary answers Questions 2 and 3 in [1].

**8. Corollary ( $ACP^\#$ ).** *If  $X$  is a compact space splittable over a space  $Y$  then  $\dim X \leq \dim Y$ . (The definition of  $ACP^\#$  can be found in [5]).*

**9. Corollary.** *If  $X$  is a metrizable compact space splittable over a space  $Y$  then  $\dim X \leq \dim Y$ .*

**10. Example.** *Compactness is essential in 7–9, for there exist infinite-dimensional second countable spaces which can be injectively mapped in  $\mathbf{R}$  [6].*

**11. Proposition.** *If  $X$  is an infinite extremally disconnected compact space splittable over a space  $Y$  then  $\beta\omega \hookrightarrow Y$ .*

PROOF: It is true in ZFC (see [7]) that  $X = X_1 \cup X_2$ ,  $X_1 \cap X_2 = \emptyset$  and every compact  $K \subset X$  is finite ( $i = 1, 2$ ). Pick a continuous map  $f : X \rightarrow Y$  with  $f^{-1}f(X_i) = X_i$ . The space  $\beta\omega$  embeds in  $X$  and  $f \upharpoonright \beta\omega$  has finite point-inverses, so that  $\beta\omega \hookrightarrow f(\beta\omega)$  [8] and our proposition is proved.

**12. Corollary.** *If  $\beta\omega$  splits over a space  $Y$  then  $\beta\omega$  embeds in  $Y$ .*

□

## REFERENCES

- [1] Arhangel'skii A.V., *A general concept of splittability of a topological space* (in Russian), in: Proceedings of the Fifth Tyraspol Symposium on General Topology and its Applications, Kishinev, Shtiintsa, 1985, 8–10.
- [2] Arhangel'skii A.V., Shakhmatov D.B., *Splittable spaces and questions of functions approximations* (in Russian), in: Proceedings of the Fifth Tyraspol Symposium on General Topology and its Applications, Kishinev, Shtiintsa, 1985, 10–11.
- [3] ———, *On pointwise approximation of arbitrary functions by countable families of continuous functions* (in Russian), Trudy Seminara I.G. Petrovskogo **13** (1988), 206–227.
- [4] Tkachuk V.V., *Approximation of  $\mathbf{R}^X$  with countable subsets of  $C_p(X)$  and calibers of the space  $C_p(X)$* , Comment. Math. Univ. Carolinae **27** (1986), 267–276.
- [5] Bregman Yu.H., Šapirovskii B.E., Šostak A.P., *On partition of topological spaces*, Časopis pro pěstování mat. **109** (1984), 27–53.
- [6] Mazurkiewicz S., *Sur les problèmes  $\kappa$  et  $\lambda$  de Urysohn*, Fund. Math. **10** (1927), 311–319.
- [7] Tkachuk V., *Remainders over discrete spaces — some applications* (in Russian), Vestnik MGU, Mat., Mech., no. 4, 1990, 18–21.
- [8] Malyhin V.I.,  *$\beta N$  is prime*, Bull. Acad. Polon. Sci., Ser. Mat. **27** (1979), 295–297.
- [9] Engelking R., *General Topology*, PWN, Warszawa, 1977.
- [10] Skljarenko E.G., *A theorem on maps, lowering dimension (in Russian)*, Bull. Acad. Polon. Sci., Ser. Mat. **10** (1962), 429–432.

DEPARTMENT OF MECHANICS AND MATHEMATICS, SECTION OF GENERAL TOPOLOGY AND GEOMETRY, MOSCOW STATE UNIVERSITY, 119899 MOSCOW, RUSSIA

(Received May 6, 1991, revised April 21, 1992)