

On a problem of Gulevich on nonexpansive maps in uniformly convex Banach spaces

SEHIE PARK

Abstract. Let X be a uniformly convex Banach space, $D \subset X$, $f : D \rightarrow X$ a nonexpansive map, and K a closed bounded subset such that $\overline{\text{co}}K \subset D$. If (1) $f|_K$ is weakly inward and K is star-shaped or (2) $f|_K$ satisfies the Leray-Schauder boundary condition, then f has a fixed point in $\overline{\text{co}}K$. This is closely related to a problem of Gulevich [Gu]. Some of our main results are generalizations of theorems due to Kirk and Ray [KR] and others.

Keywords: uniformly convex, Banach space, Hilbert space, contraction, nonexpansive map, weakly inward map, demi-closed, Rothe condition, Leray-Schauder condition, (KR)-bounded, Opial's condition

Classification: Primary 47H10; Secondary 54H25

The well-known theorem of Browder-Göhde-Kirk assures existence of a fixed point for nonexpansive maps $f : K \rightarrow K$ where K is a *bounded* closed convex subset of a uniformly convex Banach space X . In [KR], Kirk and Ray showed that f can be replaced by a weakly inward nonexpansive map $f : K \rightarrow X$ while the boundedness of K can be replaced by that of the geometric estimator

$$G(x, fx) = \{z \in K : \|z - x\| \geq \|z - fx\|\}$$

for some $x \in K$ or more general sets. Note that any fixed point of f is contained in $G(x, fx)$.

On the other hand, Gulevich [Gu] considered the situation as follows: H is a Hilbert space, K is a nonempty bounded closed (not necessarily convex) subset of H , and $f : D \subset H \rightarrow H$ is a nonexpansive map, where $\overline{\text{co}}K \subset D$. Gulevich's basic theorem [Gu, Theorem 1] states that f has a fixed point in $\overline{\text{co}}K$ if f satisfies the Rothe condition $f(\text{Bd } K) \subset K$. He also raised as a problem whether H can be replaced by a uniformly convex Banach space.

In the present paper, we obtain some fixed point theorems on nonexpansive maps defined on closed (not necessarily bounded or convex) subsets of a Banach space. Our results are closely related to Gulevich's theorem and extend some known results of Kirk and Ray [KR], Goebel and Kuczumow [Go], and Browder [B1], [B2]. Moreover, we adopt more general boundary conditions on those nonexpansive maps. In fact, the weakly inwardness or the so-called Leray-Schauder condition is used in our results instead of the Rothe condition used in [Gu].

Recall that $f : K \rightarrow X$ is a *contraction* if there exists a $k \in [0, 1)$ such that

$$\|fx - fy\| \leq k\|x - y\| \quad \text{for all } x, y \in K;$$

and a *nonexpansive map* if

$$\|fx - fy\| \leq \|x - y\| \quad \text{for all } x, y \in K.$$

We say that f is *weakly inward* if $fx \in \overline{I_K}(x)$ for any $x \in \text{Bd } K$ (equivalently, for any $x \in K$), where $\overline{}$, Bd , and Int denote the closure, boundary, and interior, respectively, and

$$I_K(x) = \{x + c(y - x) : y \in K, c \geq 1\}.$$

Note that any map satisfying the Rothe condition is weakly inward.

We begin with the following:

Theorem 0. *Let K be a closed subset of a Banach space X and $f : K \rightarrow X$ a contraction satisfying one of the following:*

- (i) $f(\text{Bd } K) \subset K$.
- (ii) f is weakly inward.
- (iii) $0 \in \text{Int } K$ and $fx \neq mx$ for all $x \in \text{Bd } K$ and $m > 1$.

Then f has a unique fixed point.

Note that Theorem 0(i) is a particular case of Assad and Kirk [AK, Theorem 1], Theorem 0(ii) is due to Martinez-Yanez [M, Theorem] or, in a more general form, to Zhang [Z, Theorem 3.3], and Theorem 0(iii) to Gatica and Kirk [GK, Theorem 2.1]. There are more general results than Theorem 0. However, Theorem 0 is sufficient for our purpose. Note also that (iii) can be replaced by the following:

(iii)' there exists a $w \in \text{Int } K$ such that

$$fx - w \neq m(x - w) \quad \text{for all } x \in \text{Bd } K \quad \text{and } m > 1.$$

Moreover, (i) \Rightarrow (ii) and, whenever K is convex and $0 \in \text{Int } K$, we have (ii) \Rightarrow (iii).

A subset K of a vector space is said to be *star-shaped* if there exists a given point $x_0 \in K$ such that $tx_0 + (1 - t)x \in K$ for any $t \in (0, 1)$ and $x \in K$, where x_0 is called a *center* of K .

For the K and f in Theorem 0, we say that K is (KR)-*bounded* or bounded in the sense of Kirk-Ray [KR] if, for some bounded set $A \subset K$, the set

$$G(A) = \bigcap_{u \in A} G(u, fu)$$

is either empty or bounded.

The following is a generalization of the almost fixed point property of bounded closed subsets of a Banach space for nonexpansive maps.

Theorem 1. *Let X be a Banach space, K a closed subset of X , and $f : K \rightarrow X$ a nonexpansive map such that K is (KR)-bounded and one of the following holds:*

- (i) K is star-shaped and $f(\text{Bd } K) \subset K$.
- (ii) K is star-shaped and f is weakly inward.
- (iii) $0 \in \text{Int } K$ and $fx \neq mx$ for all $x \in \text{Bd } K$ and $m > 1$.

Then there exists a bounded sequence $\{x_n\}$ in K such that $\|x_n - fx_n\| \rightarrow 0$ as $n \rightarrow \infty$.

PROOF: For cases (i) and (ii) we may without loss of generality assume that 0 is the center. For $\alpha \in (0, 1)$, define $f_\alpha : K \rightarrow X$ by $f_\alpha x = \alpha fx$ for $x \in K$. Then clearly f_α is a contraction. We show that one of (i)–(iii) in Theorem 0 holds for f_α :

(i) Since K is star-shaped at center 0 , we have $\alpha K \subset K$. Since $f(\text{Bd } K) \subset K$, for $x \in \text{Bd } K$, we have $f_\alpha x = \alpha fx \in \alpha K \subset K$. Therefore, $f_\alpha(\text{Bd } K) \subset K$.

(ii) From $fx \in \bar{I}_K(x)$, we have $f_\alpha x = \alpha fx \in \alpha \bar{I}_K(x) \subset \bar{I}_K(x)$ since $I_K(x)$ is a star-shaped set with center 0 . See Zhang [Z, Theorem 1.2]. Note that (i) \Rightarrow (ii).

(iii) Suppose that $f_\alpha x = mx$ for some $x \in \text{Bd } K$ and $m > 1$, then $fx = \alpha^{-1} f_\alpha x = (\alpha^{-1} m)x$ and $\alpha^{-1} m > 1$, which contradicts our assumption.

Therefore, by Theorem 0, f_α has a fixed point $x_\alpha \in K$. Suppose that the set $\{x_\alpha : \alpha \in (0, 1)\}$ is not bounded. Then it is possible to choose $\alpha \in (0, 1)$ so that

$$\sup_{u \in A} \|fu\| \leq \inf_{u \in A} \|x_\alpha - u\|$$

and in addition, if $G(A) \neq \emptyset$, then α may also be chosen so that

$$\|x_\alpha\| > \sup\{\|x\| : x \in G(A)\}.$$

Therefore, for each $u \in A$,

$$\begin{aligned} \|x_\alpha - fu\| &= \|\alpha fx_\alpha - fu\| \leq \alpha \|fx_\alpha - fu\| + (1 - \alpha)\|fu\| \\ &\leq \alpha \|x_\alpha - u\| + (1 - \alpha)\|x_\alpha - u\| = \|x_\alpha - u\|. \end{aligned}$$

This implies $x_\alpha \in G(A)$, which is a contradiction. Thus $M = \sup\{\|x_\alpha\| : \alpha \in (0, 1)\} < \infty$ and we have

$$\|x_\alpha - fx_\alpha\| = (\alpha^{-1} - 1)\|x_\alpha\| \leq (\alpha^{-1} - 1)M,$$

yielding $\|x_\alpha - fx_\alpha\| \rightarrow 0$ as $\alpha \rightarrow 1$. This completes our proof. □

Note that Kirk and Ray [KR, Theorem 2.3] obtained Theorem 1(ii) for the case K is convex. In the second half of the proof of Theorem 1, we followed that of [KR, Theorem 2.3]. Note that Theorem 1(i) generalizes Dotson [D, Theorem 1].

A Banach space X is said to satisfy *Opial's condition* if, whenever a sequence $\{x_n\}$ converges weakly to $x_0 \in X$, then

$$\liminf_{n \rightarrow \infty} \|x_n - x\| > \liminf_{n \rightarrow \infty} \|x_n - x_0\|$$

for all $x \in X, x \neq x_0$. Opial [O] showed that if C is a weakly compact subset of a Banach space X satisfying this condition and $f : C \rightarrow X$ is nonexpansive, then $I - f$ is demi-closed ([B2], [Gö]); that is, if $\{x_n\} \subset C$ satisfies $x_n \rightarrow x$ weakly while $(I - f)x_n \rightarrow y$ strongly, then $(I - f)x = y$, where I is the identity map on C .

Examples of spaces satisfying Opial's condition are Hilbert spaces, l^p ($1 \leq p < \infty$), and uniformly convex Banach spaces with weakly continuous duality maps.

From Theorem 1, we have the following:

Theorem 2. *Let X be a Banach space, K a weakly compact subset of X , and $f : K \rightarrow X$ a nonexpansive map satisfying one of (i)–(iii) in Theorem 1.*

- (a) *If $I - f$ is demi-closed on K , then f has a fixed point.*
- (b) *If X satisfies Opial's condition, then f has a fixed point.*

PROOF: Since K is closed and bounded, f satisfies all the requirements of Theorem 1. Hence, there exists a sequence $\{x_n\}$ in K such that $\|x_n - fx_n\| \rightarrow 0$ as $n \rightarrow \infty$. Since K is weakly compact, we may assume that $x_n \rightarrow x$ weakly to some $x \in K$. Since $x_n - fx_n \rightarrow 0$ strongly, $x_n - fx_n = (I - f)x_n$, and $I - f$ is demi-closed, we conclude that $(I - f)x = 0$ and hence $x = fx$. This completes our proof. \square

Note that Zhang [Z, Theorem 3.8 and Corollaries 3.10, 3.11] obtained the multi-valued version of Theorem 2(ii), with different proof, and that Theorem 2(i) generalizes Dotson [D, Theorem 2]. Note also that if K is compact, then f has a fixed point in Theorem 2 without assuming the demi-closedness of $I - f$.

From Theorem 1, we also have the following:

Theorem 3. *Let X be a uniformly convex Banach space, D a subset of X , and $f : D \rightarrow X$ a nonexpansive map. Let K be a closed (KR) -bounded subset of X such that $\overline{\text{co}} K \subset D$ and one of (i)–(iii) of Theorem 1 holds for $f|_K$. Then f has a fixed point in $\overline{\text{co}} K$.*

PROOF: Since $f|_K$ satisfies all the requirements of Theorem 1, there exists a bounded sequence $\{x_n\}$ in K such that $\|x_n - fx_n\| \rightarrow 0$. Since $\{x_n\}$ is contained in a bounded closed convex subset $L \subset \overline{\text{co}} K$ and L is weakly compact, we may assume $x_n \rightarrow x_0$ weakly to some $x_0 \in L$. Since $I - f$ is demi-closed on L ([B2], [Gö]) and $(I - f)x_n \rightarrow 0$ strongly, we conclude that $(I - f)x = 0$, and hence $x = fx$. This completes our proof. \square

If we can eliminate the star-shapedness in (i), then Theorem 3(i) will be the required affirmative answer to Gulevich's problem. Moreover, Gulevich [Gu] noted that, for case (i) of Theorem 3 in a Hilbert space H , f has a fixed point in K .

In case $0 \in \text{Int } K$, Theorem 3(iii) generalizes [Gu, Theorem 1].

Note that the set $A \subset K$ for the (KR)-boundedness can be chosen so that $A \subset D$ and $f(A) \subset K$. See the proof of Theorem 1. Therefore, Theorem 3 generalizes Ray [R, Lemma 1].

For $D = K = \overline{\text{co}} K$, Theorem 3 reduces to the following:

Theorem 4. *Let X be a uniformly convex Banach space, K a closed convex subset, and $f : K \rightarrow X$ a nonexpansive map such that K is (KR)-bounded and one of (i)–(iii) in Theorem 0 holds. Then f has a fixed point.*

Note that Theorem 4(i) and (ii) are due to Kirk and Ray [KR, Theorem 2.3], which extends Goebel and Kuczumow [Go, Theorem 6]. Also note that Theorem 4(iii) extends Browder [B2, Theorem 1] for nonexpansive maps. For a Hilbert space X and a closed ball in X , Theorem 4(i) is due to Browder [B1, Theorem 2], which was used to show existence of periodic solutions for nonlinear equations of evolution.

Recently Canetti, Marino, and Pietramala [CMP] obtained multi-valued versions of Theorem 4(ii) and, under the stronger assumption of convexity, some other results similar to Theorems 1–3 for case (ii).

Finally, we note that the so-called Rothe condition (i) was first adopted by Knaster, Kuratowski, and Mazurkiewicz [KKM]. Also, the origin of the so-called Leray-Schauder condition (iii) seems to be Schaefer [S], and the following are well-known examples of that condition:

$$(A) \quad \|fx - x\|^2 \geq \|fx\|^2 - \|x\|^2 \text{ for } x \in \text{Bd } K.$$

$$(K) \quad \text{Re} \langle fx, x \rangle = \|x\|^2 \text{ for } x \in \text{Bd } K, x \neq 0, \text{ in a Hilbert space } H.$$

Or more generally,

$$(P) \quad (fx, Jx) \leq (x, Jx) \text{ for } x \in \text{Bd } K, 0 \in \text{Int } K, \text{ where } J \text{ is any duality map of } X \text{ into } 2^{X^*}.$$

Condition (A) is due to Altman [A], (K) to Krasnosel'skii [K] and Shinbrot [Sh], and (P) to Petryshyn [P].

Acknowledgement. This work was supported in part by KOSEF-951-0102-006-2.

REFERENCES

- [A] Altman M., *A fixed point theorem for completely continuous operators in Banach spaces*, Bull. Acad. Polon. Sci. **3** (1955), 409–413.
- [AK] Assad N.A., Kirk W.A., *Fixed point theorems for set-valued mappings of contractive type*, Pac. J. Math. **43** (1972), 553–562.
- [B1] Browder F.E., *Existence of periodic solutions for nonlinear equations of evolution*, Proc. Nat. Acad. Sci. USA **53** (1965), 1100–1103.
- [B2] Browder F.E., *Semicontractions and semiaccretive nonlinear mappings in Banach spaces*, Bull. Amer. Math. Soc. **74** (1968), 660–665.
- [CMP] Canetti A., Marino G., Pietramala P., *Fixed point theorems for multivalued mappings in Banach spaces*, Nonlinear Anal. TMA **17** (1991), 11–20.

- [D] Dotson W.G., *Fixed point theorems for non-expansive mappings on star-shaped subsets of Banach spaces*, J. London Math. Soc. (2) **4** (1972), 408–410.
- [GK] Gatica J.A., Kirk W.A., *Fixed point theorems for contraction mappings with applications to nonexpansive and pseudo-contractive mappings*, Rocky Mount. J. Math. **4** (1974), 69–79.
- [Go] Goebel K., Kuczumow T., *A contribution to the theory of nonexpansive mappings*, Bull. Calcutta Math. Soc. **70** (1978), 355–357.
- [Gö] Göhde D., *Zum Prinzip der kontraktiven Abbildung*, Math. Nachr. **30** (1965), 251–258.
- [Gu] Gulevich N.M., *Existence of fixed points of nonexpansive mappings satisfying the Rothe condition*, J. Soviet Math. **26** (1984), 1607–1611.
- [KR] Kirk W.A., Ray W.O., *Fixed-point theorems for mappings defined on unbounded sets in Banach spaces*, Studia Math. **64** (1979), 127–138.
- [KKM] Knaster B., Kuratowski C., Mazurkiewicz S., *Ein Beweis des Fixpunktsatzes für n -dimensionale Simplexe*, Fund. Math. **14** (1929), 132–137.
- [K] Krasnosel'skii M.A., *New existence theorems for solutions of nonlinear integral equations*, Dokl. Akad. Nauk SSSR **88** (1953), 949–952.
- [M] Martinez-Yanez C., *A remark on weakly inward contractions*, Nonlinear Anal. TMA **16** (1991), 847–848.
- [O] Opial Z., *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591–597.
- [P] Petryshyn W.V., *A new fixed point theorem and its application*, Bull. Amer. Math. Soc. **78** (1972), 225–229.
- [R] Ray W.O., *Zeros of accretive operators defined on unbounded sets*, Houston J. Math. **5** (1979), 133–139.
- [S] Schaefer H.H., *Neue Existenzsätze in der Theorie nichtlinearer Integralgleichungen*, Ber. Verh. Sächs. Akad. Wiss. Leipzig Math.-Natur. Kl. **101** (1955), no.7, 40pp.
- [Sh] Shinbrot M., *A fixed point theorem and some applications*, Arch. Rational Mech. Anal. **17** (1964), 255–271.
- [Z] Zhang S., *Star-shaped sets and fixed points of multivalued mappings*, Math. Japonica **36** (1991), 327–334.

DEPARTMENT OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 151–742, KOREA

E-mail: shpark@math.snu.ac.kr

(Received May 14, 1995)