

Relatively compact spaces and separation properties

A.V. ARHANGEL'SKII, I.V. YASCHENKO

Abstract. We consider the property of relative compactness of subspaces of Hausdorff spaces. Several examples of relatively compact spaces are given. We prove that the property of being a relatively compact subspace of a Hausdorff spaces is strictly stronger than being a regular space and strictly weaker than being a Tychonoff space.

Keywords: compact space, separation property, extension

Classification: 54A25

Notations and terminology follow [En] and [PW]. All spaces under consideration are assumed to be Hausdorff topological spaces. Condensation is a one-to-one continuous map onto. Cardinals are initial ordinals. Symbols ω , \mathbb{Z} , \mathbb{R} stand for the first infinite cardinal, the set of all integers and the real line, respectively. A space, closed in every regular (Hausdorff) space containing it is called *R-closed* (*H-closed*). A subspace Y of a space X is said to be *relatively compact* in X iff every open cover of X has a finite subcover of Y [Ra]. A subspace Y of a space X is said to be *relatively normal* in X iff whenever F_1 and F_2 are closed subsets of Y and $\text{Cl}_X F_1 \cap \text{Cl}_X F_2 = \emptyset$, then there are disjoint open subsets U_1 and U_2 of X , such that $F_1 \subset U_1$ and $F_2 \subset U_2$.

Every relatively compact subspace of a space X is relatively normal in X and every relatively normal subspace is a regular space [AH]. On the other hand, every subspace Y of a compact space K is relatively compact in K . Hence every Tychonoff space Y can be embedded into some space (e.g. $I^{\omega(Y)}$) as a relatively compact subspace. Therefore we could consider “being a relatively compact subspace” as a separation property, between regularity and complete regularity. Below we shall show that our property is *strictly* stronger than regularity and *strictly* weaker than complete regularity. We also use the following separation property.

1.1 Definition. A space X has the *countable separation property* iff whenever F is a closed subspace of X and $x \notin F$, there are open $W_i : i \in \omega$ such that for each $i \in \omega$ $x \notin W_i$, $F \subset W_i$ and $\text{Cl}_X W_{i+1} \subset W_i$.

Clearly, every Tychonoff space has the countable separation property and each space with countable separation property is regular.

1.2 Definition. A space Y will be said to be *potentially compact*, if there is a space X such that Y is a subspace of X and Y is relatively compact in X .

Thus we have

1.3 Proposition [AH]. *Every potentially compact space is regular.*

The following observation helps to identify several regular spaces which are not relatively compact in any Hausdorff space.

1.4 Proposition. *Let Y be an R -closed space which is relatively compact in a space X . Then Y is compact.*

PROOF: Choose arbitrary $x \in X \setminus Y$ and let $Y_1 = Y \cup \{x\}$. Clearly, Y_1 is relatively compact in X . Hence Y_1 is regular (Proposition 1.3). Then Y is closed in Y_1 . So, $x \notin \text{Cl}_X Y$. It follows that Y is closed in X . Thus Y is compact in itself, i.e. Y is compact. \square

So, any regular R -closed non-compact space is not relatively compact in any Hausdorff space containing it. One of the well-known examples with such properties is the Jones space over $Y = (\omega_1 + 1) \times (\omega_1 + 1) \setminus \{(\omega_1, \omega_1)\}$ [Jo], see also [PW, p.150–153]. Let $C = \omega_1 \times \{0\}$, $D = \{0\} \times \omega_1$, \bar{Y} be the quotient space obtained from $Y \times \omega$ by identifying C_{2n+1} with C_{2n+2} and D_{2n+2} with D_{2n+3} for each $n \in \mathbb{N}$ and $q : Y \times \omega \rightarrow \bar{Y}$ be the natural quotient map. Let $\tilde{Y} = \bar{Y} \cup \{z\}$ topologized as follows: \bar{Y} is an open subspace of \tilde{Y} , and $\{\{z\} \cup \cup_{n > k} Y_n : k \in \omega\}$ is a base in z . The resulting space is regular, not completely regular space [Jo], see also [PW, p.150–153].

1.5 Proposition. *Let \tilde{Y} be the Jones space over $(\omega_1 + 1) \times (\omega_1 + 1) \setminus \{(\omega_1, \omega_1)\}$. Then \tilde{Y} is not relatively compact in any Hausdorff space.*

PROOF: In view of Proposition 1.4 we need to prove only that \tilde{Y} is R -closed. Assume the contrary: X is a regular space, $\tilde{Y} \subset X$ and $x \in \text{Cl}_X \tilde{Y} \setminus \tilde{Y}$. Clearly $x \in \text{Cl}_X Y_n$ for some $n \in \omega$. Now we need the following fact:

1.6 Claim. *Let X be a regular space and $Y = (\omega_1 + 1) \times (\omega_1 + 1) \setminus \{(\omega_1, \omega_1)\} \subset X$. Then $|X \setminus Y| \leq 1$. If, moreover, $X \setminus Y \neq \emptyset$, then $\text{Cl}_X Y = (\omega_1 + 1) \times (\omega_1 + 1) = \beta Y$.*

It follows that $x \in \text{Cl}_X Y_{n+1}$, and by induction we have that $x \in \text{Cl}_X Y_k$ for each $k \geq n$. So, $x = z$ contradicting $x \notin \tilde{Y}$. \square

1.7 Remark. The above arguments also work to show that \tilde{Y} is not relatively normal in any regular space.

To construct a non-Tychonoff space Y which is relatively compact in some Hausdorff space X , we need the following lemma.

1.8 Lemma. *There are a Hausdorff space X and a Tychonoff zero-dimensional relatively compact subspace Y of X and two uncountable closed disjoint G_δ -subsets F_1 and F_2 of a space Y , such that $\text{Cl}_X F_1 \cap \text{Cl}_X F_2 = \emptyset$, F_1 and F_2 can be separated (in Y) by disjoint open sets, but whenever $f : Y \rightarrow \mathbb{R}$ is a continuous function then $|f^{-1}(0) \cap F_1| > \omega$ implies $|F_2 \setminus f^{-1}(0)| \leq \omega$. In particular, F_1 and F_2 cannot be separated (in Y) by a continuous real-valued function.*

PROOF: Let Y be the set

$$[-1, 1] \times [0, 1] \setminus \{(-1, 0), (1, 0)\}.$$

Basic elements for topology of Y are either:

1. $\{x\}$ for $x \in (-1, 1) \times (0, 1]$;

2.

$$\{-1, -1 + \varepsilon\} \times \{y\} : 0 < \varepsilon < 1\}$$

for $(x, y) \in \{-1\} \times (0, 1]$;

$$\{1 - \varepsilon, 1\} \times \{y\} : 0 < \varepsilon < 1\}$$

for $(x, y) \in \{1\} \times (0, 1]$;

3.

$$\{(x + e(1 - |x|)t, t) : t \in [0, 1] \setminus K, e \in \{-1/2, 0, 1/2\}\} K \in [(0, 1]]^{<\omega}$$

for $(x, y) \in (-1, 1) \times \{0\}$.

A typical neighborhood V_a of a point $(a, 0)$ can be described in the following way. Take the vertical line $l_0 : x = a$ through $(a, 0)$ and the two lines l_+ and l_- through $(a, 0)$ symmetrical with respect to l_0 having the slope $\pm 2/(1 - |a|)$. Then V_a is the intersection of the union $l_0 \cup l_+ \cup l_-$ with the rectangle $[-1, 1] \times [0, 1]$ from which any finite set of points different from $(a, 0)$ is removed.

Clearly Y is a Hausdorff zero-dimensional (hence Tychonoff) space. Let

$$F_1 = \{-1\} \times (0, 1]$$

$$F_2 = \{1\} \times (0, 1]$$

$$U_1 = [-1, -1 + 1/10] \times (0, 1]$$

$$U_2 = (1 - 1/10, 1] \times (0, 1].$$

Then F_1, F_2 are disjoint closed G_δ -subsets of Y , U_1, U_2 are disjoint open neighborhoods of F_1 and F_2 respectively. Moreover for

$$W_i = \left[-1, -1 + \frac{1}{2^{2i+1}}\right) \times (0, 1] : i \in \omega$$

we have: $\bigcap_{i \in \omega} W_i = F_1$ and $\text{Cl}_X W_{i+1} \subset W_i$.

First we prove

1.9 Claim. *Let $f : Y \rightarrow \mathbb{R}$ be a continuous function such that $|f^{-1}(0) \cap F_1| > \omega$. Then $|F_2 \setminus f^{-1}(0)| \leq \omega$.*

PROOF: Assume the contrary: $f : Y \rightarrow \mathbb{R}$ is a continuous function such that $|f^{-1}(0) \cap F_1| > \omega$ and $|F_2 \setminus f^{-1}(0)| > \omega$. Then there are $\varepsilon > 0$ and $P \in [F_2]^{>\omega}$ such that $\forall p \in P f(p) > 3\varepsilon$. Since f is continuous there are $\delta > 0$, $L \in [P]^{>\omega}$, $M \in [F_1]^{>\omega}$ such that $f(x, y) > 2\varepsilon$ whenever $1 - \delta < x < 1$, $y \in L$ and $f(x, y) < \varepsilon$ whenever $-1 < x < -1 + \delta$, $y \in M$. By the definition of the base of Y and continuity of f we have $f(x, 0) < \varepsilon$ for each $-1 < x < -1 + \delta$ and $f(x, 0) > 2\varepsilon$

for each $1 - \delta < x < 1$. Moreover, there is a family $\{K_x : x \in (-1, -1 + \delta)\} \subset [(0, 1]]^{<\omega}$ such that

$$f\left(\bigcup\left\{\{(x + e(1 - |x|)t, t) : t \in [0, 1] \setminus K_x, e \in \{\frac{-1}{2}, 0, \frac{1}{2}\}\}x \in (-1, -1 + \delta)\right\}\right) \subset [0, \varepsilon).$$

It follows that $f(x, 0) < \varepsilon$ for each $-1 < x < -1 + \frac{4}{5}\delta + \frac{1}{2}\frac{4}{5}\delta$ except for finitely many times. Therefore, $|\{x \in (-1, -1 + \frac{6}{5}\delta) : f(x, 0) > \varepsilon\}| < \omega$. Applying the argument above finitely many times we obtain that $|\{x \in (-1, \frac{1}{5}\delta) : f(x, 0) > \varepsilon\}| < \omega$. Similarly, starting from the right end of the segment $[-1, 1]$, we can prove that $|\{x \in (\frac{-1}{5}, 1) : f(x, 0) < 2\varepsilon\}| < \omega$. This contradiction completes the proof of the claim. \square

Now we shall construct a space X . Consider the Stone-Ćech extension βY of the space Y .

Let

$$\begin{aligned} \tilde{G}_1 &= \text{Cl}_{\beta Y}(F_1) \setminus Y \\ \tilde{G}_2 &= \text{Cl}_{\beta Y}(F_2) \setminus Y \\ \tilde{G}_3 &= \beta Y \setminus (\tilde{G}_1 \cup \tilde{G}_2 \cup Y). \end{aligned}$$

Let $X = G_1 \cup G_2 \cup G_3 \cup Y$ be the disjoint union of copies of sets $\tilde{G}_1, \tilde{G}_2, \tilde{G}_3, \tilde{Y}$. Basic elements for topology on X are either:

1. U for some open $U \subset Y$;
2. $\{g\} \cup (U \cap Y)$, for some $g \in G_3$, and some neighborhood U of g in βY ;
3. $\{g\} \cup (U \cap U_1)$, for some $g \in G_1$, and some neighborhood U of g in βY ;
4. $\{g\} \cup (U \cap U_2)$, for some $g \in G_2$, and some neighborhood U of g in βY .

Now $U_1 \cap U_2 = \emptyset$ implies that X is a Hausdorff space. Clearly, every open cover γ of X induces an open cover γ' of βY , members of which are unions of at most two elements of γ . It follows that Y is relatively compact in X . Finally, $\text{Cl}_X F_1 = G_1, \text{Cl}_X F_2 = G_2$ yields $\text{Cl}_X F_1 \cap \text{Cl}_X F_2 = \emptyset$. Thus Y and X satisfy all the required conditions.

We now turn to the second example.

1.10 Example. *There is a regular non-Tychonoff space \tilde{Y} with the countable separation property which is relatively compact in some Hausdorff space.*

PROOF: We use the notation of Lemma 1.8. Feed Y and X into the ‘‘Jones Machine’’ ([Jo], see also [PW]). Let $A = F_1, B = F_2, C = \text{Cl}_X F_1, D = \text{Cl}_X F_2$ and let \tilde{X} be the quotient space obtained from $X \times \omega$ by identifying C_{2n+1} with C_{2n+2} and D_{2n+2} with D_{2n+3} for each $n \in \mathbb{N}$ and $q : X \times \omega \rightarrow \tilde{X}$ be the natural quotient map. Let $\tilde{X} = \tilde{X} \cup \{z\}$ topologized as follows: \tilde{X} is an open subspace of \tilde{X} , and $\{\{z\} \cup \cup_{n>k} X_n : k \in \omega\}$ is a base in z . Let $\tilde{Y} = q(Y \times \omega) \cup \{z\}$. Clearly, \tilde{X} is Hausdorff and \tilde{Y} is a regular, non-Tychonoff subspace of \tilde{X} ([Jo], see also

[PW]). Since for each $n \in \omega$, $Y \times n$ is relatively compact in $X \times n$ and hence in \tilde{X} and every neighborhood of z contains all except at most finitely many $Y \times n$, \tilde{Y} is relatively compact in \tilde{X} . Finally, since $\tilde{Y} \setminus \{z\}$ is Tychonoff and $F_1 = \bigcap_{i \in \omega} W_i$ where $\text{Cl}_Y W_{i+1} \subset W_i$, it follows that Y has the countable separation property. \square

If we use the “Double Jones Machine” instead of the “Jones Machine” in Example 1.10 (i.e. consider the factor space of the product $X \times \mathbb{Z}$ and add two points $-\infty$ and ∞) we obtain a bit stronger

1.11 Example. *There is a regular space Z which is relatively compact in some Hausdorff space and has the countable separation property, but which is not functionally Hausdorff.*

Now using Herrlich technique [He] one can obtain

1.12 Example. *There is a regular space Z which is relatively compact in some Hausdorff space, such that all real-valued continuous functions on Z are constants.*

What if X has some separation property stronger than Hausdorff? First, since every Hausdorff space can be embedded as a closed subspace into some semiregular space the following assertion holds.

2.1 Proposition. *A space Y can be embedded as a relatively compact subspace into a Hausdorff space if and only if Y can be embedded as a relatively compact subspace into a semiregular space.*

On the other hand, if Y is relatively compact in some Urysohn space, then Y must be Tychonoff. Indeed, we have

2.2 Theorem. *Let Y be a dense relatively compact subspace of an Urysohn space X . Then there is a compact space Z , and condensation $f : X \rightarrow Z$, such that for each $y \in X$ the restriction $f|_{Y \cup \{y\}}$ of f to $Y \cup \{y\}$ is a homeomorphism of $Y \cup \{y\}$ onto the image.*

To prove this we need the following

2.3 Proposition. *Let Y be a dense relatively compact subspace of a space X . Then X is H-closed.*

PROOF: Direct check. \square

PROOF OF THE THEOREM: Let Z be the semiregularization of a space Z , and let $f : X \rightarrow Z$ be the natural condensation. Then Z is a semiregular Urysohn space, and $f(Y)$ is relatively compact in Z . Proposition 2.3 yields that Z is H-closed. So Z is a semiregular Urysohn H-closed space. Hence Z is compact. Now take arbitrary $y \in X$. Then $Y \cup \{y\}$ is relatively compact in X . Therefore, the semiregularization of $Y \cup \{y\}$ is again $Y \cup \{y\}$. It follows that $f|_{Y \cup \{y\}}$ is a homeomorphism. \square

2.4 Definition. A subspace Y of a space X is said to be *real-normal* in X iff every two subspaces of Y having disjoint closures in X can be separated in X by a continuous real-valued function.

2.5 Corollary. *Let Y be a dense relatively compact subspace of an Urysohn space X . Then Y is Tychonoff, X is functionally Hausdorff and Y is real-normal in X .*

PROOF: We shall prove that Y is real-normal in X , other properties are obvious. Let $F_1, F_2 \subset Y$, $\text{Cl}_X F_1 \cap \text{Cl}_X F_2 = \emptyset$. Use the notations of 2.2. Since for each $y \in X$ $f|_{Y \cup \{y\}}$ is a homeomorphism $\text{Cl}_Z f(F_1) \cap \text{Cl}_Z f(F_2) = \emptyset$. Hence $\text{Cl}_Z f(F_1)$ and $\text{Cl}_Z f(F_2)$ can be separated in compact space Z by a continuous real-valued function. Therefore, the same is true for F_1 and F_2 in X . \square

2.6 Problem. *Find an “inner” characterization of the potential compactness.*

2.7 Problem. *Is there a regular space without a dense Tychonoff subspace?*

In the opinion of the authors, the last question should have the negative answer, but to construct the corresponding example entirely new ideas will be needed. Indeed, regular non-Tychonoff spaces are always constructed by adding new “bad” points to a “not-so-nice” Tychonoff space.

REFERENCES

- [AH] Arhangel'skii A.V., Hamdi M.M. Genedi, *Foundations of the theory of relative topological properties*, General Topology. Spaces and mappings., MGU Moscow, 1989, pp. 3–48.
- [En] Engelking R., *General Topology*, PWN Warszawa, 1987.
- [He] Herrlich H., *Ordnungsfähigkeit total-diskontinuierlicher Räume*, Math. Ann. **159** (1965), 77–80.
- [Jo] Jones F.B., *Hereditary separable, non-completely regular spaces*, Topology Conf., Virginia Polytechnic Inst. and State U. 1973, Springer Lecture Notes in Mathematics **375** (1974), 149–152.
- [PW] Porter J.R., Woods R.G., *Extensions and Absolutes of Hausdorff Spaces*, Springer-Verlag, 1988.
- [Ra] Ranchin D.B., *On compactness modulo ideal*, DAN SSSR **202** (1972), 761–764.

DEPARTMENT OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY MOSCOW, RUSSIA

and

OHIO UNIVERSITY, ATHENS, OHIO, USA

E-mail: aarhange@oucsace.cs.ohiou.edu

MOSCOW STATE UNIVERSITY, MOSCOW, RUSSIA

E-mail: ivan@sch57.mcn.msk.su

(Received March 14, 1995)