

## A note on Banach spaces with $\ell^1$ -saturated duals

DENNY H. LEUNG

*Abstract.* It is shown that there exists a Banach space with an unconditional basis which is not  $c_0$ -saturated, but whose dual is  $\ell^1$ -saturated.

*Keywords:* dual space,  $\ell^1$ -saturated spaces

*Classification:* 46B10

Let  $E$  and  $F$  be Banach spaces. We say that  $E$  is  $F$ -saturated if every infinite dimensional closed subspace of  $E$  contains an isomorphic copy of  $F$ . In [2], it is shown that there exists a  $c_0$ -saturated Banach space with an unconditional basis whose dual contains an isomorphic copy of  $\ell^2$ . In this note, we give an example where the dual situation occurs. It is shown that there is a Banach space with an unconditional basis which contains an isomorphic copy of  $\ell^2$ , and whose dual is  $\ell^1$ -saturated.

We follow standard Banach space terminology as used in [3]. Our example is a certain subspace of the weak  $L^2$  space  $L^{2,\infty}[0, \infty)$ . Recall that this is the space of all measurable functions  $f$  on  $[0, \infty)$  such that

$$(1) \quad \|f\| = \sup_{c>0} c(\lambda\{|f| > c\})^{1/2} < \infty,$$

where  $\lambda$  is the Lebesgue measure on  $[0, \infty)$ . Although equation (1) only defines a quasi-norm on  $L^{2,\infty}[0, \infty)$ , it is well known that it is equivalent to a norm on  $L^{2,\infty}[0, \infty)$ , and that  $L^{2,\infty}[0, \infty)$  is norm complete. The reader may consult [4] for further information concerning the family of Lorentz spaces, of which  $L^{2,\infty}[0, \infty)$  is a member. Finally, for a measurable function  $f$ , we let  $f^*$  be the decreasing rearrangement of  $|f|$ , as defined in §2a of [4].

**Proposition 1.** *For each  $n \in \mathbb{N}$ , define*

$$f_n(t) = \begin{cases} \min(2^{n/2}, (t - n + 1)^{-1/2}) & \text{if } n - 1 < t \leq n, \\ 0 & \text{otherwise.} \end{cases}$$

*Then for any  $m \in \mathbb{N}$ , and any sequence of scalars  $(a_n)$ ,*

$$\left(\frac{1}{2} \sum_{n=1}^m a_n^2\right)^{1/2} \leq \left\| \sum_{n=1}^m a_n f_n \right\| \leq \left(\sum_{n=1}^m a_n^2\right)^{1/2},$$

where  $\|\cdot\|$  refers to the quasi-norm defined by (1).

PROOF: Given a scalar sequence  $(a_n)_{n=1}^m$ , let  $a = \max\{|a_n| : 1 \leq n \leq m\}$ , and suppose this maximum is attained at  $n_0$ . Let  $N$  be the set of all natural numbers  $\leq m$  such that  $a_n > 2^{-n/2}a$ . If  $0 < c < a$ , and  $n \in N$ , then  $c < 2^{n/2}a_n$ . Hence  $\lambda\{|a_n f_n| > c\} = \min(a_n^2/c^2, 1)$ . Therefore,

$$\begin{aligned} \left\| \sum_{n=1}^m a_n f_n \right\| &\geq \sup_{0 < c < a} c \left( \lambda \left\{ \sum_{n \in N} |a_n f_n| > c \right\} \right)^{1/2} \\ &= \sup_{0 < c < a} \left( \sum_{n \in N} \min(a_n^2, c^2) \right)^{1/2} \\ &= \left( \sum_{n \in N} a_n^2 \right)^{1/2}. \end{aligned}$$

If  $1 \leq n \leq m$ , and  $n \notin N$ , then  $a_n \leq 2^{-n/2}a$ . Thus

$$\sum_{\substack{1 \leq n \leq m \\ n \notin N}} a_n^2 \leq a^2 \sum_{n=1}^m 2^{-n} < a^2.$$

Hence

$$\begin{aligned} \left\| \sum_{n=1}^m a_n f_n \right\| &\geq \left( \sum_{n=1}^m a_n^2 - \sum_{\substack{1 \leq n \leq m \\ n \notin N}} a_n^2 \right)^{1/2} \\ &\geq \left( \sum_{n=1}^m a_n^2 - a^2 \right)^{1/2} \\ &= \left( \sum_{\substack{1 \leq n \leq m \\ n \neq n_0}} a_n^2 \right)^{1/2}. \end{aligned}$$

Clearly,  $\|\sum_{n=1}^m a_n f_n\| \geq \|a_{n_0} f_{n_0}\| = |a_{n_0}|$  as well. This proves the first half of the inequality. Observe that for any  $c > 0$ ,

$$\lambda \left\{ \left| \sum_{n=1}^m a_n f_n \right| > c \right\} = \sum_{n=1}^m \lambda \left\{ |f_n| > \frac{c}{|a_n|} \right\} \leq \frac{1}{c^2} \sum_{n=1}^m a_n^2.$$

The second half of the inequality follows. □

For  $n \in \mathbb{N}$ , and  $1 \leq j \leq 2^n$ , let  $g_{n,j}$  be the characteristic function of the interval  $[n-1 + (j-1)2^{-n}, n-1 + j2^{-n})$ . Let  $E$  be the closed linear span of the sequence  $(g_{n,j})_{j=1}^{2^n} \prod_{n=1}^{\infty}$ . Clearly  $(g_{n,j})_{j=1}^{2^n} \prod_{n=1}^{\infty}$  is an unconditional basis of  $E$ .

**Proposition 2.** *The space  $E$  contains an isomorphic copy of  $\ell^2$ .*

PROOF: Recall the sequence  $(f_n)$  defined in Proposition 1. For each  $n$ , let  $h_n = \sum_{j=1}^{2^n} \sqrt{2^n/j} g_{n,j}$ . Then  $0 \leq h_n \leq f_n \leq \sqrt{2} h_n$  for all  $n$ . It follows from Proposition 1 that the subspace  $\{h_n\}$  of  $E$  is isomorphic to  $\ell^2$ .  $\square$

It remains to show that  $E'$  is  $\ell^1$ -saturated. Let  $F$  be the closed subspace of  $L^{2,\infty}[0, \infty)$  generated by  $L^1 \cap L^\infty$ . It is well known that  $F'$  is canonically isomorphic to  $L^{2,1}[0, \infty)$ , where the latter is the space of all measurable functions  $f$  on  $[0, \infty)$  such that

$$\|f\|_{2,1} = \int_0^\infty \frac{f^*(t)}{\sqrt{t}} dt < \infty.$$

Let  $\Sigma$  be the  $\sigma$ -algebra generated by the sets  $[n-1 + (j-1)2^{-n}, n-1 + j2^{-n})$ ,  $1 \leq j \leq 2^n$ ,  $n \in \mathbb{N}$ . Then clearly  $E$  is the subspace of  $F$  consisting of all  $\Sigma$ -measurable functions. It follows easily that  $E'$  can be identified canonically with the subspace of  $L^{2,1}[0, \infty)$  consisting of all  $\Sigma$ -measurable functions. Now if  $G$  is a subspace of  $E'$ , then it contains a basic sequence equivalent to a normalized disjointly supported sequence  $(u_n)$  in  $L^{2,1}[0, \infty)$ . By [1, Corollary 2.4],  $\{u_n\}$ , and hence  $G$ , contains a copy of  $\ell^1$ .

We end this note with the following problem.

**Problem.** Suppose  $E$  is a Banach space (with or without an unconditional basis) such that  $E'$  has the Schur property. Is  $E$  necessarily  $c_0$ -saturated?

#### REFERENCES

- [1] Carothers N.L., Dilworth S.J., *Subspaces of  $L^{p,q}$* , Proc. Amer. Math. Soc. **104** (1988), 537–545.
- [2] Leung D.H., *On  $c_0$ -saturated Banach spaces*, Illinois J. Math. **39** (1995), 15–29.
- [3] Lindenstrauss J., Tzafriri L., *Classical Banach Spaces I*, Springer-Verlag, 1977.
- [4] Lindenstrauss J., Tzafriri L., *Classical Banach Spaces II*, Springer-Verlag, 1979.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119260

*E-mail:* matlhh@leonis.nus.sg

(Received November 27, 1995)