

The a priori estimate of the maximum modulus to solutions of doubly nonlinear parabolic equations*

LIANG XITING, WU ZAIDE

Abstract. The a priori estimate of the maximum modulus of the generalized solution is established for a doubly nonlinear parabolic equation with special structural conditions.

Keywords: doubly nonlinear parabolic equation, generalized solution, bounded solution, maximum modulus, a priori estimate

Classification: 35K65

Introduction

Since Lions [1] proposed in 1969 to consider the doubly nonlinear parabolic equation,

$$\frac{\partial}{\partial t}(|u|^{\lambda-2}u) - \frac{\partial}{\partial t}(|u_{x_i}|^{p-2}u_{x_i}) = f,$$

there are papers devoted to the existence of generalized solutions of more general doubly nonlinear parabolic equations (see, for example, [2]–[3]). But the investigations concerning the properties of generalized solutions still seldom appear. In [4], Liang Xiting & Liang Xuexin proved the local and global boundedness to generalized solutions of doubly nonlinear parabolic equations with a more general structural condition than (1), (2) below. However, in that paper, it does not supply any a priori estimate for the maximum modulus of solutions. In this present paper, we give a certain a priori estimate of the maximum modulus for the generalized solutions of the doubly nonlinear parabolic equation (1) with structure conditions (2). These are the extensions of the corresponding results for elliptic equations.

Let G be a bounded domain in the n -dimensional Euclidean space E^n and $T > 0$ a finite real number. Consider on $Q = G \times (0, T)$ the following doubly nonlinear parabolic equation

$$(1) \quad \frac{\partial}{\partial t}(|u|^{\lambda-2}u) - \operatorname{div} \underline{A}(x, t, u, \nabla u) + B(x, t, u, \nabla u) = 0,$$

where $2 \leq \lambda < np/(n-p)$ as $1 < p < n$ and $2 \leq \lambda < \infty$ as $p \geq n$, $\underline{A}(x, t, u, \xi)$ and $B(x, t, u, \xi)$ are defined on $Q \times E^1 \times E^n$, continuous with respect to u and ξ for

* This work was in part supported by The Natural Science Foundation of China and by The Foundation of Zhongshan University Advanced Research Center

fixed x and t , measurable with respect to x and t for fixed u and ξ and satisfying the following structural conditions, respectively:

$$(2) \quad \begin{aligned} \xi \cdot \underline{A}(x, t, u, \xi) &\geq |\xi|^p, \quad p > 1 \\ |\underline{A}(x, t, u, \xi)| &\leq \kappa |\xi|^{p-1}, \quad \kappa \geq 1 \\ B(x, t, u, \xi) &= B_1(x, t, u, \xi) + \mu |u|^{\alpha-2} u \\ |B_1(x, t, u, \xi)| &\leq b(x, t) |\xi|^\beta + f(x, t) \end{aligned}$$

where $\mu \geq 0$;

$$(3) \quad \begin{aligned} q = p(n + \lambda)/(n + p) \leq \alpha &\leq p(1 + \lambda/n) = l \quad \text{and} \\ \beta_0 = p - (n + p)/(n + \lambda) &\leq \beta < p \\ b(x, t) &\in L_r(Q) \end{aligned}$$

$$(4) \quad \begin{aligned} 1/r = 1 - \beta/p - 1/l \quad \text{as} \quad \beta_0 &\leq \beta < \beta_1 = p - n/(n + \lambda) \\ r = \infty \quad \text{as} \quad \beta &= \beta_1 \\ r > (n + p)/(p - \beta) \quad \text{as} \quad \beta_1 &< \beta < p \end{aligned}$$

$$(5) \quad f(x, t) \in L_s(Q), \quad s > (n + p)/p.$$

We call u a generalized solution of (1), if

$$\begin{aligned} u &\in C(0, T; L_\lambda(G)) \cap L_p(0, T; W_p^1(G)) \quad \text{as} \quad \beta_0 \leq \beta \leq \beta_1 \\ u &\in C(0, T; L_\lambda(G)) \cap L_p(0, T; W_p^1(G)) \cap L_{t^*}(Q) \\ (1 - (p - \beta)(n + \lambda)/(n + p))/t^* &+ (p - \beta)(n + \lambda)/((n + p)l) + \beta/p + 1/r = 1, \\ &\text{as} \quad \beta_1 < \beta < p \end{aligned}$$

and the following holds

$$(1)' \quad \begin{aligned} &\int_0^t \int_G \{-\nu_t |u|^{\lambda-2} \mu + \nabla \nu \cdot \underline{A}(x, t, u, \nabla u) + \nu B(x, t, u, \nabla u)\} dx dt \\ &+ \int_G \nu(x, t) |u(x, t)|^{\lambda-2} u(x, t) \Big|_{t=0}^{t=t} dx = 0 \\ &\forall t \in (0, T), \quad \phi \in W_\lambda^1(0, T; L_\lambda(G)) \cap L_p(0, T; \dot{W}_p^1(G)); \end{aligned}$$

our results are the following:

Theorem 1. *Suppose (2)–(5) are fulfilled in which $\mu > 0$ and $s = \infty$. Let u be a generalized solution of (1). In addition, there is a constant $M > 0$ such that*

$$(6) \quad (u - M)^+ = \max(u - M, 0) \in L_p(0, T; \dot{W}_p^1(G)) \quad \text{and} \quad (u - M)^+ \Big|_{t=0} = 0.$$

Then holds:

$$(7) \quad \operatorname{ess\,sup}_Q u^+ \leq \max\{M, (\|f\|_{L_\infty(Q)}/\mu)^{1/(\alpha-1)}\}.$$

In particular, if $1 < p < 2n/(n + \lambda - 2)$ holds,

$$(8) \quad \operatorname{vrai\,max}_Q u^+ \leq M \quad \text{or} \quad \operatorname{vrai\,max}_Q u^+ < (\|f\|_{L_\infty(Q)}/\mu)^{1/(\alpha-1)}$$

and there exists a constant $\theta > 0$ depending only on $n, p, \mu, \lambda, \kappa, b(x, t)$ and $|Q|$, the $n + 1$ dimensional Lebesgue measure of Q , such that

$$(9) \quad \operatorname{ess\,sup}_Q u^+ \leq \max\{M, (\|f\|_{L_\infty(Q)}/\mu(1 + \theta))^{1/(q-1)}\},$$

provided $\alpha = q$ and $\beta = \beta_0$.

Theorem 2. Suppose (2)–(5) are fulfilled in which $\mu \geq 0$ and $\beta = \beta_0$. Let u be a generalized solution of (1) satisfying (6). Then there exists a constant C depending only on $n, p, \mu, \lambda, \kappa, b(x, t)$ and $|Q|$ such that

$$(10) \quad \operatorname{ess\,sup}_Q u^+ \leq M + C\|f\|_{L_s(Q)}^{1/(q-1)}.$$

Proof of Theorems

In order to prove the theorems we need the following lemmata. The proof of Lemma 1 is essentially the same as in [5, Chapter II, § 3] and Lemma 2 is a special case of [6, Chapter II, Lemma 5.1].

Lemma 1. Let $u \in C(0, T; L_\lambda(G)) \cap L_p(0, T; \dot{W}_p^1(G))$. Then

$$\begin{aligned} \|u\|_{L_l(Q)} &\leq C(n, p, \lambda)\|u\|_Q^{1/q}, \quad l = p(1 + \lambda/n), \quad q = p(n + \lambda)/(n + p), \\ \|u\|_Q &= \operatorname{ess\,sup}_{t \in (0, T)} \int_G |u|^\lambda dx + \int \int_Q |\nabla u|^p dx dt. \end{aligned}$$

Lemma 2. Let $u \in L_1(Q)$ satisfying

$$\int \int_Q (u - k)^+ dx dt \leq F|Q \cap \{u > k\}|^{1+\tau} \quad \forall k \geq k_0 \geq 0,$$

where F, τ are positive constants. Then

$$\operatorname{ess\,sup}_Q u^+ \leq k_0 + (1 + 1/\tau)F|Q|^\tau.$$

PROOF OF THEOREM 1: We take for $\theta \geq 0$

$$k(\theta) = (\|f\|_{L_\infty(Q)} / (\mu(1 + 2\theta)))^{1/(\alpha-1)}.$$

If $M < k(\theta)$, then there is a $\theta' > 0$ such that $M < k(\theta)$ for $\theta \in (0, \theta')$. Let $k_0 = \max\{M, k(\theta)\}$. For any $k \geq k_0$, we have on the set $Q \cap \{u > k\}$ that

$$(11) \quad f(x, t) - \mu|u|^{\alpha-2} \leq F = \begin{cases} 0 & \text{as } M \geq k(\theta), \\ (2\theta/(1 + 2\theta))\|f\|_{L_\infty(Q)} & \text{as } M < k(\theta). \end{cases}$$

For the sake of simplicity we assume $u \in W_\lambda^1(0, T; L_\lambda(G))$. Then for $k \geq k_0$, we may take $\nu = (u - k)^+$ as a test function (in general we should take the time average of ν as a test function and invoke a limit process). Inserting such a ν into (1)' we obtain by integrating by part with respect to t that

$$(12) \quad \int_0^t \int_G (u - k)^+ (|u|^{\lambda-2}u)_t dx dt + \int_0^t \int_{G \cap \{u > k\}} |\nabla u|^p dx dt \\ \leq \int_0^t \int_G (u - k)^+ (b(x, t)|\nabla u|^\beta + F) dx dt, \quad t \in (0, T).$$

Represent

$$(13) \quad I = \int_0^t \int_G (u - k)^+ (|u|^{\lambda-2}u)_t dx dt.$$

In order to estimate I we put

$$\tilde{u} = \begin{cases} u & \text{as } u > k, \\ k & \text{as } u \leq k; \end{cases}$$

then

$$I = \int_0^t \int_G (\tilde{u} - k)(|\tilde{u}|^{\lambda-2}\tilde{u})_t dx dt \\ = (\lambda - 1) \int_0^t \int_G \left(\frac{\bar{u}^\lambda}{\lambda} - \frac{k\bar{u}^{\lambda-1}}{\lambda-1} + \frac{k^\lambda}{\lambda(\lambda-1)} \right)_t dx dt \\ = (\lambda - 1) \int_G \left(\frac{\bar{u}^\lambda}{\lambda} - \frac{k\bar{u}^{\lambda-1}}{\lambda-1} + \frac{k^\lambda}{\lambda(\lambda-1)} \right) dx \\ \geq (\lambda - 1) \int_{G \cap \{k < u < h\}} \left(\frac{u^\lambda}{\lambda} - \frac{ku^{\lambda-1}}{\lambda-1} + \frac{k^\lambda}{\lambda(\lambda-1)} \right) dx.$$

Let

$$g(\eta) = \frac{1}{\lambda} - \frac{\eta}{\lambda-1} + \frac{\eta^\lambda}{\lambda(\lambda-1)} - \frac{(1-\eta)^\lambda}{\lambda^2}, \quad \eta \in (0, 1).$$

On account of $g(0) = 1/\lambda - 1/\lambda^2 > 0$ and $g(1) = 0$ and

$$\eta^{\lambda-1} + \left(\frac{\lambda-1}{\lambda}\right)(1-\eta)^{\lambda-1} < 1 \quad \text{for } \lambda \geq 2 \quad \text{and } \eta \in (0, 1)$$

we have

$$g'(\eta) = \frac{1}{\lambda-1} + \frac{\eta^{\lambda-1}}{\lambda-1} + \frac{(1-\eta)^{\lambda-1}}{\lambda} < 0 \quad \text{for } \eta \in (0, 1).$$

This implies

$$(14) \quad \frac{u^\lambda}{\lambda} - \frac{ku^{\lambda-1}}{\lambda-1} + \frac{k^\lambda}{\lambda(\lambda-1)} \geq \frac{(u-k)^\lambda}{\lambda^2} \quad \text{on } Q \cap \{u > k\},$$

$$I \geq C(\lambda) \int_{G \cap \{u > k\}} (u-k)^\lambda dx.$$

Combining (13), (14) with (12) we get that

$$(15) \quad \int_{G \cap \{u > k\}} (u-k)^\lambda dx + \int_0^t \int_{G \cap \{u > k\}} |\nabla u|^p dx dt$$

$$\leq C \int_0^t \int_G (u-k)^+ (b(x,t)|\nabla u|^\beta + F) dx dt.$$

Take the supremum for $t \in (0, T)$ yields

$$(16) \quad \text{ess sup}_{t \in (0, T)} \int_G |(u-k)^+|^\lambda dx + \int \int_{Q \cap \{u > k\}} |\nabla u|^p dx dt$$

$$\leq C \int \int_Q (u-k)^+ (b(x,t)|\nabla u|^\beta + F) dx dt.$$

In what follows we write

$$(17) \quad F(k) = \int \int_Q (u-k)^+ F dx dt$$

which is a nondecreasing function of k . In virtue of $(u-k)^+ \in L_p(0, T; \dot{W}_p^1(G))$ for $k \geq M$ we have by the use of Lemma 1 that

$$(18) \quad \int \int_Q (u-k)^+ b(x,t)|\nabla u|^\beta dx dt$$

$$= \int \int_Q |(u-k)^+|^{1-(p-\beta)(n+p)/(n+\lambda)+(p-\beta)(n+p)/(n+\lambda)} b(x,t)|\nabla u|^\beta dx dt$$

$$\leq \varepsilon(k) \left(\int \int_Q |(u-k)^+|^l dx dt \right)^{(p-\beta)(n+p)/(n+\lambda)l} \times$$

$$\times \left(\int \int_Q |\nabla(u-k)^+|^p dx dt \right)^{\beta/p}$$

$$\leq C\varepsilon(k) \| |(u-k)^+ \|_{lQ},$$

where $\varepsilon(k) = \|b(x, t)\|_{L_r(Q \cap \{u > k\})} U$,

$$(19) \quad \begin{aligned} U &= \|u\|_{L_i(Q)}^{1-(p-\beta)(n+p)/(n+\lambda)} \quad \text{as } \beta_0 \leq \beta \leq \beta_1 \quad \text{and} \\ U &= \|u\|_{L_{i^*}(Q)}^{1-(p-\beta)(n+p)/(n+\lambda)} \quad \text{as } \beta_1 < \beta < p. \end{aligned}$$

In virtue of

$$|Q \cap \|u > k\|| \leq k^{-l} \iint_Q |u|^l dx dt \rightarrow 0 \quad \text{as } k \rightarrow \infty$$

and the absolute continuity of a Lebesgue integral we have $\varepsilon(k) \rightarrow 0$ as $k \rightarrow \infty$. So, we can take an $h_0 > k_0$ such that

$$(20) \quad C\varepsilon(k) \leq 1/2 \quad \text{as } k \geq h_0.$$

From (16)–(19) it follows

$$(21) \quad \| (u - h_0)^+ \| \| \|_Q \leq CF(h_0).$$

Let $h > k \geq k_0$ be arbitrary; it follows from (16) that

$$(22) \quad \begin{aligned} \text{ess sup}_{t \in (0, T)} \int_G ((u - k)^+ - (u - h)^+) dx + \int \int_{Q \cap \{k < u < h\}} |\nabla u|^p dx dt \\ \leq C \left\{ \int \int_Q (u - k)^+ b(x, t) |\nabla u|^\beta dx dt + F(k) \right\}. \end{aligned}$$

The effective domain of the integral appearing on the right hand side of (22) excludes any set of the form $Q \cap \{u = \text{const}\}$ having a positive measure. For simplicity we assume $|Q \cap \{u = \text{const}\}| = 0$. For any $k > k_0$, we take

$$h_{-1} = \infty > h_0 > h_1 > \dots > h_m > h_{m+1} = k$$

such that $Q_i = Q \cap \{h_{i-1} > u > h_i\}$ satisfies

$$|Q_i|^{1/r} = \varepsilon/N \quad (i = 1, 2, \dots, m) \quad \text{and} \quad |Q_{m+1}|^{1/r} \leq \varepsilon/N,$$

where ε will be specified and N is defined by

$$(23) \quad \varepsilon(N) = \left(\int \int_{Q \cap \{b(x, t) > N\}} b(x, t)^r dx dt \right)^{1/r} \leq \varepsilon.$$

It follows from (23) that

$$(24) \quad \begin{aligned} \|b(x, t)\|_{L_r(Q_i)} &\leq \left(\int \int_{Q \cap \{b(x, t) > N\}} b(x, t)^r dx dt \right)^{1/r} + N|Q_i|^{1/r} \\ &\leq \varepsilon(N) + N|Q_i|^{1/r} \leq 2\varepsilon \quad i = 1, 2, \dots, m + 1. \end{aligned}$$

Denote for $i = 1, 2, \dots, m + 1$

$$(25) \quad u_i = (u - h_i)^+ - (u - h_{i-1})^+.$$

Then for almost everywhere $(x, t) \in Q$ we have

$$(u - h_i)^+ = \sum_{j=0}^i u_j \quad \text{on } Q \cap \{u > h_i\},$$

$\nabla u_i = \nabla u$ as $(x, t) \in Q_i$ and $\nabla u_i = 0$ as (x, t) otherwise. It follows from (22) by taking $k = h_{i+1}$ and $h = h_i$ that

$$(26) \quad \|u_{i+1}\|_Q \leq C(II + F(k)),$$

where

$$(27) \quad \begin{aligned} II &= \iint_{Q \cap \{u > h_{i+1}\}} (u - h_{i+1}) b(x, t) |\nabla u|^\beta dx dt \\ &= \iint_{Q \cap \{u > h_{i+1}\}} \left(u_{i+1} + \sum_{j=0}^i u_j \right) b(x, t) |\nabla u|^\beta dx dt = III + IV \end{aligned}$$

$$(28) \quad \begin{aligned} III &= \iint_{Q \cap \{u > h_{i+1}\}} u_{i+1} b(x, t) |\nabla u|^\beta dx dt \\ &\leq \iint_{Q \cap \{u > h_{i+1}\}} u_{i+1} b(x, t) \left(\sum_{j=0}^{i+1} |\nabla u_j| \right)^\beta dx dt \\ &\leq 2^\beta \iint_{Q \cap \{u > h_{i+1}\}} u_{i+1} b(x, t) \left[|\nabla u_{i+1}|^\beta + \left(\sum_{j=0}^i |\nabla u_j| \right)^\beta \right] dx dt \\ &\leq 2^\beta \iint_{Q_{i+1}} u_{i+1} b(x, t) |\nabla u_{i+1}|^\beta dx dt \\ &\quad + (2m)^\beta \sum_{j=0}^i \iint_{Q_i \cap Q_j} u_{i+1} b(x, t) |\nabla u_j|^\beta dx dt \\ &\leq 2^\beta U \|b(x, t)\|_{L_r(Q_i)} \|u_{i+1}\|_{L_t(Q)}^{(p-\beta)(n+p)/(n+\lambda)} \|\nabla u_{i+1}\|_{L_p(Q)}^\beta \\ &\quad + C(m, U) \|b(x, t)\|_{L_r(Q_i)} \|u_{i+1}\|_{L_t(Q)}^{(p-\beta)(n+p)/(n+\lambda)} \sum_{j=0}^i \|\nabla u_j\|_{L_p(Q)}^\beta \\ &\leq 2^\beta U 2^\varepsilon C \|u_{i+1}\|_Q + C(m, U) \|u_{i+1}\|_Q^{(p-\beta)/p} \sum_{j=0}^i \|u_j\|_Q^{\beta/p}; \end{aligned}$$

in the deduction of (28) we have used (24) and for the sake of simplicity we have absorbed the norm $\|b(x, t)\|_{L_r(Q)}$ into the constant $C(m, U)$, the U is appearing in (19). Similarly

$$\begin{aligned}
 IV &= \int \int_{Q \cap \{u > h_{i+1}\}} \sum_{j=0}^i u_j b(x, t) |\nabla u|^\beta dx dt \\
 &\leq 2^\beta \int \int_{Q \cap \{u > h_{i+1}\}} \sum_{j=0}^i u_j b(x, t) \left[|\nabla u_{i+1}|^\beta + \left(\sum_{j=0}^i |\nabla u_j| \right)^\beta \right] dx dt \\
 (29) \quad &\leq 2^\beta U \|b(x, t)\|_{L_r(Q)} \sum_{j=0}^i \|u_j\|_{L_l(Q)}^{(p-\beta)(n+p)/(n+\lambda)} \|\nabla u_{i+1}\|_{L_p(Q)}^\beta \\
 &\quad + (2m)^\beta U \|b(x, t)\|_{L_r(Q)} \sum_{j=0}^i \|u_j\|_{L_l(Q)}^{(p-\beta)(n+p)/(n+\lambda)} \|\nabla u_j\|_{L_p(Q)}^\beta \\
 &\leq 1/4 \|u_{i+1}\|_Q + C(m, U) \sum_{j=0}^i \|u_j\|_Q.
 \end{aligned}$$

Now we see from (28) that if at the beginning we take ε so small that

$$(30) \quad 2^{\beta+1} CU \varepsilon \leq 1/4,$$

then from (26)–(29) we infer

$$(31) \quad \|u_{i+1}\|_Q \leq C(m, U) \left\{ \sum_{j=0}^i \|u_j\|_Q + F(k) \right\}, \quad i = 1, 2, \dots, m.$$

Because of (30) the ε is controlled by C, U and β . By the definition of Q_i we have

$$m(\varepsilon/N)^r \leq \sum_{i=1}^m |Q_i| \leq |Q|.$$

So, m is finite and is also controlled by C, U and β . (31) is now written as

$$(32) \quad \|u_{i+1}\|_Q \leq C(U) \left\{ \sum_{j=0}^i \|u_j\|_Q + F(k) \right\}, \quad i = 1, 2, \dots, m.$$

Combining this with (21), the latter can be written as

$$(33) \quad \|u_0\|_Q \leq CF(h_0) \leq CF(k),$$

we deduce by induction that

$$(34) \quad \|u_i\|_Q \leq C(U)F(k), \quad i = 1, 2, \dots, m+1.$$

Thus

$$(35) \quad \begin{aligned} \|(u-k)^+\|_{L_l(Q)}^{q/l} &\leq C\|(u-k)^+\|_Q \leq C\|\sum_{i=0}^{m+1} u_i\|_Q \\ &\leq C\sum_{i=0}^{m+1} \|u_i\|_Q \leq C(U)F(k) \\ &\leq C(U)F\|(u-k)^+\|_{L_l(Q)}|Q \cap \{u > k\}|^{1-1/l}, \end{aligned}$$

i.e.

$$\begin{aligned} \|(u-k)^+\|_{L_l(Q)} &\leq C(U)F^{1/(q-1)}|Q \cap \{u > k\}|^{(1-1/l)/(q-1)}, \\ \int \int_Q (u-k)^+ dx dt &\leq C(U)F^{1/(q-1)}|Q \cap \{u > k\}|^{1+\tau}, \quad \forall k \geq k_0 \end{aligned}$$

where $\tau = p/(n(p-1) + p(\lambda-1)) > 0$. By Lemma 2 we have

$$(36) \quad \begin{aligned} \operatorname{ess\,sup}_Q u^+ &\leq k_0 + (1+1/\tau)C(U)F^{1/(q-1)}|Q|^\tau \\ &\leq \max\{M, k(\theta)\} + C(U)F^{1/(q-1)} \end{aligned}$$

(for simplicity we have absorbed $(1+1/\tau)|Q|^\tau$ into the constant $C(U)$). From (36), it follows (7) by letting $\theta \rightarrow 0$.

In particular, if $1 < p < 2n/(n+\lambda-2)$, from (36) and (11) we have

$$\operatorname{ess\,sup}_Q u^+ \leq \max\{M, k(\theta)\} = M \quad \text{as } M \geq k(0) = (\|f\|_{L_\infty(Q)}/\mu)^{1/(\alpha-1)}$$

and

$$(37) \quad \begin{aligned} \operatorname{ess\,sup}_Q u^+ &\leq k(\theta) + C(U)F^{1/(q-1)} \\ &= (\|f\|_{L_\infty(Q)}/(\mu(1+2\theta)))^{1/(\alpha-1)} \\ &\quad + C(U)(\|f\|_{L_\infty(Q)}2\theta/(1+2\theta))^{1/(q-1)} \quad \text{as } M < k(0). \end{aligned}$$

Observing $q = p(n+\lambda)/(n+p) \in (1, 2)$ as $1 < p < 2n/(n+\lambda-2)$, we can show that for $\theta > 0$ small enough, which depends on $C(U)$, $\|f\|_{L_\infty(Q)}$, μ , λ , α and p ,

$$\begin{aligned} &(\|f\|_{L_\infty(Q)}/\mu(1+2\theta))^{1/(\alpha-1)} + C(U)(\|f\|_{L_\infty(Q)}2\theta/(1+2\theta))^{1/(q-1)} \\ &\leq (\|f\|_{L_\infty(Q)}/\mu(1+\theta))^{1/(\alpha-1)} < (\|f\|_{L_\infty(Q)}/\mu)^{1/(\alpha-1)}. \end{aligned}$$

The conclusion (8) then follows.

Finally, (37) can be written as

$$\operatorname{vrai\,max}_Q u^+ \leq (\|f\|_{L_\infty(Q)}/\mu(1 + 2\theta))^{1/(q-1)} + C(\|f\|_{L_\infty(Q)}2\theta/(1 + 2\theta))^{1/(q-1)}$$

provided $\alpha = q$ and $\beta = \beta_0$ owing to the fact that

$$U = \|u\|_{L_l(Q)}^{1-(p-\beta)(n+p)/(n+\lambda)} = 1 \quad \text{as } \beta = \beta_0.$$

For $q \in (1, 2)$ and θ small enough (θ is now independent of u) we have

$$C(2\theta/(1 + 2\theta))^{1/(q-1)} \leq (\mu(1 + \theta))^{-1/(q-1)} - (\mu(1 + 2\theta))^{-1/(q-1)}$$

and then the conclusion (9) follows. □

PROOF OF THEOREM 2: Under the assumptions of Theorem 2 instead of (11) we have

$$(11)' \quad f(x, t) - \mu|u|^{\alpha-2}u \leq f(x, t) \quad \text{on } Q \cap \{u > k\}$$

and then (35) is replaced by

$$\begin{aligned} \|(u - k)^+\|_{L_l(Q)}^{q/l} &\leq C \iint_Q (u - k)^+ f(x, t) \, dx \, dt \\ &\leq C\|f\|_{L_s(Q)}\|(u - k)^+\|_{L_l(Q)}|Q \cap \{u > k\}|^{1-1/l-1/s}, \end{aligned}$$

where the constant C is independent of u owing to $\beta = \beta_0$. Then it follows

$$\iint_Q (u - k)^+ \, dx \, dt \leq C\|f\|_{L_s(Q)}^{1/(q-1)}|Q \cap \{u > k\}|^{1+\tau_1},$$

where $\tau_1 = (p - (n + p)/s)/(n(p - 1) + p(\lambda - 1)) > 0$. By the use of Lemma 2 we infer (10). □

REFERENCES

- [1] Lions J.L., *Quelques methods de resolution des problèmes aux limites non linéaires*, Dunod. Gauthier-Villars, Paris, 1969.
- [2] Alt H.W., Luckhaus S., *Quasilinear elliptic-parabolic differential equations*, Math. Z. **183** (1983), 311–341.
- [3] Blanchard D., Francfort G.A., *Study of a doubly nonlinear heat equations with no growth assumptions on the parabolic term*, SIAM J. Math. Anal. **19** (1988), 1032–1057.
- [4] Liang Xiting, Liang Xuexin, *Boundedness properties of solutions doubly nonlinear parabolic equations*, Acta Math. Sci., English ed. 1994, supplement, pp. 110–117.
- [5] Ladyzenskaja O.A., Solonnikov V.A., Ural'ceva N.N., *Linear and quasilinear equations of parabolic type*, Transl. Math. Monographs # 23, AMS, Providence, Rhode Island, 1968.

- [6] Ladyzenskaja O.A., Ural'ceva N.N., *Linear and quasilinear elliptic equations*, Academic Press. Beijing, 1987 (Chinese version).
- [7] Miranda C., *Alcune osservazioni sulla maggiorazione in L^{ν} delle soluzioni deboli delle equazioni ellittiche del secondo ordine*, Ann. Mat. Pura Appl. **61** (1963), 151–169.
- [8] Yu Minggu, Liang Xiting, *A new estimate to the maximum of generalized solutions of quasilinear elliptic and parabolic equations* (in Chinese), J. Shanxi Univ. (Nat. Sci. Ed.) **3** (1988), 7–12.
- [9] Lu Youwen, Liang Xiting, *On a priori estimate of maximum modulus for generalized solutions of quasilinear elliptic equations* (in Chinese), J. Tianjin Normal Univ. (Nat. Sci. Ed.) **1** (1989), 5–14.
- [10] Liang Xiting, *An improvement to estimate of maximum modulus of generalized solutions of linear elliptic and parabolic equations of second order* (in Chinese), Acta Sci. Nat. Univ. Sunyatseni **28.2** (1989), 78–80.

DEPARTMENT OF MATHEMATICS, ZHONGSHAN UNIVERSITY, GUANGZHOU 510275, CHINA

DEPARTMENT OF MATHEMATICS, TIANJIN UNIVERSITY, TIANJIN 300072, CHINA

(Received August 28, 1995)