

Connected transversals to subnormal subgroups

TOMÁŠ KEPKA, JON D. PHILLIPS

Abstract. Subnormal subgroups possessing connected transversals are briefly discussed.

Keywords: subgroup, transversal

Classification: 20F22

In [8] J.D.H. Smith introduced the notion of a stably nilpotent quasigroup, showing that a quasigroup Q is stably nilpotent if and only if the inner permutation groups of Q are subnormal in the multiplication group of Q . Generalizing this for abstract groups, we come by groups which are, in a certain sense, relatively nilpotent with respect to a subgroup. The present short note collects some basic information on such groups.

1. Preliminaries

1.1. Let H be a subgroup of a group G . Then $L_G(H)$ denotes the core and $N_G(H)$ the normalizer of H in G . Further, $N_{G,0}(H) = H$ and $N_{G,n+1}(H) = N_G(N_{G,n}(H))$ for every $n \geq 0$.

The subgroup H is said to be subnormal of depth at most $n \geq 0$ in G if there are subgroups H_0, H_1, \dots, H_n of G such that $H_0 = H$, and $H_n = G$ and H_i is normal in H_{i+1} for every $0 \leq i \leq n - 1$.

1.2. Let G be a group. For $n \geq 0$, $Z_n(G)$ denotes the n th member of the usual central series. That is, $Z_0(G) = 1$, and $Z_{n+1}(G)/Z_n(G) = Z(G/Z_n(G))$.

Now, let H be a subgroup of G . We define two series of normal subgroups of G : $Z_{H,0}(G) = Z_{H,0}^*(G) = L_G(H)$, $Z_{H,n}(G) \subseteq Z_{H,n+1}^*(G)$ and $Z_{H,n+1}^*(G)/Z_{H,n}(G) = Z(G/Z_{H,n}(G))$, $Z_{H,n+1}(G) = L_G(H \cdot Z_{H,n+1}^*(G))$.

1.3 Remark. (i) A subgroup H is subnormal of depth at most $n \geq 0$ in a group G , provided that $N_{G,n}(H) = G$. The converse is not true in general (see, e.g., 4.1).

(ii) If G is a finite group, then subnormal subgroups form a sublattice in the lattice of all subgroups of G (see, e.g., [6, Theorem 6.5]). This is not true in general ([7, §13.1, p. 375]), albeit subnormal subgroups of arbitrary (i.e., even infinite) groups are closed under finite intersections.

2. Technical results

2.1 Lemma. *Let H be a subgroup of a group G . Then:*

- (i) $L_G(H) = Z_{H,0}(G) \subseteq Z_{H,1}(G) \subseteq Z_{H,2}(G) \subseteq \dots$;
- (ii) $L_G(H) = Z_{H,0}^*(G) \subseteq Z_{H,1}^*(G) \subseteq Z_{H,2}^*(G) \subseteq \dots$;
- (iii) $Z_{H,n}(G) \subseteq Z_{H,n+1}^*(G) \subseteq Z_{H,n+1}(G) \subseteq Z_{H,n+2}^*(G) \subseteq \dots$ for every $n \geq 0$;
- (iv) $Z_{H,n}(G) \subseteq L_G(N_{G,n}(H))$ for every $n \geq 0$.

PROOF: The first three assertions are clear from definition 1.2, (iv) is clear for $n = 0$, and we shall proceed further by induction.

Let $f : G \rightarrow \overline{G} = G/Z_{H,n}(G)$, $g : G \rightarrow \tilde{G} = G/L_G(N_{G,n}(H))$ and $h : \overline{G} \rightarrow \tilde{G}$ denote the natural projections, $g = hf$. Then $Z_{H,n+1}^*(G) = f^{-1}(Z(\overline{G})) \subseteq g^{-1}(Z(\tilde{G})) = K$, $HK \subseteq N_{G,n}(H)K \subseteq N_G(N_{G,n}(H)) = N_{G,n+1}(H)$ and $Z_{H,n+1}(G) = L_G(H \cdot Z_{H,n+1}^*(G)) \subseteq L_G(HK) \subseteq L_G(N_{G,n+1}(H))$. □

2.2 Lemma. *Let $H \subseteq K \subseteq G$ be subgroups of a group G . Then $Z_{H,n}(G) \subseteq Z_{K,n}(G)$ and $Z_{H,n}^*(G) \subseteq Z_{K,n}^*(G)$ for every $n \geq 0$.*

PROOF: By induction on n (see the proof of 2.1 (iv)). □

2.3 Lemma. *Let H be a subgroup of a group G . Then $Z_n(G) \subseteq Z_{H,n}^*(G) \subseteq Z_{H,n}(G)$ for every $n \geq 0$.*

PROOF: Clearly, $Z_n(G) \subseteq Z_{1,n}^*(G) \subseteq Z_{1,n}(G)$ and we can use 2.2. □

2.4 Lemma. *Let H be a subgroup of a group G . Then:*

- (i) $Z_{H,0}(G) = G$ iff $H = G$;
- (ii) $Z_{H,1}(G) = G$ iff $G' \subseteq H$;
- (iii) $Z_{H,n}(G) = G$ for $n \geq 0$ iff $G = H \cdot Z_{H,n}^*(G)$;
- (iv) if G is nilpotent of class at most $n \geq 0$, then $Z_{H,n}(G) = G$;
- (v) if $Z_{H,n}(G) = G$ for $n \geq 0$, then $N_{G,n}(H) = G$ (and hence H is subnormal of depth at most n in G).

PROOF: The first assertions are easy, (iv) follows from 2.3, and (v) follows from 2.1 (iv). □

2.5 Lemma. *Let H be a subgroup of a group G such the $L_G(H) = 1$. Then:*

- (i) $Z_{H,1}^*(G) = Z(G)$ and $Z_{H,1}(G) = L_G(HZ(G))$;
- (ii) $Z_{H,1}(G) = G$ iff G is abelian;
- (iii) $Z_{H,2}(G) = G$ iff $G' \subseteq HZ(G)$.

PROOF: Obvious. □

2.6 Lemma. *Let H be a subgroup of a group G . Then:*

- (i) $HZ_{H,n}(G) = HZ_{H,n}^*(G)$ for every $n \geq 0$;
- (ii) if K is a subgroup conjugate to H , then $Z_{H,n}(G) = Z_{K,n}(G)$ and $Z_{H,n}^* = Z_{K,n}^*(G)$ for every $n \geq 0$.

PROOF: The assertions follow easily from definition 1.2. □

2.7 Proposition. *Let H be a subgroup of a group G . The following conditions are equivalent for $n \geq 1$:*

- (i) $Z_{H,n}^*(G) = G$;
- (ii) $Z_{H,n}(G) = G$;
- (iii) $HZ_{H,n}(G) = G$;
- (iv) $HZ_{H,n}^*(G) = G$;
- (v) $G' \subseteq Z_{H,n-1}(G)$;
- (vi) $G' \subseteq HZ_{H,n-1}(G)$;
- (vii) $G' \subseteq HZ_{H,n-1}^*(G)$.

PROOF: (i) implies (ii) by 2.1 (iii); (ii) implies (iii) and (v) implies (vi) trivially; (iii) implies (iv) and (vi) implies (vii) by 2.6 (i).

We now show (iv) implies (v). Put $N = Z_{H,n-1}(G)$. We have $\overline{G} = G/N = HZ_{H,n}^*(G)/N = \overline{H}Z(\overline{G})$, and hence $(\overline{G})' \subseteq \overline{H}$, $G' \subseteq HN = HZ_{H,n-1}^*(G)$ and $N = L_G(HZ_{H,n-1}^*(G)) = HZ_{H,n-1}^*(G)$. Consequently $G' \subseteq N$. Finally, we show (vii) implies (i). Since $G' \subseteq HZ_{H,n-1}^*(G)$, we have $Z_{H,n-1}(G) = HZ_{H,n-1}(G)$, $G' \subseteq Z_{H,n-1}(G)$ and $Z_{H,n}^*(G) = G$ (see 1.2). □

2.8. Let H be a subgroup of a group G , $n \geq 0$, $N = Z_{H,n}(G)$, $N^* = Z_{H,n}^*(G)$, $\overline{G} = G/N$, and $\overline{H} = HN/N \subseteq \overline{G}$.

- (i) $HN = HN^*$, $N = L_G(HN^*) = L_G(HN)$ and this implies that $L_{\overline{G}}(\overline{H}) = 1$ and $\overline{H} \cong H/H \cap N$.
- (ii) $Z_{H,n+1}^*(G)/N = Z(\overline{G}) = Z_{\overline{H},1}^*(\overline{G})$, $Z_{H,n+1}(G) = L_G(H \cdot Z_{H,n+1}^*(G))$ and $Z_{H,n+1}(G)/N = L_{\overline{G}}(\overline{H}Z(\overline{G})) = Z_{\overline{H},1}(\overline{G})$.
- (iii) $Z_{H,n+m}^*(G)/N = Z_{\overline{H},m}^*(\overline{G})$ and $Z_{H,n+m}(G)/N = Z_{\overline{H},m}(\overline{G})$ for every $m \geq 1$.

2.9. Let H be a subgroup of a group G . Put $H_n = H \cap Z_{H,n}(G)$ for every $n \geq 0$. Then $L_G(H) = H_0 \subseteq H_1 \subseteq H_2 \subseteq \dots$ and H_n is normal in G .

2.10 Lemma. *Let H be a subgroup of a group G such that $L_G(H) = 1$ and let $\alpha = [G : HZ(G)]$. Then:*

- (i) $Z_{H,1}(G) = L_G(HZ(G))$ can be embedded into the Cartesian product of α copies of $Z(G)$;
- (ii) $Z_{H,1}(G)$ is an abelian group;
- (iii) H_1 (see 2.9) can be embedded into the Cartesian product of $\alpha - 1$ copies of $Z(G)$ ($\alpha - 1 = \alpha$ for α infinite).

PROOF: Put $N = Z_{H,1}(G)$. For every $x \in G$, $N = N^x = L_G(H^x \cdot Z(G))$, $H^x \cap Z(G) \subseteq L_G(H^x) = L_G(H) = 1$, $H^x \cdot Z(G)$ is the direct product of H^x and $Z(G)$ and consequently the restriction f_x of the natural projection $H^x \cdot Z(G) \rightarrow Z(G)$ to N is a homomorphism of N onto $Z(G)$ (we have $Z(G) \subseteq N$).

Now, let A be a right transversal to $HZ(G)$ in G such that $1 \in A$. Define a homomorphism $f : N \rightarrow \prod_{\alpha} Z(G)$ by $f(u) = \prod_{a \in A} f_a(u)$, $u \in N$. If $u \in \text{Ker}(f)$, then $aua^{-1} \in H$ for every $a \in A$. Consequently, $u \in H$ and if $x \in G$, $x = zva$, $a \in A$, $v \in H$, $z \in Z(G)$, then $xux^{-1} = zvaua^{-1}v^{-1}z^{-1} = vaua^{-1}v^{-1} \in H$. Thus $u \in L_G(H) = 1$ and we have proved that f is injective. Finally, for $g = \prod_{a \neq 1} f_a$ we get $\text{Ker}(g) \cap H = 1$, and hence $g|H_1$ is injective. □

2.11 Proposition. *Let H be a subgroup of a group G and let $\alpha_n = [G : H \cdot Z_{H,n+1}(G)]$ for every $n \geq 0$. Then $Z_{H,n+1}(G)/Z_{H,n}(G)$ is an abelian group which can be embedded into the Cartesian product of α_n copies of $Z(G/Z_{H,n}(G)) = Z_{H,n+1}^*(G)/Z_{H,n}(G)$.*

PROOF: The result follows by an easy combination of 2.10 and 2.8 (i),(ii). □

2.12 Corollary. *Let H be a subgroup of a group G such that $Z_{H,n}(G) = G$ for some $n \geq 0$. If H is soluble of derived length $m \geq 0$, then G is also soluble and its derived length is at most $n + m$.*

2.13 Lemma. *Let H be a subgroup of a group G such that $Z_{H,2}(G) = G$. Then $H \subseteq L_G(H)$.*

PROOF: By 2.10, $H/L_G(H)$ is abelian. □

2.14 Proposition. *Let H be a subgroup of a finite group G such that $[G : H]$ is a power of a prime p and $L_G(H)$ is a p -group. Then $G = Z_{H,n}(G)$ for some $n \geq 0$ iff G is a p -group.*

PROOF: If G is a p -group, then G is nilpotent and our result follows from 2.3. Now assume that $Z_{H,n}(G) = G$. We shall proceed by induction on $\text{card}(G)$. Further, considering the factor $G/L_G(H)$, we can restrict ourselves to the case $L_G(H) = 1$. Then $H \cap Z(G) = 1$, $[HZ(G) : H] = \text{card}(Z(G))$, and hence $Z(G)$ is a p -group. From this, $N = Z_{H,1}(G)$ is a p -group by 2.10 (i). Since $N \neq 1$ (otherwise $G = 1$), G/N is a p -group by induction. □

2.15. Let H be a subgroup of a group G such that $G/Z_{H,n}(G)$ is a two element group for some $n \geq 0$.

- (i) If $n = 0$, then $G/L_G(H)$ is a two element group, which means that H is normal and of index 2 in G .
- (ii) Assume that $n \geq 1$. Clearly, $Z_{H,n+1}(G) = Z_{H,n+1}^*(G) = G$ and $G' \subseteq Z_{H,n}(G) = H \cdot Z_{H,n}^*(G)$. Put $N = Z_{H,n-1}(G)$, $\overline{G} = G/N$ and $\overline{G} = HN/N = HZ_{H,n-1}^*(G)/L_G(HZ_{H,n-1}^*(G))$. We have $L_{\overline{G}}(\overline{H})=1$, $Z(\overline{G}) =$

$Z_{H,n}^*(G)/N, (\overline{G})' \subseteq Z_{H,n}(G)/N = \overline{H} \cdot Z(\overline{G})$ and $\overline{G}/\overline{H}Z(\overline{G}) \cong G/Z_{H,n}(G)$, so that $\overline{G}/\overline{H}Z(\overline{G})$ is a two element group.

- (iii) Assume that $n = 1$ and that $L_G(H) = 1$ (cf. (ii)). Then $Z_{H,2}(G) = Z_{H,2}^*(G)$ and $G' \subseteq Z_{H,1}(G) = HZ(G)$. Take $w \in G - HZ(G)$ and put $W = Z(G) \cup wZ(G)$. Then $w^2 = uz$ for suitable $u \in H, z \in Z$ and $w^{-1}uw = w^{-1}w^2z^{-1}w = u$. This implies that $u \in L_G(H) = 1$, so that $w^2 \in Z(G)$ and we see that W is an abelian subgroup of $G, W \cap H = 1$ and $G = HW$.

3. Connected transversals to subnormal subgroups

3.1. In this section, let H be a subgroup of a group G such that there exist H -connected transversals A, B to H in G (i.e., A, B are left transversals and $[A, B] \subseteq H$).

3.2 Lemma.

- (i) $HZ_{H,n}(G) = HZ_{H,n}^*(G) = N_{G,n}(H)$ for every $n \geq 0$.
- (ii) $Z_{H,n}(G) = L_G(N_{G,n}(H))$ for every $n \geq 0$.

PROOF: This is clear for $n = 0$ and we shall proceed by induction on n .

Put $N = Z_{H,n}(G)$ and consider the factors $\overline{G} = G/N$ and $\overline{H} = HN/N$. Then $L_{\overline{G}}(\overline{H}) = 1$, and so $N_{\overline{G}}(\overline{H}) = \overline{H}Z(\overline{G})$ by [3, Proposition 2.7]. This implies that $N_G(HN) = HZ_{H,n+1}^*(G)$. However, $HN = N_{G,n}(H)$ by the induction and we have $N_{G,n+1}(H) = HZ_{H,n}^*(G) = HZ_{H,n}(G)$ (2.6 (ii)). The rest is clear. \square

3.3 Proposition. *The following conditions are equivalent for $n \geq 1$:*

- (i) $Z_{H,n}(G) = G$;
- (ii) $HZ_{H,n-1}(G)$ is normal in G ;
- (iii) $H \subseteq Z_{H,n-1}(G)$;
- (iv) $H_{n-1} = H$ (see 2.9);
- (v) H is subnormal of depth at most n in G ;
- (vi) $N_{G,n}(H) = G$;
- (vii) $N_G(H)$ is subnormal of depth at most $n - 1$ in G .

PROOF: (i) implies (ii) by 2.7 (ii),(vi) (in fact, $G' \subseteq HZ_{H,n-1}(G)$); (ii) implies (iii), since $Z_{H,n-1}(G) = L_G(HZ_{H,n-1}(G))$; (iii) implies (iv) trivially; (iv) implies (ii), since $Z_{H,n-1}(G) = L_G(HZ_{H,n-1}(G))$; (i) implies (vi) by 2.1 (iv); (vi) implies (vii) and (vii) implies (v) trivially; (vi) implies (i) by 3.2 (ii).

We now show (ii) implies (i). The existence of H -connected transversals easily yields that $G' \subseteq HZ_{H,n-1}(G)$ (consider the factor $G/Z_{H,n-1}(G)$), and the result follows from 2.7.

We proceed by induction on n to show (v) implies (vi). If $n = 1$, then H is normal in G and (vi) is clear. Let $n \geq 2$ and let $L_G(H) = 1$ (considering the factor $G/L_G(H)$, we can restrict ourselves to this case). There is a subgroup K of G such

that H is a normal subgroup of K and K is subnormal of depth at most $n-1$ in G . Put $L = L_G(K)$, $\overline{G} = G/L$ and $\overline{K} = K/L$. Then $L_{\overline{G}}(\overline{K}) = 1$ and \overline{K} is subnormal of depth at most $n-1$ in \overline{G} . Consequently, $N_{\overline{G},n-1}(\overline{K}) = \overline{G}$ and $N_{G,n-1}(K) = G$. On the other hand, $K \subseteq N_G(H) = HZ(G)$ ([3, Proposition 2.7]), and hence $N_G(H) = KZ(G)$ is normal in $N_G(K)$. We have proved that $N_G(H)$ is subnormal of depth at most $n-1$ in G . Using the induction hypothesis again (for $N_G(H)$), we get $N_{G,n}(H) = N_{G,n-1}(N_G(H)) = G$. \square

3.4 Proposition. *Suppose that $G = \langle A, B \rangle$ and that $G/Z_{H,n}(G)$ is a two element group for some $n \geq 0$. Then $n = 0$ and H is a normal subgroup of index 2 in G .*

PROOF: Assume on the contrary, $n \geq 1$. With respect to 2.15, we can in fact assume that $n = 1$ and $L_G(H) = 1$. Then $Z_{H,1}(G) = HZ(G)$ and $H \cap Z(G) = 1$. By [1, Lemma 1.4], $Z(G) \subseteq A \cap B$. Now, let $a \in A$ and $z \in Z(G)$. Then $az = bu$ for some $b \in A$ and $u \in H$. We have $u = b^{-1}az$ and $c^{-1}uc = c^{-1}b^{-1}cb \cdot b^{-1}c^{-1}ac \cdot z = c^{-1}b^{-1}cb \cdot b^{-1}az \cdot a^{-1}c^{-1}ac \in H$ for every $c \in B$. This shows that $u \in L_G(H) = 1$ and $az = b \in A$. Now, since $[G : HZ(G)] = 2$, it is clear that $A = Z(G) \cup aZ(G)$ for each $a \in A - Z(G)$. Quite similarly, $B = Z(G) \cup bZ(G)$ for each $b \in B - Z(G)$. In particular, both A and B are abelian subgroups of G (see 2.15 (iii)).

Finally, let $a \in A$. Then $a^{-1}b \in H$ for some $b \in B$ and, for every $c \in B$, $c^{-1}a^{-1}bc = c^{-1}a^{-1}ca \cdot a^{-1}b \in H$. Thus $a^{-1}b \in L_G(H) = 1$ and $a = b \in B$. We have proved that $A = B$ and consequently $G = \langle A, B \rangle = A$ is an abelian group, $H = 1$, $Z_{H,1}(G) = G$ and $G/Z_{H,1}(G)$ is trivial, a contradiction. \square

3.5 Lemma. *Suppose that $L_G(H) = 1$, H is not abelian, every proper factor group of H is cyclic and that $G = \langle A, B \rangle$. Then $Z_{H,n}(G) \neq G$ for every $n \geq 0$, i.e., H is not subnormal in G (see 3.3).*

PROOF: Put $N = Z_{H,1}(G) (= L_G(HZ(G)))$, $\overline{G} = G/N$ and $\overline{H} = HN/N \cong H/H_1$, $H_1 = H \cap N$. If $H_1 \neq 1$, then \overline{H} is cyclic, and so $\overline{A} = \overline{B}$ is an abelian subgroup of \overline{G} by [1, Corollary 2.3]. However, this implies that $\overline{G} = \overline{A}$ is an abelian group, $\overline{H} = 1$, $H \subseteq N = HZ(G)$ and $H = H_1$ is abelian by 2.10 (iii), which is a contradiction.

We have proved that $H_1 = 1$, so that $N = H_1Z(G) = Z(G)$ and $\overline{H} \cong H$. Proceeding by induction, we get $Z_{H,m}(G) = Z_m(G)$ for every $m \geq 0$. Now, if $Z_{H,n}(G) = G$ for some $n \geq 0$, then G (and hence H) is nilpotent. But in such a case, $Z(H) \neq 1$, $H/Z(H)$ is cyclic and this implies that H is abelian a contradiction. \square

3.6 Proposition. *Suppose that every proper factorgroup of H is cyclic, that H is subnormal in G and that $G = \langle A, B \rangle$. Then $G' \subseteq N_G(H)$ and H is subnormal depth at most 2 in G . Moreover, if H is not abelian, then $G' \subseteq H$ and H is normal in G .*

PROOF: First, assume that $L_G(H) \neq 1$. Then $\overline{H} = H/L_G(H)$ is a cyclic subgroup of $\overline{G} = G/L_G(H)$ and $G' \subseteq H$ by [1, Theorem 2.2].

Next, let $L_G(H) = 1$. Then H is abelian by 3.5 and if H is cyclic, then we can use [1, Theorem 2.2] again to show that $H = 1$ and G is abelian. Finally, if H is not cyclic, then $H \cong Z_p^{(2)}$ for a prime p and the result follows from [5, Lemma 4.2]. \square

3.7 Remark. According to [2], G is soluble, provided that G is finite and $H \cong S_3$. On the other hand, by 3.5, if $L_G(H) = 1$ and $G = \langle A, B \rangle$, then H is not subnormal in G .

3.8 Proposition. Suppose that $L_G(H) = 1$ and G is nilpotent of class at most 2. Then $[A, B] = 1$ and A, B are isomorphic subgroups of G .

PROOF: $[A, B] \subseteq H \cap G' \subseteq H \cap Z(G) \subseteq L_G(H) = 1$. The rest follows from [4, Lemma 2.3]. \square

4. Examples

4.1. Let G be the subgroup of S_6 (the symmetric group on $\{1, 2, \dots, 6\}$) generated by the following permutations: $(1\ 2), (3\ 4), (5\ 6), (1\ 3)(2\ 4), (1\ 3\ 5)(2\ 4\ 6)$. Further, let $K = \langle (1\ 2), (3\ 4), (5\ 6) \rangle \subseteq G$ and $H = \langle (1\ 2) \rangle \subseteq K$. Then H is normal in K , K is normal in G , $\text{card}(G) = 48$, $K \cong Z_2^{(3)}$, $H \cong Z_2$, $L_G(H) = 1$ and H is subnormal of depth 2 in G . On the other hand, $N_G(H) = \langle K, (3\ 5)(4\ 6) \rangle$, $\text{card}(N_G(H)) = 16$, $K = L_G(N_G(H))$, $N_{G,2}(H) = N_G(N_G(H)) = N_G(H)$, $G/K \cong S_3$ and $Z(G) = 1$. Now, $Z_{H,n}(G) \neq G$ for every $n \geq 0$ and there exist no H -connected transversals to H in G (see 2.4(v) and 3.3).

4.2. Let G be the subgroup of S_{18} generated by $A = \{\text{id}, (1\ 2)(3\ 10\ 15\ 4\ 9\ 16)(5\ 12\ 17\ 6\ 11\ 18)(7\ 8)(13\ 14), (1\ 3\ 11\ 7\ 9\ 17\ 13\ 15\ 5)(2\ 10\ 18)(4\ 12\ 14)(6\ 8\ 16), (1\ 4\ 11\ 14\ 3\ 12\ 7\ 10\ 17\ 2\ 9\ 18\ 13\ 16\ 5\ 8\ 15\ 6), (1\ 5\ 10\ 14\ 6\ 9\ 7\ 11\ 16\ 2\ 12\ 15\ 13\ 17\ 4\ 8\ 18\ 3), (1\ 6\ 10\ 7\ 12\ 16\ 13\ 18\ 4)(2\ 11\ 15)(3\ 8\ 17)(5\ 9\ 14), (1\ 7\ 13)(2\ 8\ 14)(3\ 9\ 15)(4\ 10\ 16)(5\ 11\ 17)(6\ 12\ 18), (1\ 8\ 13\ 2\ 7\ 14)(3\ 16\ 9\ 4\ 15\ 10)(5\ 18\ 11\ 6\ 17\ 12), (1\ 9\ 5\ 7\ 15\ 11\ 13\ 3\ 17)(2\ 16\ 12)(4\ 18\ 8)(6\ 14\ 10), (1\ 10\ 5\ 14\ 9\ 6\ 7\ 16\ 11\ 2\ 15\ 12\ 13\ 4\ 17\ 8\ 3\ 18), (1\ 11\ 4\ 14\ 12\ 3\ 7\ 17\ 10\ 2\ 18\ 9\ 13\ 5\ 16\ 8\ 6\ 15), (1\ 12\ 4\ 7\ 18\ 10\ 13\ 6\ 16)(2\ 17\ 9)(3\ 19\ 11)(5\ 15\ 8), (1\ 13\ 7)(2\ 14\ 8)(3\ 15\ 9)(4\ 16\ 10)(5\ 17\ 11)(6\ 18\ 12), (1\ 14\ 7\ 2\ 13\ 8)(3\ 4)(5\ 6)(9\ 10)(11\ 12)(15\ 16)(17\ 18), (1\ 15\ 17\ 7\ 3\ 5\ 13\ 9\ 11)(2\ 4\ 6)(8\ 10\ 12)(14\ 16\ 18), (1\ 16\ 17\ 14\ 15\ 18\ 7\ 4\ 5\ 2\ 3\ 6\ 13\ 10\ 11\ 8\ 9\ 12), (1\ 17\ 16\ 14\ 18\ 15\ 7\ 5\ 4\ 2\ 6\ 3\ 13\ 11\ 10\ 8\ 12\ 9), (1\ 18\ 16\ 7\ 6\ 4\ 13\ 12\ 10)(2\ 5\ 3)(8\ 11\ 9)(14\ 17\ 15)\}$ and let H be the stabilizer of 1 in G . Then $L_G(H) = 1$, $\text{card}(H) = 972 = 2^2 3^5$, H is not nilpotent, A is an H -selfconnected transversal to H in $G = \langle A \rangle$, $\text{card}(G) = 17496 = 2^3 3^7$, and $Z_{H,3}(G) = G$ (cf. 2.13).

4.3. Let G be the subgroup of S_6 generated by $A = \{\text{id}, (1\ 2)(3\ 4)(5\ 6), (1\ 3\ 5)(2\ 4\ 6), (1\ 4\ 5\ 2\ 3\ 6), (1\ 5\ 4\ 2\ 6\ 3), (1\ 6\ 4)(2\ 5\ 3)\}$ and let H be the stabilizer of 1 in G . Then $L_G(H) = 1$, $H \cong Z_2^{(2)}$, A is an H -selfconnected transversal to H in $G = \langle A \rangle$, $\text{card}(G) = 24$, $Z_{H,2}(G) = G$, $\text{card}(Z(G)) = 2$, G is not nilpotent, $\text{card}(N_G(H)) = 8$, $N_G(H) = HZ(G) = Z_{H,1}(G) \cong Z_2^{(3)}$ and $G/Z_{H,1}(G) \cong Z_3$ (cf. 2.4(iv) and 3.4).

4.4. Let G be the subgroup of S_6 generated by $A = \{\text{id}, (1\ 2\ 3)(4\ 5\ 6), (1\ 3\ 2)(4\ 5\ 6), (1\ 4)(2\ 6\ 3\ 5), (1\ 5\ 3\ 6)(2\ 4), (1\ 6\ 2\ 5)(3\ 4)\}$ and let H be the stabilizer of 1 in G . Then $L_G(H) = 1$, $H \cong S_3$ is soluble, A is an H-selfconnected transversal to H in $G = \langle A \rangle$, $\text{card}(G) = 36$, $G \neq Z_{H,n}(G)$ for every $n \leq 0$ and H is not subnormal in G (see 3.5).

4.5. Let G be the subgroup of S_4 generated by $(1\ 2)$, $(3\ 4)$, $(1\ 3\ 2\ 4)$, $(1\ 4\ 2\ 3)$, let H be the stabilizer of 1 in G and let $A = \{\text{id}, (1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3)\}$. Then $L_G(H) = 1$, $H \cong Z_2$, A is an H-selfconnected transversal to H in G , $A \cong Z_2^{(2)}$ is a subgroup of G , G is a dihedral eight-element group, $Z_{H,1}(G) \cong Z_2^{(2)}$ and $G/Z_{H,1}(G) \cong Z_2$ (cf. 3.4).

REFERENCES

- [1] Kepka T., Niemenmaa M., *On loops with cyclic inner mapping groups*, Arch. Math. **60** (1993), 233–23.
- [2] Niemenmaa M., *Transversals, commutators and solvability in finite groups*, Bolletino U.M.I. **9-A** (1995), 203–208.
- [3] Niemenmaa M., Kepka T., *On multiplication groups of loops*, J. of Algebra **135** (1990), 112–122.
- [4] Niemenmaa M., Kepka T., *On connected transversals to abelian subgroups in finite groups*, Bull. London Math. Soc. **24** (1992), 343–346.
- [5] Niemenmaa M., Kepka T., *On connected transversals to abelian subgroups*, Bull. Austral. Math. Soc. **49** (1994), 121–128.
- [6] Passman D., *Permutation groups*, Mathematics Lecture Notes Series, W.A. Benjamin, Inc., New York-Amsterdam, 1966.
- [7] Robinson D.J.L., *A course in the theory of groups*, Graduate Text in Mathematics 80 (1982), Springer-Verlag, New York-Heidelberg-Berlin.
- [8] Smith J.D.H., *Multiplication groups of quasigroups*, Preprint 603 (1981), Technische Hochschule, Darmstadt.

DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83, 186 00 PRAHA 8, CZECH REPUBLIC

DEPARTMENT OF MATHEMATICAL SCIENCES, SAINT MARY'S COLLEGE, MORAGA, CALIFORNIA 94575, USA

(Received January 9, 1996)