

## An elementary proof of a theorem on sublattices of finite codimension

MAREK WÓJTOWICZ

*Abstract.* This paper presents an elementary proof and a generalization of a theorem due to Abramovich and Lipecki, concerning the nonexistence of closed linear sublattices of finite codimension in nonatomic locally solid linear lattices with the Lebesgue property.

*Keywords:* linear lattice, Lebesgue property, lattice homomorphism

*Classification:* 46A40, 47B65

In 1990 Y.A. Abramovich and Z. Lipecki proved, by means of Boolean algebra techniques and Liapunov's convexity theorem, the following result ([1, Theorem 2]; cf. [3, Example 27.8]):

*Let  $X = (X, \tau)$  be a Hausdorff locally solid linear lattice such that:*

- (i)  *$X$  is nonatomic and Dedekind complete,*
- (ii)  *$X$  has the Lebesgue property.*

*Then  $X$  contains no proper closed sublattices of finite codimension.*

(By a sublattice of a linear lattice we always mean a *linear* sublattice.  $X$  has the Lebesgue property (or,  $\tau$  is a Lebesgue topology on  $X$ ) provided that for every MS-sequence  $(x_\alpha)$  in  $X$  with  $x_\alpha \downarrow 0$  we have  $x_\alpha \rightarrow 0(\tau)$ . For other undefined notions and basic results concerning linear lattices (= Riesz spaces) in this paper we refer the reader to the monographs [2], [3]).

Here we give an elementary and short proof of a more general result, namely, we show that the two assumptions in the above theorem, i.e. that  $\tau$  is Hausdorff and  $X$  is Dedekind complete are superfluous. It should be noted that every infinite dimensional linear lattice possesses sublattices of arbitrary finite codimension ([1, Theorem 3]).

**Theorem.** *Let  $X$  be a nonatomic linear lattice, and let  $Y$  be a sublattice of  $X$  with  $\dim X/Y < \infty$ . Then  $Y$  is order dense in  $X$ .*

*If, additionally,  $\tau$  is a Lebesgue topology on  $X$ , then  $Y$  is  $\tau$ -dense in  $X$ .*

*In particular, the topological dual  $X'$  of any nonatomic locally solid linear lattice  $(X, \tau)$  with the Lebesgue property is nonatomic (equivalently,  $X$  has no nontrivial continuous Riesz homomorphisms  $X \rightarrow \mathbf{R}$ ).*

Order denseness of  $Y$  in  $X$  is understood in the sense of ([2, Definition 1.9]), i.e. that  $Y_+ \setminus \{0\}$  is cofinal in  $X_+ \setminus \{0\}$ .

PROOF: Let  $Q$  denote the quotient map  $X \rightarrow X/Y$ . Since  $X$  is nonatomic, every principal ideal  $A_e = \{x \in X : |x| \leq \lambda e \text{ for some } \lambda \geq 0\}$ ,  $e \in X^+$ , is of infinite dimension. If  $Y$  were not order dense in  $X$ , then  $A_e \cap Y = \{0\}$  for some  $e \in X^+$ , and hence  $Q$  restricted to  $A_e$  would be a linear isomorphism; thus  $\dim A_e \leq \dim Q(X) < \infty$ , a contradiction. This proves the first part of the theorem, and since for Lebesgue topologies order denseness implies topological denseness, the second part also follows; the particular case is implied by ([2, Theorem 3.13]; [3, Theorem 18.3 (iii)]).  $\square$

**Examples.** 1. Let  $K$  be a topological Hausdorff space. The lattice  $C(K)$  is nonatomic whenever  $K$  has no isolated points, thus every such lattice has the property described in the first part of the Theorem.

2. Let  $S_p$  denote the nonatomic sublattice, consisting of all step functions, of the (nonatomic) lattice  $L_p = L_p(0, 1)$ ,  $0 < p < \infty$ . It is easily seen that  $S_p$  endowed with the  $p$ -norm topology has the Lebesgue property, and hence  $S_p$  possesses the property described in the second part of the Theorem without being even  $\sigma$ -Dedekind complete (compare with (i) above).

3. This example seems to be known; we include it for completeness of the paper. Let  $1 < p < \infty$ . Since every continuous linear functional  $f$  on the lattice  $L_p$  is order continuous ([2, Theorems 9.1, 22.1 and 22.4]), any family of seminorms  $(q_f)$  of the form  $q_f(x) = |f|(|x|)$ ,  $x \in L_p$ , determines a Lebesgue topology on  $L_p$  ([2, p. 40]). This topology is Hausdorff whenever the family  $(q_f)$  is total.

#### REFERENCES

- [1] Abramovich Y.A., Lipceki Z., *On ideals and sublattices in linear lattices and  $F$ -lattices*, Math. Proc. Cambridge Phil. Soc. **108** (1990), 79–87.
- [2] Aliprantis C.D., Burkinshaw O., *Locally Solid Riesz Spaces*, Academic Press, 1978.
- [3] Luxemburg W.A.J., Zaanen A.C., *Riesz Spaces I*, North-Holland, 1971.

INSTITUTE OF MATHEMATICS, PEDAGOGICAL UNIVERSITY, PL. SŁOWIAŃSKI 9,  
65–069 ZIELONA GÓRA, POLAND

(Received July 1, 1997, revised November 4, 1997)