An inequality in Orlicz function spaces with Orlicz norm

Jincai Wang

Abstract. We use Simonenko quantitative indices of an $\mathcal{N}$-function Φ to estimate two parameters q_Φ and Q_Φ in Orlicz function spaces $L^\Phi[0,\infty)$ with Orlicz norm, and get the following inequality: $\frac{B_\Phi}{B_\Phi - 1} \le q_\Phi \le Q_\Phi \le \frac{A_\Phi}{A_\phi - 1}$, where A_Φ and B_Φ are Simonenko indices. A similar inequality is obtained in $L^\Phi[0,1]$ with Orlicz norm.

Keywords: Orlicz spaces, Simonenko indices, $\triangle_2$-condition

Classification: 46B20, 46E30

1. Introduction

Definition 1.1. A function $M : \mathbb{R} \longrightarrow \mathbb{R}$ is called an $\mathcal{N}$-function, if

(i) M is continuous, convex and even;
(ii) $M(u) > 0$ for $u \neq 0$, $M(0) = 0$;
(iii) $\lim\limits_{u\to 0} M(u)/u = 0$, $\lim\limits_{u\to\infty} M(u)/u = \infty$.

Let

$$\Phi(u) = \int_0^{|u|} \phi(t)\,dt \quad \text{and} \quad \Psi(v) = \int_0^{|v|} \psi(s)\,ds$$

be a pair of complementary $\mathcal{N}$-functions. The Orlicz function space is defined as follows: $L^\Phi[0,1] = \{x(t) : x(t)$ is measurable on $[0,1]$ and $\rho_\Phi(\lambda x(t))\,dt < \infty$ for some $\lambda > 0\}$, where $\rho_\Phi(x(t)) = \int_{[0,1]} \Phi(x(t))\,dt$; $L^\Phi[0,\infty) = \{x(t) : x(t)$ is measurable on $[0,\infty)$, $\rho_\Phi(\lambda x(t))\,dt < \infty$ for some $\lambda > 0\}$, and $\rho_\Phi(x(t)) = \int_{[0,\infty)} \Phi(x(t))\,dt$. We define the Orlicz norm on the Orlicz space as

$$\|x\|_\Phi = \inf_{k>0} \frac{1}{k}[1 + \rho_\Phi(kx)].$$

An $\mathcal{N}$-function $\Phi(u)$ is said to satisfy the $\triangle_2$-*condition for small* u (in symbol $\Phi \in \triangle_2(0)$), if there exists $u_0 > 0$ and $C > 0$, such that $\Phi(2u) \le C\Phi(u)$ for $0 \le u \le u_0$. $\Phi(u)$ is said to satisfy the $\triangle_2$-*condition for large* u (in symbol $\Phi \in \triangle_2(\infty)$), if there exists $u_0 > 0$ and $C > 0$ such that $\Phi(2u) \le C\Phi(u)$ for $u \ge u_0$. $\Phi(u)$ is said to satisfy the $\triangle_2$-*condition for all* $u \ge 0$ (in symbol $u \in \triangle_2$), if there exist $C > 0$ such that $\Phi(2u) \le C\Phi(u)$ for $u \ge 0$. An $\mathcal{N}$-function

$\Phi(u)$ is said to satisfy the ∇_2-*condition for small* u (*for large* u, *for all* $u \geq 0$), in symbol $\Phi \in \nabla_2(0)$ ($\Phi \in \nabla_2(\infty)$, $\Phi \in \nabla_2$), if its complementary $\mathcal{N}$-function $\Psi \in \triangle_2(0)$ ($\Psi \in \triangle_2(\infty)$, $\Psi \in \triangle_2$).

The basic results on Orlicz spaces can be found in Krasnosel'skii and Rutickii [2], Lindenstrauss and Tzafriri [3], Rao and Ren [6], Chen [1].

The Simonenko indices of an $\mathcal{N}$-function Φ are defined as

$$A_\Phi = \inf_{t>0} \frac{t\phi(t)}{\Phi(t)}, \qquad B_\Phi = \sup_{t>0} \frac{t\phi(t)}{\Phi(t)}. \tag{1}$$

Simonenko introduced these indices in [9] and [8], and we can find a detailed description in Maligranda [4].

Clearly, $1 \leq A_\Phi \leq B_\Phi \leq \infty$.

Proposition 1.1. *Let* Φ *be an* $\mathcal{N}$*-function. Then*
$\Phi \in \nabla_2 \iff 1 < A_\Phi$*;* $\Phi \in \triangle_2 \iff B_\Phi < \infty$.

The proof of the proposition can be found in Krasnosel'skii and Rutickii [2, p. 24–26].

Lemma 1.2. *Let* Φ *and* Ψ *be a pair of complementary* $\mathcal{N}$*-functions. Then*

$$\frac{1}{A_\Phi} + \frac{1}{B_\Psi} = 1. \tag{2}$$

The proof of Lemma 1.2 can be found in Simonenko [9] or Rao & Ren [6].

The next lemma can be found in [1], [10] or [5].

Lemma 1.3. *Let* $\Phi(u) = \int_0^{|u|} \phi(t)\,dt$ *and* $\Psi(v) = \int_0^{|v|} \psi(s)\,ds$ *be a pair of complementary* $\mathcal{N}$*-functions. We denote*

$$k_x^* = \inf\{k > 0 : \rho_\Psi[\phi(k|x|)] \geq 1\}, \quad k_x^{**} = \sup\{k > 0 : \rho_\Psi[\phi(k|x|)] \leq 1\}.$$

Then $k \in [k_x^*, k_x^{**}]$ *if and only if*

$$\|x\|_\Phi = \frac{1}{k}[1 + \rho_\Phi(kx)].$$

2. Main results

Y. Yan estimated the two parameters Q_Φ and q_Φ in the Orlicz sequence space l^Φ, and got the following result (see [11], [7] or [13]).

Proposition 2.1. *Let Φ and Ψ be a pair of complementary $\mathcal{N}$-functions. Then*

$$\frac{b_\Phi^*}{b_\Phi^*-1} \le q_\Phi \le Q_\Phi \le \frac{a_\Phi^*}{a_\Phi^*-1}\,, \tag{3}$$

where

$$a_\Phi^* = \inf\left\{\frac{t\phi(t)}{\Phi(t)} : 0 < t \le \psi[\Psi^{-1}(1)]\right\},$$
$$b_\Phi^* = \sup\left\{\frac{t\phi(t)}{\Phi(t)} : 0 < t \le \psi[\Psi^{-1}(1)]\right\}.$$

The upper estimate in (3) can also be found in [12]. Now we establish a similar inequality in the Orlicz function space with Orlicz norm. Firstly, we have

Theorem 2.1. *Let Φ, Ψ be a pair of complementary $\mathcal{N}$-functions. For $L^\Phi[0,\infty)$, we denote*

$$Q_\Phi = \sup_{\|x\|_\Phi=1} k_x^{**} = \sup_{\|x\|_\Phi=1}\left\{k>0 : \|x\|_\Phi = \frac{1}{k}(1+\rho_\Phi(kx))\right\},$$
$$q_\Phi = \inf_{\|x\|_\Phi=1} k_x^{*} = \inf_{\|x\|_\Phi=1}\left\{k>0 : \|x\|_\Phi = \frac{1}{k}(1+\rho_\Phi(kx))\right\}.$$

Then

$$A_\Psi = \frac{B_\Phi}{B_\Phi-1} \le q_\Phi \le Q_\Phi \le \frac{A_\Phi}{A_\Phi-1} = B_\Psi, \tag{4}$$

where A_Φ, B_Φ, A_Ψ and B_Ψ are defined by (1).

Proof: The left and right equations in (4) follow from Lemma 1.2. Now we prove

$$q_\Phi \ge \frac{B_\Phi}{B_\Phi-1}\,. \tag{5}$$

For $\Phi \notin \triangle_2$, by Proposition 1.1, we have $B_\Phi = \infty$ or $A_\Psi = 1$. The result is obvious.

For $\Phi \in \triangle_2$, we only prove that for every $x \in L^\Phi[0,\infty)$ which satisfies $\|x\|_\Phi = 1$, we have $k_x^* \ge \frac{B_\Phi}{B_\Phi-1}$. Firstly, we have $\rho_\Psi(\phi(k_x^*|x(t)|)) \ge 1$. In fact, if $\Phi \in \triangle_2$, then $\rho_\Phi[(k_x^*+1)x] < \infty$. So

$$\begin{aligned}\rho_\Psi(\phi((k_x^*+1)|x(t)|)) &\le \rho_\Psi(\phi((k_x^*+1)|x(t)|)) + \rho_\Phi((k_x^*+1)|x(t)|)\\ &= \int_G (k_x^*+1)|x(t)|\cdot\phi((k_x^*+1)|x(t)|)\,dt\\ &\le B_\Phi\rho_\Phi((k_x^*+1)|x(t)|) < \infty.\end{aligned}$$

Choose $k_x^* < k_n < k_x^* + 1$ such that $k_n \searrow k_x^*$. By the right continuity of ϕ and Lebesgue dominated convergence theorem, we have

$$\rho_\Psi(\phi(k_x^*|x(t)|)) = \lim_{n\to\infty} \rho_\Psi(\phi(k_n|x(t)|)) \geq 1.$$

For every $x \in L^\Phi[0,\infty)$ which satisfies $\|x\|_\Phi = 1$, we have

$$\begin{aligned}
1 + \rho_\Phi(k_x^* x) &\leq \rho_\Psi(\phi(k_x^*|x(t)|)) + \rho_\Phi(k_x^*|x(t)|) \\
&= \int_{[0,\infty)} \Psi\{\phi[(k_x^*|x(t)|)]\}\,dt + \int_{[0,\infty)} \Phi(k_x^*|x(t)|)\,dt \\
&= \int_{[0,\infty)} k_x^*|x(t)|\phi(k_x^*|x(t)|)\,dt \\
&\leq B_\Phi \int_{[0,\infty)} \Phi(k_x^*|x(t)|)\,dt = B_\Phi \rho_\Phi(k_x^* x).
\end{aligned}$$

This implies

$$\rho_\Phi(k_x^* x) \geq \frac{1}{B_\Phi - 1}\,. \tag{6}$$

By Lemma 1.3, we get

$$1 = \|x\|_\Phi = \frac{1}{k_x^*}\{1 + \rho_\Phi(k_x^* x)\}.$$

So $\rho_\Phi(k_x^* x) = k_x^* - 1$. By (6)

$$k_x^* \geq \frac{B_\Phi}{B_\Phi - 1}\,.$$

Next, we prove

$$Q_\Phi \leq \frac{A_\Phi}{A_\Phi - 1}\,. \tag{7}$$

If $\Phi \notin \nabla_2$, then $A_\Phi = 1$ or $B_\Psi = \infty$. The result is obvious.

If $\Phi \in \nabla_2$, then $A_\Phi > 1$. For every $x \in L^\Phi[0,\infty)$ which satisfies $\|x\|_\Phi = 1$, and for any $k \in [k_x^*, k_x^{**}]$, we have

$$1 = \|x\|_\Phi = \frac{1}{k}[1 + \rho_\Phi(kx)].$$

For any $0 < \varepsilon < 1 < k$, we have

$$1 = \|x\|_\Phi = \inf_{t>0} \frac{1}{t}[1 + \rho_\Phi(tx)] \leq \frac{1}{k-\varepsilon}[1 + \rho_\Phi((k-\varepsilon)x)]. \tag{8}$$

By the definition of k_x^{**} and $k-\varepsilon < k_x^{**}$, we have

$$\begin{aligned} 1+\rho_\Phi[(k-\varepsilon)x] &\geq \rho_\Psi\{\phi[(k-\varepsilon)x]\} + \rho_\Phi[(k-\varepsilon)x] \\ &= \int_{[0,\infty)} (k-\varepsilon)x(t)\phi[(k-\varepsilon)x(t)]\,dt \\ &\geq A_\Phi \rho_\Phi((k-\varepsilon)x(t)). \end{aligned} \tag{9}$$

Therefore by (8) and (9), we have

$$1 \geq (A_\Phi - 1)\rho_\Phi((k-\varepsilon)x(t)) \geq (A_\Phi - 1)(k-\varepsilon-1))$$

or

$$k-\varepsilon \leq \frac{A_\Phi}{A_\Phi - 1}\,.$$

Since ε is arbitrary, we have

$$k \leq \frac{A_\Phi}{A_\Phi - 1}\,.$$

This implies (7) since x and k are arbitrary. □

Corollary 2.1. (i) *If* $\Phi \in \nabla_2$, *then* $Q_\Phi < \infty$; (ii) *If* $\Phi \in \triangle_2$, *then* $q_\Phi > 1$.

For $0 \neq x \in L^\Phi[0,1]$, we still denote

$$\begin{aligned} k_x^* &= \inf\{k>0 : \rho_\Psi[\phi(kx)] \geq 1\}, \\ k_x^{**} &= \sup\{k>0 : \rho_\Psi[\phi(kx)] \leq 1\}, \\ Q_\Phi &= \sup_{\|x\|_\Phi = 1} k_x^{**} = \sup_{\|x\|_\Phi=1} \left\{ k>0 : \|x\|_\Phi = \frac{1}{k}(1+\rho_\Phi(kx)) \right\}, \\ q_\Phi &= \inf_{\|x\|_\Phi = 1} k_x^{*} = \inf_{\|x\|_\Phi=1} \left\{ k>0 : \|x\|_\Phi = \frac{1}{k}(1+\rho_\Phi(kx)) \right\}. \end{aligned}$$

Let $\varepsilon_0 = \min\{\frac{1}{2\phi(1)}, 1\}$. Denote

$$\begin{aligned} A_\Phi^* &= \inf\left\{ \frac{t\phi(t)}{\Phi(t)} : t \in [\varepsilon_0, \infty) \right\}, \\ B_\Phi^* &= \sup\left\{ \frac{t\phi(t)}{\Phi(t)} : t \in [\varepsilon_0, \infty) \right\}. \end{aligned}$$

Obviously, $\varepsilon_0\phi(\varepsilon_0) \leq \frac{\phi(\varepsilon_0)}{2\phi(1)} \leq \frac{1}{2}$.

Theorem 2.2. *If Φ, Ψ is a pair of complementary $\mathcal{N}$-functions, then*

$$\frac{B_\Phi^* - \varepsilon_0\phi(\varepsilon_0)}{B_\Phi^* - 1} \le q_\Phi \le Q_\Phi \le \frac{A_\Phi^* + A_\Phi^*\Phi(\varepsilon_0)}{A_\Phi^* - 1}\,.$$

Proof: Firstly, we prove $q_\Phi \ge \frac{B_\Phi^* - \varepsilon_0\phi(\varepsilon_0)}{B_\Phi^* - 1}$. If $\Phi \notin \triangle_2(\infty)$, then $B_\Phi^* = \infty$, and the result is clear. If $\Phi \in \triangle_2(\infty)$, then $B_\Phi^* < \infty$. By the proof of Theorem 2.1, for $x \in L^\Phi[0,1]$ with $\|x\|_\Phi = 1$, we have $\rho_\Psi(\phi(k_x^* x)) \ge 1$. So

$$\begin{aligned} 1 + \rho_\Phi(k_x^* x) &\le \rho_\Psi(\phi(k_x^* x)) + \rho_\Phi(k_x^* x) \\ &= \int_{[0,1]} k_x^*|x(t)|\phi(k_x^*|x(t)|)\,dt \\ &\le \int_{G_1=\{t:k_x^*|x(t)|<\varepsilon_0\}} \varepsilon_0\phi(\varepsilon_0)\,dt + \int_{G\setminus G_1} k_x^*|x(t)|\phi(k_x^*|x(t)|)\,dt \\ &< \varepsilon_0\phi(\varepsilon_0) + B_\Phi^*\rho_\Phi(k_x^* x). \end{aligned}$$

Therefore

$$1 - \varepsilon_0\phi(\varepsilon_0) \le (B_\Phi^* - 1)\rho_\Phi(k_x^* x).$$

Noting that $\rho_\Phi(k_x^* x) = k_x^* - 1$, we have

$$\frac{1 - \varepsilon_0\phi(\varepsilon_0)}{B_\Phi^* - 1} \le k_x^* - 1,$$

i.e.

$$k_x^* \ge \frac{B_\Phi^* - \varepsilon_0\phi(\varepsilon_0)}{B_\Phi^* - 1}\,.$$

Since x is arbitrary,

$$q_\Phi \ge \frac{B_\Phi^* - \varepsilon_0\phi(\varepsilon_0)}{B_\Phi^* - 1}\,.$$

Next we prove $Q_\Phi \le \frac{A_\Phi^*(1+\Phi(\varepsilon_0))}{A_\Phi^* - 1}$. If $\Phi \notin \nabla_2(\infty)$, the result is obvious. If $\Phi \in \nabla_2(\infty)$, then $\forall x \in S(L^\Phi[0,1])$, $\forall k \in [k_x^*, k_x^{**}]$ and $0 < \varepsilon < 1$, we get

$$\begin{aligned} 1 + \rho_\Phi[(k-\varepsilon)x] &\ge \rho_\Psi\{\phi[(k-\varepsilon)|x|]\} + \rho_\Phi[(k-\varepsilon)x] \\ &= \int_{[0,1]} (k-\varepsilon)|x(t)|\phi[(k-\varepsilon)|x(t)|]\,dt \\ &\ge \int_{\{t\in[0,1]:(k-\varepsilon)|x(t)|\ge\varepsilon_0\}} (k-\varepsilon)|x(t)|\phi[(k-\varepsilon)|x(t)|]\,dt \\ &\ge A_\Phi^* \int_{\{(k-\varepsilon)|x(t)|\ge\varepsilon_0\}} \Phi((k-\varepsilon)|x(t)|)\,dt \\ &= A_\Phi^*\{\rho_\Phi[(k-\varepsilon)x(t)] - \int_{\{t\in[0,1]:(k-\varepsilon)|x(t)|<\varepsilon_0\}} \Phi((k-\varepsilon)x(t))\,dt\} \\ &\ge A_\Phi^*\{\rho_\Phi[(k-\varepsilon)x(t)] - \Phi(\varepsilon_0)\}. \end{aligned}$$

So

$$1+A_\Phi^*\Phi(\varepsilon_0)\geq (A_\Phi^*-1)\rho((k-\varepsilon)x(t))\geq (A_\Phi^*-1)(k-\varepsilon-1),$$

i.e.

$$k\leq \frac{A_\Phi^*[1+\Phi(\varepsilon_0)]}{A_\Phi^*-1}+\varepsilon.$$

Therefore,

$$k\leq \frac{A_\Phi^*[1+\Phi(\varepsilon_0)]}{A_\Phi^*-1}.$$

Since $x\in S(L^\Phi[0,1])$ is arbitrary,

$$Q_\Phi\leq \frac{A_\Phi^*(1+\Phi(\varepsilon_0))}{A_\Phi^*-1}.$$

□

Corollary 2.2 (S.T. Chen [1, p. 21])**.**
(i) *If* $\Phi\in\triangle_2(\infty)$, *then* $q_\Phi>1$.
(ii) *If* $\Phi\in\nabla_2(\infty)$, *then* $Q_\Phi<\infty$.

From the proof of Theorem 2.2, we know Theorem 2.2 is true for any $0<\varepsilon<\varepsilon_0$. Letting ε to tend to 0, we get

Corollary 2.3. *Let* Φ,Ψ *be a pair of complementary* $\mathcal{N}$*-functions. Then*

$$A_\Psi=\frac{B_\Phi}{B_\Phi-1}\leq q_\Phi\leq Q_\Phi\leq\frac{A_\Phi}{A_\Phi-1}=B_\Psi, \tag{10}$$

where A_Φ, B_Φ, A_Ψ *and* B_Ψ *are defined by* (1).

Example 1. For the $\mathcal{N}$-function $\Phi(u)=|u|^p$, which generates $L^p[0,\infty)$, we have $A_\Phi=B_\Phi=p$. By Theorem 2.1 and Corollary 2.3, we have $q_\Phi=Q_\Phi=\frac{p}{p-1}$.

Example 2. For the $\mathcal{N}$-function $\Phi(u)=e^{|u|}-|u|-1$, we have

$$1\leq q_\Phi\leq Q_\Phi\leq 2. \tag{11}$$

Indeed, $F_\Phi(t)=\frac{t(e^t-1)}{e^t-t-1}$ is increasing in $(0,+\infty)$. So $A_\Phi=\lim\limits_{t\to 0^+}F_\Phi(t)=2$ and $B_\Phi=\lim\limits_{t\to+\infty}F_\Phi(t)=\infty$. Therefore (11) follows from Theorem 2.1 and Corollary 2.3.

Acknowledgment. The author would like to thank the referee for reading this paper carefully and for giving some suggestions.

References

[1] Chen S.T., *Geometry of Orlicz spaces*, Dissertationes Mathematicae, Warszawa, 1996.

[2] Krasnosel'skii M.A., Rutickii Ya.B., *Convex Functions and Orlicz Space*, Noordhoff Ltd., Groningen, 1961.

[3] Lindenstrauss J., Tzafriri L., *Classical Banach Spaces*, I and II, Springer, Berlin, 1977 and 1979.

[4] Maligranda L., *Indices and interpolation*, Dissertationes Math. (Rozprawy Mat.) **234** (1985).

[5] Orlicz W., *Linear Functional Analysis*, World Scientific Publishing, Singapore, 1992.

[6] Rao M.M., Ren Z.D., *Theory of Orlicz Spaces*, Marcel Dekker, New York, 1991.

[7] Rao M.M., Ren Z.D., *Applications of Orlicz Spaces*, Marcel Dekker, New York, 2002.

[8] Simonenko I.B., *Interpolation and extrapolation of linear operators in Orlicz spaces*, Dokl. Akad. Nauk SSSR **151** (1963), 1288–1291 (Russian).

[9] Simonenko I.B., *Interpolation and extrapolation of linear operators in Orlicz spaces*, Mat. Sb. (N.S.) **63** (105) (1964), 536–553 (Russian).

[10] Wu C.X., Zhao S.Z., Chen J.O., *Calculation of Orlicz norm and rotundity of Orlicz spaces*, J. Harbin Inst. Techn. **10** (1973), no. 2, 1–12 (Chinese).

[11] Yan Y., *On a pair of geometric parameters of Orlicz norm*, Comment. Math. Prace Mat. **41** (2001), 257–263.

[12] Yan Y., *Some results on packing in Orlicz sequence spaces*, Studia Math. **147** (1) (2001), 73–88.

[13] Yan Y.Q., *Packing constants, weakly convergent sequence coefficients and Riesz angles in Orlicz sequence spaces*, Ph.D. Dissertation, Suzhou Univ., Suzhou, P. R. China, 2002.

Department of Mathematics, Suzhou University, Suzhou 215006, P.R. China

(*Received* April 9, 2002, *revised* November 11, 2002)